

Open camera or QR reader and
scan code to access this article
and other resources online.



SPECIAL ARTICLE

Social Determinants of Health and Health Equity in the Diagnosis and Management of Pediatric Mild Traumatic Brain Injury: A Content Analysis of Research Underlying Clinical Guidelines

Nathan E. Cook,^{1-3,*} Alicia Kissinger-Knox,^{1,3} Ila A. Iverson,⁴ Brian C. Liu,^{2,3} Charles E. Gaudet,¹⁻³ Marc A. Norman,⁵ and Grant L. Iverson^{1-3,6}

Abstract

We conducted a content analysis of the literature underlying the Centers for Disease Control and Prevention (CDC) Guideline on the Diagnosis and Management of Mild Traumatic Brain Injury Among Children (i.e., the “Guideline”) to determine the extent to which social determinants of health (SDoH) were examined or addressed. The systematic review forming the basis for the Guideline included 37 studies addressing diagnosis, prognosis, and treatment/rehabilitation. We examined those studies to identify SDoH domains derived from the U.S. Department of Health and Human Services’ Healthy People 2020 and 2030 websites. No study explicitly mentioned “social determinants of health,” by name, and few studies addressed SDoH domains as a primary focus (ranging from 0% to 27% of studies across SDoH domains). The most frequently represented SDoH domains, described in an inferential or a descriptive manner, were Education Access and Quality (29.7% of studies), Social and Community Context (27.0% of studies), and Economic Stability (21.6% of studies). Health Care Access (13.5% of studies) was less well represented and no studies (0%) examined Neighborhood and Built Environment. In terms of the CDC clinical questions, SDoH were only examined as predictors of outcome (prognosis) and no studies examined SDoH in relation to diagnosis or treatment/rehabilitation. The Guideline includes some commentary on health literacy and socioeconomic status. Overall, social determinants of health are largely unrepresented as important or meaningful variables influencing the Guideline on the Diagnosis and Management of Mild Traumatic Brain Injury Among Children, or in the studies that informed the Guideline.

Keywords: brain injury; concussion; pediatric; socioeconomic status

¹Department of Physical Medicine and Rehabilitation, Harvard Medical School, Boston, Massachusetts, USA.

²MassGeneral Hospital for Children Sports Concussion Program, Waltham, Massachusetts, USA.

³Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Charlestown, Massachusetts, USA.

⁴Department of Psychology, University of British Columbia, Vancouver, British Columbia, Canada.

⁵Department of Psychiatry, UC San Diego School of Medicine, San Diego, California, USA.

⁶Department of Physical Medicine and Rehabilitation, Shoen Adams Research Institute at Spaulding Rehabilitation, Charlestown, Massachusetts, USA.

*Address correspondence to: Nathan E. Cook, PhD, Sports Concussion Program, Mass General Waltham, 52 2nd Avenue, Suite 1150, Waltham, MA 02451, USA E-mail: necook@mgh.harvard.edu

© Nathan E. Cook et al., 2023; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons License (CC-BY) (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited.

Introduction

Social and economic circumstances significantly influence children's health and well-being.^{1,2} Social determinants of health (SDoH) refer to the environments and conditions in which we are born, raised, educated, live, and work that influence individual and group differences in health status and outcomes.³ Several SDoH domains have been identified including, Economic Stability (e.g., poverty, ability to afford healthcare, and housing instability), Education Access and Quality (e.g., language and literacy, high school graduation, and enrollment in higher education), Health Care Access and Quality (e.g., access to healthcare including primary care, health literacy, and health and dental insurance), Neighborhood and Built Environment (e.g., environmental conditions such as safe air and water, biking and walking accessibility, and crime and violence), and Social and Community Context (e.g., experiences of discrimination or racism, parental mental health, and positive versus negative relationships within the family and community).^{4,5} Social determinants are causal factors in health inequity.

Health equity means that all people should have fair and equal opportunity to enjoy their full health potential⁶ and health inequities are systematic differences in health status between groups, with significant social and economic costs both to individuals and societies.⁷ Sociocultural and demographic factors associated with health inequity (that partially overlap with and are associated with SDoH in direct and indirect ways) include race, ethnicity, language, culture, and socioeconomic status (SES). For example, disparities in pediatric healthcare access and health outcomes have been reported in association with race and ethnicity,⁸⁻¹⁰ SES,¹¹ English language proficiency,¹² and acculturation.^{13,14}

There are important associations between SDoH, health inequity, and pediatric injury. For example, social gradients have been found with regard to causes of medically treated adolescent injuries, such that higher income is associated with greater likelihood of sport-related injury whereas poverty is associated with greater likelihood of injuries due to fighting.¹⁵ Moreover, children from low-income households, children from families that receive income assistance, and children born to teenage mothers are at an elevated risk for burn injuries.¹⁶

A pediatric injury that is a major source of concern for parents, healthcare providers, school personnel, and legislators is concussion, or mild traumatic brain injury (mTBI).¹⁷⁻²³ Approximately 1 in 100 school-aged children in the United States sustain a mTBI each year.²⁴ Emerging research findings have begun to document the clear and expected relevance of SDoH with regard to pediatric mTBI. For example, there is accumulating evidence of racial, ethnic, and socioeconomic disparities in accessing concussion care,²⁵⁻²⁷ recovery time, and supports received upon returning to school following the in-

jury,^{28,29} as well as health literacy (i.e., general concussion knowledge, familiarity with concussion laws, and awareness of concussion symptoms).³⁰⁻³²

Given the significance of concussion and mTBI as a public health concern, the Centers for Disease Control and Prevention (CDC) undertook a massive and pioneering effort to develop the first clinical management guideline in the United States for pediatric mTBI, which was published as the Centers for Disease Control and Prevention Guideline on the Diagnosis and Management of Mild Traumatic Brain Injury Among Children (i.e., the "Guideline").³³ The Guideline includes 19 sets of recommendations regarding the diagnosis, prognosis, and management/treatment of pediatric mTBI. The recommendations in the Guideline were developed based on clinical science identified in a major systematic review.³⁴

The systematic review was organized around six clinical questions related to: 1) diagnosing mTBI; 2) accuracy of routine head imaging; 3) features associated with increased risk of intracranial injury/abnormal head imaging; 4) factors associated with worse outcome in the first year following mTBI; 5) factors associated with worse outcome beyond 1 year following mTBI; and 6) treatments associated with improved mTBI outcome. The systematic review screened 15,150 articles, a total of 2984 full-text articles were reviewed, and 75 articles were ultimately included in the synthesis across the clinical questions (some articles were reviewed across multiple clinical questions). The CDC established a multi-disciplinary workgroup that drafted recommendations relevant to each of the six clinical questions based on the studies identified in the systematic review, along with related evidence, scientific principles, and expert inference.³³ Authors from multiple disciplines including primary care pediatrics, athletic training, physical therapy, and sports medicine have highlighted the broad applicability and importance of the Guideline.³⁵⁻³⁸

The Guideline does not reference "social determinants" or "health equity" by name. However, the Guideline mentions and discusses aspects of SDoH. For instance, the Guideline notes the importance of health literacy and social support as means of promoting better patient outcomes. In addition, family and social stressors (Recommendation 8B), race, ethnicity, and SES (Recommendation 9B) are highlighted as potentially associated with worse outcome following injury and thus healthcare professionals are encouraged to monitor youth with these and other risk factors more closely (Recommendation 11A).

The prevailing Guideline for the diagnosis and management of mTBI in children represents an extraordinary synthesis of a broad and diverse literature leading to specific and practical recommendations for healthcare providers. Leveraging that multi-year effort from multi-disciplinary experts, the purpose of this paper was to

carefully review the studies identified in the systematic review³⁴ and relied upon to support the practice recommendations in the Guideline,³³ and to extract information pertaining to SDoH and health equity. More specifically, we conducted a content analysis to determine the extent to which the clinical science underlying the Guideline for diagnosing and managing pediatric mTBI has examined or discussed SDoH or health equity. Further, we sought to identify critical knowledge gaps to encourage and suggest future directions to enhance the incorporation of SDoH in pediatric mTBI research and clinical practice.

Methods

Selection of studies

We selected studies underlying the CDC Guideline,³³ which were identified through a systematic review.³⁴ We collected the articles identified for four of the six clinical questions from the systematic review (Questions 1, 4, 5, and 6 on the topics of diagnosis, short-term prognosis [within 1 year of injury], longer-term prognosis [beyond 1 year post-injury], and treatment/rehabilitation, respectively). Our focus was on diagnosis and management; thus, we did not examine the studies relevant to the two clinical questions that addressed neuroimaging. A total of 38 studies were identified; one study was excluded because it was a conference presentation, resulting in a final sample of 37 published studies.^{39–75}

Data extraction

We developed a coding sheet to identify SDoH and associated subcategories derived from both the Healthy People 2020⁵ and Healthy People 2030⁴ websites. Each of these websites lists the same five SDoH domains: Economic Stability, Education Assess and Quality, Health Care Access and Quality, Neighborhood and Built Environment, and Social and Community Context. Healthy People 2020 provided a bulleted list of these five domains, with several key issues and underlying factors within each domain. For example, a key issue listed under Economic Stability is “poverty.” We included each of the key issues and underlying factors as separate variables to code. We then cross-referenced this list with the information on Healthy People 2030, which provides a separate webpage for each SDoH domain, with an “Overview and Objectives” section that presents brief text descriptions. Three authors (NEC, AKK, BCL) read each text description and extracted any additional key issues or underlying factors that were not included in the Healthy People 2020 bulleted list. For example, in the Health Care Access and Quality domain, health and/or dental insurance was listed in Healthy People 2030, but it was not on the Healthy People 2020 list; therefore, it was added to the coding sheet.

The coding sheet also included summary information about the study sample including the sample size, average age, age range, as well as the gender, racial, and ethnic composition of the sample (coders collected the exact sample details or identified that the information was “Not Reported” by the study authors). The coding sheet also included study design, whether studies included exclusionary criteria based on demographic, sociocultural, or health factors, and whether studies mentioned future directions/research needs regarding social determinants or health equity. Lastly, we added summary variables to characterize whether, or the extent to which, each study analyzed or provided information about the following five key health equity variables: race, ethnicity, culture/acculturation, SES, and language. Coders could indicate whether the study: 1) provided no mention of the variable; 2) included the variable as a demographic category only; or 3) examined the variable in depth (e.g., the health equity variable was a primary variable of interest in the study or outcome/prognostic results were stratified and reported across levels of the variable). The study coding sheet is included in the online supplemental materials.

After the coding sheet was developed, four authors (NEC, AKK, BCL, IAI) reviewed the 37 articles. Five studies were assigned to all four raters as training articles and the raters met as a group to discuss and calibrate ratings. The remaining 32 articles were assigned to two raters each. Authors performed a content analysis independently by extracting details for each study and completing the coding sheet described above. Discrepancies were resolved by discussion. Lastly, two authors (NEC and AKK) independently determined whether articles addressed SDoH by examining the variables in an inferential or intentional way (e.g., the SDoH variable represented a primary focus or emphasis of the study, or the SDoH variable served as a primary predictor of outcome), or in a descriptive or demographic way (e.g., the SDoH variable was summarized as a demographic variable only, mentioned in the discussion section as an area of future study, or utilized as a design feature of the study such as might pertain to recruitment, inclusion/exclusion criteria).

Data analysis and synthesis

The percentages of studies that were identified as including each of the five SDoH domains and each of the five key health equity variables were calculated and summarized descriptively. No statistical significance testing or quantitative synthesis/meta-analytic techniques were used.

Results

The 37 studies included a total of 15,887 participants (median=150 participants; range 30-3091). The mean sample age was reported in 14 studies and the average

sample age was 12.9 years (standard deviation=3.2). Samples included individuals ranging in age from 0 to 22 years, although the median age for the lower bound of age ranges was 6 years and the median upper bound was 16 years. Gender was reported in nearly all studies (k=34; 91.9%). Racial composition of the samples was only reported in about one-third of studies (k=11; 29.7%). Seven of these 11 studies and one additional study that did not report racial composition reported ethnic composition of the samples (k=8; 21.6%). Study samples were mostly recruited from hospitals/emergency departments (k=30; 81.1%), with very few studies recruiting from high school sports (k=3; 8.1%) or specialty concussion clinics (k=3; 8.1%). One study (2.7%) reported recruiting broadly from a healthcare sys-

tem that included the emergency department, hospital admissions, and outpatient clinics. No study explicitly recruited from pediatric primary care. Study designs included prospective cohort (k=26; 70.3%), retrospective cohort (k=4; 10.8%), retrospective case-control (k=3; 8.1%), and randomized controlled trials (k=2; 5.4%), with one cross-sectional cohort (2.7%) and one cross-sectional case-control (2.7%).

Social determinants of health

No study explicitly mentioned “social determinants of health,” by name. The content analysis results related to SDoH domains among the 37 included studies are summarized in Table 1 and Figure 1. The most common SDoH domains addressed in an inferential or intentional

Table 1. Summary of Content Analysis Results Regarding Inclusion of Social Determinants of Health


First author (year)	CDC question(s)	Social determinants of health (SDoH)					
		Economic stability	Education access and quality	Health care access and quality	Neighborhood and built environment	Social and community context	
Agrawal (2005)	4	Outcome <1 year	–	Descriptive	–	–	–
Babikian (2011)	4	Outcome <1 year	–	Descriptive	Descriptive	–	–
Babikian (2013)	5	Outcome ≥1 year	–	Inferential	–	–	Inferential
Barlow (2010)	4,5	Outcome <1 year and ≥1 year	–	–	–	–	Inferential
Berger (2002)	1	Diagnosis	–	–	–	–	–
Blume (2012)	4	Outcome <1 year	Descriptive	–	Descriptive	–	–
Bouvier (2012)	4	Outcome <1 year	–	–	–	–	–
Castile (2012)	4	Outcome <1 year	–	–	–	–	–
Chrisman (2013)	4	Outcome <1 year	–	–	–	–	–
Fay (2010)	5	Outcome ≥1 year	–	Descriptive	–	–	–
Gagnon (2004)	1	Diagnosis	–	–	Descriptive	–	–
Geyer (2009)	1	Diagnosis	–	–	–	–	–
Grubenhoff (2010)	1	Diagnosis	–	Descriptive	–	–	–
Hessen (2008)	5	Outcome ≥1 year	Descriptive	Descriptive	–	–	–
Levin (2008)	4,5	Outcome <1 year and ≥1 year	Inferential	–	–	–	–
Lovell (2003)	1	Diagnosis	–	–	–	–	–
Lumba-Brown (2014)	6	Treatment	–	–	–	–	–
Massagli (2004)	5	Outcome ≥1 year	–	–	–	–	–
Max (2013b)	4	Outcome <1 year	Inferential	–	–	–	Inferential
Max (2013a)	5	Outcome ≥1 year	Inferential	–	–	–	Inferential
Moran (2009)	4	Outcome <1 year	–	–	–	–	–
Mucha (2014)	1	Diagnosis	–	–	–	–	–
O’Connor (2012)	4,5	Outcome <1 year and ≥1 year	–	Inferential	–	–	Inferential
Olsson (2013)	4,5	Outcome <1 year and ≥1 year	–	–	–	–	Inferential
Papoustis (2014)	5	Outcome ≥1 year	–	–	–	–	–
Ponsford (1999)	4	Outcome <1 year	–	Descriptive	–	–	Inferential
Ponsford (2001)	6	Treatment	–	Descriptive	–	–	Inferential
Rivara (2011)	4,5	Outcome <1 year and ≥1 year	Inferential	–	Descriptive	–	Inferential
Schatz (2006)	1	Diagnosis	–	Descriptive	–	–	–
Smyth (2014)	4,5	Outcome <1 year and ≥1 year	–	–	–	–	Inferential
Taylor (2015)	5	Outcome ≥1 year	Descriptive	Descriptive	–	–	–
Teasdale (2003)	5	Outcome ≥1 year	–	–	–	–	–
Thomas (2015)	6	Treatment	–	–	–	–	–
van der Veen (2015)	4	Outcome <1 year	–	–	–	–	–
Yeates (1999)	4	Outcome <1 year	–	–	–	–	–
Zonfrillo (2014)	4,5	Outcome <1 year and ≥1 year	Inferential	–	Inferential	–	–
Zuckerman (2012)	4	Outcome <1 year	–	–	–	–	–

–=Not addressed.

Inferential=SDoH variable represented in an inferential or intentional way (e.g., a primary focus or emphasis of the study, a primary predictor of outcome).

Descriptive=SDoH variable represented in a descriptive or demographic way (e.g., summarized as a demographic variable only, mentioned in the discussion section as an area of future study, or utilized as a design feature of the study such as might pertain to recruitment, inclusion/exclusion criteria, etc.).

CDC, Centers for Disease Control and Prevention.



manner were Social and Community Context (27.0% of studies), followed by Economic Stability (13.5%). Very few studies addressed Education Access and Quality (5.4%) or Healthcare Access and Quality in an inferential (2.7%) fashion. No studies examined the SDoH domain Neighborhood and Built Environment, either in an inferential or a descriptive manner. Overall, about two-thirds of the studies (67.5%) either did not address a SDoH domain at all or addressed a SDoH domain in a descriptive or demographic fashion (e.g., as a demographic variable only, mentioned in the discussion section as an area of future study, as a design feature of the study such as might pertain to recruitment, inclusion/exclusion criteria). In terms of the CDC clinical questions, SDoH were only examined as predictors of outcome (prognosis) and no studies examined SDoH in relation to diagnosis or treatment/rehabilitation.

Health equity factors

No study explicitly mentioned “health equity” or similar phrasing. The term “disparities” appeared in three studies, but only one study used the term in relation to health disparities or health equity (one study described “disparities” with regard to differing results from adjusted versus unadjusted statistical analyses and another study noted “gender disparities” with regard to biomechanics in head and neck acceleration). The content analysis results related to health equity factors among the 37 included studies are summarized in Table 2 and Figure 2. Approximately one in five studies (k=8; 21.6%) examined a health equity variable in depth, such as by investigating SES as a direct predictor or modifier of outcome. Another third of the studies (32.4%) mentioned at least one of the health equity factors, but only as a demographic category (e.g., reporting the racial or ethnic composition of the

Table 2. Summary of Content Analysis Results Regarding Health Equity Factors, Exclusionary Criteria, and Future Directions

First author (year)	Health equity factors					Exclusion criteria*	Future directions [^]
	Race	Ethnicity	SES	Culture	Language		
Agrawal (2005)	–	–	–	–	–	Yes	No
Babikian (2011)	Demographic	Demographic	Demographic	–	–	Yes	No
Babikian (2013)	–	–	Demographic	–	–	No	No
Barlow (2010)	–	–	Demographic	–	–	No	No
Berger (2002)	Demographic	–	–	–	–	No	No
Blume (2012)	Demographic	Demographic	Demographic	–	–	No	No
Bouvier (2012)	–	–	–	–	–	No	No
Castile (2012)	–	–	–	–	–	No	No
Chrisman (2013)	–	–	–	–	–	No	No
Fay (2010)	In depth	–	In depth	–	–	Yes	Yes
Gagnon (2004)	–	–	–	–	–	Yes	No
Geyer (2009)	–	–	–	–	–	No	No
Grubenhoff (2010)	–	–	–	–	–	Yes	No
Hessen (2008)	–	–	–	–	–	No	No
Levin (2008)	–	–	In depth	–	–	Yes	No
Lovell (2003)	–	–	–	–	–	Yes	No
Lumba-Brown (2014)	–	–	–	–	–	Yes	No
Massagli (2004)	Demographic	–	–	–	–	Yes	No
Max (2013)	Demographic	Demographic	Demographic	–	–	Yes	Yes
Max (2013)	In depth	In depth	In depth	–	–	Yes	No
Moran (2009)	Demographic	–	Demographic	–	–	Yes	No
Mucha (2014)	–	–	–	–	–	Yes	No
O'Connor (2012)	Demographic	Demographic	In depth	–	–	No	No
Olsson (2013)	–	–	–	–	–	Yes	No
Papoustis (2014)	–	–	In depth	–	–	Yes	No
Ponsford (1999)	–	–	Demographic	–	–	Yes	No
Ponsford (2001)	–	–	Demographic	–	–	Yes	Yes
Rivara (2011)	Demographic	Demographic	In depth	–	–	No	No
Schatz (2006)	–	–	–	–	–	No	No
Smyth (2014)	–	Demographic	–	–	–	No	Yes
Taylor (2015)	Demographic	–	In depth	–	–	Yes	No
Teasdale (2003)	–	–	–	–	–	No	No
Thomas (2015)	–	–	–	–	–	Yes	No
van der Veen (2015)	–	–	–	–	–	No	No
Yeates (1999)	Demographic	–	Demographic	–	–	Yes	Yes
Zonfrillo (2014)	In depth	In depth	In depth	–	–	No	Yes
Zuckerman (2012)	–	–	–	–	–	Yes	No

*Studies included exclusionary criteria based on demographic, sociocultural, or health factors.

[^]Studies mentioned future directions/research needs regarding social determinants or health equity.

SES, socioeconomic status.

sample). The most commonly examined health equity factor was SES, with roughly half the studies (45.9%) including SES as either a demographic category or examining it in depth. No studies examined or reported on culture/ac-culturation or language.


Demographic, sociocultural, or health factors as exclusionary criteria

Over half of the studies ($k = 20$; 54.1%) excluded participants based on demographic, sociocultural, or health factors, namely language proficiency and the presence of pre-injury health conditions/disabilities. Many studies excluded children with limited English proficiency,^{51,62–65,69,71,75} or if a guardian/caregiver was unable to complete the research consent process in English.^{65,69,71} One study included English speaking children of Spanish speaking parents.⁵¹ Many studies excluded children with pre-injury health conditions or disabilities, such as a history of a developmental delay, disability, or disorder,^{56,63,76,77} including autism spectrum disorder,^{57,58,75,78} intellectual disability,^{57–59,69,71,73,79} learn-

ing disabilities, including attention-deficit/hyperactivity disorder (ADHD),^{49,54,71,75} enrollment in special education,^{49,75} or diagnosed psychiatric or psychological disorder⁶⁰ including schizophrenia,^{57,58,78} severe psychiatric history including hospitalization,^{59,69,73,75,79} or behavioral problems.⁴⁹ No studies provided rationale or explanation for such exclusionary criteria, nor described such exclusions as a limitation of their study, such that the generalizability of the reported results might be limited. Only one study emphasized the importance of assessing sociocultural factors, such as a youth's pre-injury family background.⁴¹

Social determinants or health equity as future clinical and research considerations

No study explicitly mentioned social determinants of health, by name, as an important framework to guide future mTBI research. One study (2.7%) referenced the need for future research to further explore health disparities in relation to outcome following brain injury, such as



disparities in care, adherence to treatment and rehabilitation plans, and limitations in social support among disadvantaged children and families, though the authors specifically mention disparities in outcome following severe TBI, as opposed to mTBI.⁷⁴ These authors further implore healthcare providers to “ensure multidimensional support and resources” and attend to such factors as socioeconomic disadvantage when providing clinical care for youth with TBI, specifically referencing those who are at socioeconomic disadvantage, including families with lower parental education, lower SES, and Medicaid insurance.⁷⁴ A few other studies (k=5; 13.5%) mentioned or referred to SDoH or health equity indirectly. For example, one study recommended that future research examine family functioning and parental adjustment in relation to outcome and recovery following pediatric mTBI.⁷³ Another study referred generally to “preinjury life stressors” and “a history of previous stressful life events” as important to consider in both

clinical and research considerations regarding pediatric mTBI.⁶⁸ One study implied the importance of health literacy by recommending that information and suggested coping strategies be provided to children and their parents following pediatric mTBI.⁶⁵ Three studies highlighted factors such as disability status (i.e., neurodevelopmental disorders)^{73,79} and SES⁵⁷ as important variables to address in terms of sample inclusion and matching criteria.

Discussion

This content analysis found that SDoH and health equity have been significantly underrepresented in the clinical research literature underlying the CDC Guideline on the Diagnosis and Management of Mild Traumatic Brain Injury Among Children. Studies underlying the Guideline predominantly addressed only a subset of SDoH domains, most commonly Social and Community Context (mainly examining positive vs. negative

relationships at home, four studies examining parental mental health, and two studies examining parental incarceration) and Education Access and Quality (e.g., parental education level, child's pre-injury learning disability status, and child's school functioning). The remaining SDoH domains are of importance to pediatric mTBI management but have very little representation in the literature.

For example, Economic Stability is a major domain of interest given emerging evidence of disparities in accessing concussion care.²⁵⁻²⁷ Health Care Access and Quality is also an important area for future research given initial evidence of health literacy disparities relating to general concussion knowledge, familiarity with concussion laws, and awareness of concussion symptoms.³⁰⁻³² Moreover, no studies examined Neighborhood and Built Environment, which is of significant interest when considering the scientific support for exercise-based concussion rehabilitation, which might include, for some youth, the ability to safely access outdoor spaces to be physically active. Regarding the clinical questions that form the basis for the Guideline, no studies addressing diagnosis (clinical question 1) and only one study addressing treatment (clinical question 6) included SDoH in an inferential or primary fashion. It is important to note that our approach to coding SDoH was liberal and might overrepresent the extent to which social determinants have been examined or woven into this literature.

This content analysis also examined the extent to which health equity and related sociocultural variables were represented or discussed. Health equity was not mentioned explicitly by name, and it was not represented as a primary framework by which researchers designed studies or interpreted results in nearly all of the studies included in the Guideline. Only one study described its findings as revealing disparities in outcomes among disadvantaged youth and encouraged future work to investigate mTBI-related health disparities. Nearly half of the studies did not mention race, ethnicity, language, or culture/acclimation in any context, not even as a demographic characteristic describing the sample composition. Only about one in five studies examined a health equity variable in depth and most frequently this involved SES, such as by including SES as a predictor of outcome. Race and ethnicity were rarely examined in depth (only three studies examined either race and/or ethnicity). No studies underlying the pediatric mTBI Guidelines examined or represented culture/acclimation or language as contributors to health inequity.

Just over half the studies underlying the prevailing Guideline utilized exclusionary criteria based on demographic, sociocultural, or health factors. Specifically, studies excluded youth from participating based on English language proficiency and/or the presence of pre-injury health conditions/disabilities. Moreover, no study

discussed these exclusionary criteria as a limitation nor provided a rationale or explanation for the exclusion. Language-based exclusionary criteria may reflect resource limitations on researchers' part, such as limited funding to adequately translate informed consent documents in multiple languages or the lack of availability of interpreters to ensure appropriate informed consent procedures. Regardless of potential rationales or practical considerations from a research perspective, these exclusionary criteria represent clear limitations to the generalizability of findings, such that youth from underprivileged backgrounds are under-represented in the literature and the extent to which some of the findings from the literature apply to them is unknown.

Lastly, no study explicitly mentioned SDoH, health equity, or similar phrasing as an important framework to guide future mTBI research. Only one study explicitly mentioned the need for future health disparities research regarding clinical care and outcomes, but the authors were referencing severe TBI, not mTBI. A few SDoH and health equity factors were mentioned indirectly, such as authors noting the need for future research to explore family functioning and adjustment following pediatric mTBI, or pre-injury stressful life events in relation to outcome from mTBI.

It is important to note that the articles included in this content analysis, and that form the basis for the Guideline, were drawn from a systematic review of literature published between 1990 and mid-2015.³⁴ It is hoped that more recent literature will increasingly incorporate and examine SDoH and health equity. In turn, the literature base that informs future updated guidelines will more adequately address SDoH and health equity. Moreover, the systematic review that formed the basis for the Guideline did not include a specific question regarding SDoH or health equity, and thus the representation of SDoH and health equity in the larger pediatric mTBI literature may differ, to some degree, from the studies included in this content analysis. Our focus, however, was to review specifically the articles that formed the basis for the development of the current Guideline.

Clinical implications

Clinicians treating children with concussion are encouraged to carefully consider how SDoH might be relevant or important during clinical assessment, patient/family education, and during treatment or rehabilitation, and incorporate relevant information into their care, treatment, and rehabilitation plans. For example, for a youth with persistent symptoms following concussion, clinical science strongly supports the use of exercise-based active rehabilitation.⁸⁰⁻⁸² However, social considerations such as SES and resource availability are crucial to consider. For many youth, access to an in-home treadmill or exercise bike is not feasible or realistic for economic, space,

or other reasons. Additionally, some youth may lack access to fitness facilities with such equipment. Moreover, even suggestions for youth to exercise outdoors may require modification, such as those who live in neighborhoods with limited green space or parks, on or near busy roadways that would not be safe and conducive to jogging, or in neighborhoods where being outside is unsafe in terms of potential for violence.

The reading level and readability scores of patient education materials is an important consideration. Parental literacy and educational attainment may reduce a family's ability to follow recommendations offered by healthcare providers. It is also important to have access to medical interpreters and to have educational materials and supports available in primary languages for the patients served.

For youth experiencing prolonged recovery following mTBI, referral to specialty care is advised.⁸³ However, facilitating referrals to specialty clinics is often a complex process and families may be poorly equipped to navigate the complexities of identifying specialty clinics, seeking referrals, and dealing with health insurance policies. Recent research indicates that specialty concussion clinic patients are more likely to: 1) have private insurance and 2) be White.⁸⁴ Recommendations or referrals for additional forms of treatments to support concussion recovery such as physical therapy or mental health also may present challenges for families with economic constraints or insurance-based limitations. Clinicians are encouraged to proactively support families and facilitate their engagement in care by making direct contact with referral providers to the greatest extent possible.

Youth often require time-limited school accommodations during concussion recovery. However, pursuing and securing accommodations by communicating with school personnel might be more or less challenging for certain caregivers based on factors such as comfort with the school support personnel structure, knowledge of how to contact school nurses or guidance counselors (and comfort in doing so), or trust in the school team that the needs of their children will be prioritized and met.

Implications for future research

There is a need to emphasize SDoH and health equity in pediatric mTBI research and to refine our understanding of how such factors relate to diagnosing, treating, and managing this injury. From a methodological perspective, concussion researchers should make concerted efforts to use qualitative and community-based participatory research designs and methodologies. These methodologies have proven particularly beneficial in terms of elucidating health disparities as well as conceptualizing and designing ways to intervene upon social determinants to reduce disparities among vulnerable groups.^{85,86} Fur-

ther, research teams are encouraged to partner with school-based athletic trainers, healthcare providers, community-based health clinics, and athletic organizations in underserved communities to focus upon the specific and direct needs of underserved youth.

Pediatric mTBI researchers could consider some of the following specific examples for how to address SDoH and health equity in future studies. Focus groups could be conducted to identify potential SDoH needs and barriers to mTBI care. Researchers could directly evaluate the association between SDoH and recovery time from mTBI. Researchers could consider the feasibility of delivering interventions designed to target SDoH. For instance, the readability and reading ease scores of mTBI patient discharge instructions and patient understanding of the instructions might be examined. Studies could examine mTBI treatment and rehabilitation uptake, compliance, and barriers to accessing specialty care. Research designs aimed at informing the nature and scope of interventions designed to improve SDoH, health inequities, and clinical outcomes will be important. Additionally, research focused on uptake and utilization of the Guideline by healthcare providers in regards to SDoH and health equity factors will be important to examine.

Concussion researchers are encouraged to consistently and routinely measure and report complete demographic information about their samples. It seems particularly important for researchers to characterize gender, racial, and ethnic identities of their participants, as well as measure and report additional sociocultural characteristics such as SES and language proficiency. The website for the EQUATOR (Enhancing the QUALity and Transparency Of health Research) network hosts additional templates for reporting guidelines.⁸⁷ *Annals of Behavioral Medicine* recently published instructions for authors reporting on study sample socioeconomic and sociodemographic characteristics that may serve as a guide for mTBI research.⁸⁸ Further, researchers are encouraged to measure and describe their samples in terms of relevant SDoH such as health literacy surrounding mTBI, insurance status, access, and barriers to primary care for mTBIs, as well as access and barriers to specialty concussion care.

As of about 2000, SDoH were rarely screened for in pediatric primary care settings.⁸⁹ Encouragingly, there has been increased emphasis and considerable research interest related to screening for SDoH and related risk factors in the context of primary care,⁹⁰ community health,⁹¹ and pediatric emergency medicine⁹² settings. Still, limitations in the available screening tools and resources have been highlighted along with the need for continued development and refinement of effective and practical tools.⁹³ The American Academy of Pediatrics recommends screening during all patient encounters and provides a website with screening resources.⁹⁴ It is

unknown how frequently, if at all, SDoH are assessed in specialty concussion clinics—and this appears to be an important area of future research. Researchers can examine SDoH and attrition from treatment trials and address disparities accordingly.

Funding agencies and organizations are encouraged to emphasize SDoH in concussion and brain injury research. Funders could consider encouraging researchers to collect and report SDoH in their research studies. For example, the PhenX measures for social determinants of health (SDoH) project, supported by the National Institute on Minority Health and Health Disparities, developed a collection of common data elements to improve the quality and consistency of SDoH data collection.⁹⁵ Additionally, funders could consider developing targeted mechanisms or opportunities regarding the potential role of and means to address SDoH and health inequities for pediatric mTBI diagnosis and clinical care.

Conclusions

Over the past two decades, there has been extraordinary advancement in knowledge about pediatric mTBI. This has included, for example, many studies published examining potential risk or vulnerability factors associated with worse outcome or slower recovery from pediatric mTBI,^{96–101} as well as substantial efforts to develop and validate evidence-supported treatment and rehabilitation approaches for pediatric mTBI, such as exercise-based rehabilitation.^{102–104} The development and publication of the Guideline was a synthesis of this rapidly evolving clinical and scientific knowledge, with the dissemination and implementation goal of improving the treatment and management of pediatric mTBI. With that said, SDoH and health equity have been significantly underrepresented in the clinical research literature, including the literature underlying the Guideline.

A major priority for the field is to better understand the role and relevance of SDoH and health inequities for pediatric mTBI diagnosis and clinical management. Moreover, it is a priority to determine the means and methods that can be used to address SDoH and health inequities and to implement those means and methods, with the goal of promoting health equity. Results of this content analysis revealed that SDoH are underrepresented as important, meaningful, or primary variables of interest directly addressed in the studies that formed the basis for the prevailing Guideline for managing pediatric mTBI and the recommendations for diagnosis, prognosis, and treatment contained therein. SDoH have not represented a primary framework by which researchers designed studies or interpreted results. SDoH are relevant and important for the diagnosis and management of pediatric mTBI. We need more work in the area to inform future updated guidelines.

Transparency, Rigor, and Reproducibility Summary

This review was not pre-registered. This review was designed as a secondary, content analysis of clinical studies that were identified in a large systematic review³⁴ that was conducted as part of the Centers for Disease Control and Prevention's effort to develop the first clinical management guideline in the United States for pediatric mTBI, which was published as the "Centers for Disease Control and Prevention Guideline on the Diagnosis and Management of Mild Traumatic Brain Injury Among Children."³³ The coding sheet for this content analysis was developed by the authors and is provided in the supplemental materials that accompany this manuscript. Two co-authors independently coded every article and discrepancies were resolved by discussion. The results of this content analysis (i.e., the study coding results) are presented in the Tables 1 and 2 and Supplementary Tables S1 and S2. The specific quotes from each manuscript that were coded are also identified and provided in the supplementary materials (i.e., actual quote is reproduced, and the section and pages from the source article are identified).

Authors' Contributions

Nathan Cook: Conceptualization, methodology, validation, formal analysis, investigation, data curation, writing—original draft, writing—review and editing, visualization, supervision, project administration. Alicia Kissinger-Knox: Validation, investigation, data curation, writing—original draft, writing—review and editing. Ila Iverson: Investigation, data curation, writing—review and editing. Brian Liu: Validation, investigation, data curation, writing—review and editing. Charles Gaudet: Validation, investigation, data curation, writing—original draft, writing—review and editing. Marc Norman: Investigation, writing—review and editing. Grant Iverson: Methodology, writing—original draft; writing—review and editing, visualization, supervision.

Funding Information

Dr. Iverson acknowledges unrestricted philanthropic support from the Mooney-Reed Charitable Foundation, Heinz Family Foundation, ImPACT[®] Applications, Inc., National Rugby League, and the Schoen Adams Research Institute at Spaulding Rehabilitation. No entity was involved in study design; the collection, analysis, and interpretation of data; the writing of the report; or the decision to submit the manuscript for publication. No honorarium, grant, or other form of payment was given to anyone to produce the manuscript.

Author Disclosure Statement

Dr. Cook has a clinical and consulting practice in forensic neuropsychology involving individuals who have sustained mild TBIs. Dr. Iverson serves as a

scientific advisor for NanoDx™, Sway Operations, LLC, and Highmark, Inc. He has received research funding from several test publishing companies, including ImPACT Applications, Inc., CNS Vital Signs, and Psychological Assessment Resources (PAR, Inc.). He has a clinical and consulting practice in forensic neuropsychology involving individuals who have sustained mild TBIs.

For the other authors, no competing financial interests exist.

Supplementary Material

Supplementary Table S1
Supplementary Table S2

References

- Tarazi C, Skeer M, Fiscella K, et al. Everything is connected: social determinants of pediatric health and disease. *Pediatr Res* 2016;79(1-2):125–126; doi: 10.1038/pr.2015.220
- Viner RM, Ozer EM, Denny S, et al. Adolescence and the social determinants of health. *Lancet* 2012;379(9826):1641–1652; doi: 10.1016/S0140-6736(12)60149-4
- Centers for Disease Control and Prevention. Social Determinants of Health: Know What Affects Health. Available from: <https://www.cdc.gov/socialdeterminants/index.htm> [Last Accessed; 6/8/2022].
- U.S. Department of Health and Human Services Office of Disease Prevention and Health Promotion. Healthy People 2030. Available from: <https://health.gov/healthypeople/objectives-and-data/social-determinants-of-health> [Last Accessed; March 9, 2021].
- U.S. Department of Health and Human Services OoDPaHP. Healthy People 2020. Washington, DC: U.S.; Available from: <https://www.healthypeople.gov/2020/topics-objectives/topic/social-determinants-of-health> [Last Accessed; March 9, 2021].
- World Health Organization. Health equity. Available from: https://www.who.int/health-topics/health-equity#tab=tab_1 [Last Accessed; 6/8/2022].
- World Health Organization. Health inequities and their causes. Available from: <https://www.who.int/news-room/facts-in-pictures/detail/health-inequities-and-their-causes> [Last Accessed; 6/8/2022].
- Charron-Chénier R, Mueller CW. Racial disparities in medical spending: healthcare expenditures for black and white households (2013–2015). *Race Soc Problems* 2018;10(2):113–133; doi: 10.1007/s12552-018-9226-4
- Flores G; Committee On Pediatric Research. Technical report—racial and ethnic disparities in the health and health care of children. *Pediatrics* 2010;125(4):e979–e1020; doi: 10.1542/peds.2010-0188
- Flores G, Lin H. Trends in racial/ethnic disparities in medical and oral health, access to care, and use of services in US children: has anything changed over the years? *Int J Equity Health* 2013;12:10; doi: 10.1186/1475-9276-12-10
- Lazar M, Davenport L. Barriers to health care access for low income families: a review of literature. *J Community Health Nurs* 2018;35(1):28–37; doi: 10.1080/07370016.2018.1404832
- Flores G, Abreu M, Tomany-Korman SC. Limited English proficiency, primary language at home, and disparities in children's health care: how language barriers are measured matters. *Public Health Rep* 2005;120(4):418–430; doi: 10.1177/003335490512000409
- Lind C, Mirchandani GG, Castrucci BC, et al. The effects of acculturation on healthy lifestyle characteristics among Hispanic fourth-grade children in Texas public schools, 2004–2005. *J Sch Health* 2012;82(4):166–174; doi: 10.1111/j.1746-1561.2011.00682.x
- Hernandez DC, Kimbro RT. The association between acculturation and health insurance coverage for immigrant children from socioeconomically disadvantaged regions of origin. *J Immigr Minor Health* 2013;15(3):453–461; doi: 10.1007/s10903-012-9643-1
- Pickett W, Molcho M, Simpson K, et al. Cross national study of injury and social determinants in adolescents. *Inj Prev* 2005;11(4):213–218; doi: 10.1136/ip.2004.007021
- Padalko A, Gawaziuk J, Chateau D, et al. Social determinants associated with pediatric burn injury: a population-based, case-control study. *J Burn Care Res* 2020;41(4):743–750; doi: 10.1093/jbcr/iraa045
- Chrisman SPD, Whitlock KB, Kroshus E, et al. Parents' perspectives regarding age restrictions for tackling in youth football. *Pediatrics* 2019;143(5); doi: 10.1542/peds.2018-2402
- Conde C. Playing it safe. *Tex Med* 2011;107(9):37-40
- McCadden MD, Cusimano MD. Staying true to Rowan's Law: how changing sport culture can realize the goal of the legislation. *Can J Public Health* 2019;110(2):165–168; doi: 10.17269/s41997-019-00174-8
- Sarmiento K, Donnell Z, Bell E, et al. From the CDC: a qualitative study of middle and high school professionals' experiences and views on concussion: identifying opportunities to support the return to school process. *J Safety Res* 2019;68:223–229; doi: 10.1016/j.jsr.2018.10.010
- Stelzer JW, Esplin NE, Sbeih F, et al. Commentary: The Florida Youth Concussion Law: a survey-based observational study of physician practices and their legislative awareness. *Neurosurgery* 2019;85(3):E592–E598; doi: 10.1093/neuros/nyz217
- Waltzman D, Sarmiento K. What the research says about concussion risk factors and prevention strategies for youth sports: a scoping review of six commonly played sports. *J Safety Res* 2019;68:157–172; doi: 10.1016/j.jsr.2018.11.005
- Fishman M, Taranto E, Perlman M, et al. Attitudes and counseling practices of pediatricians regarding youth sports participation and concussion risks. *J Pediatr* 2017;184:19–25; doi: 10.1016/j.jpeds.2017.01.048
- Cook NE, Iverson GL. Concussion among children in the U.S. general population: incidence and risk factors. *Frontiers in Neurology* 2021;12(773927):1–6; doi: 10.3389/fneur.2021.773927
- Lyons TW, Miller KA, Miller AF, et al. Racial and ethnic differences in emergency department utilization and diagnosis for sports-related head injuries. *Front Neurol* 2019;10:690; doi: 10.3389/fneur.2019.00690
- Copley M, Jimenez N, Kroshus E, et al. Disparities in use of subspecialty concussion care based on ethnicity. *J Racial Ethn Health Disparities* 2020;7(3):571–576; doi: 10.1007/s40615-019-00686-6
- Wallace JS, Mannix RC. Racial disparities in diagnosis of concussion and minor head trauma and mechanism of injury in pediatric patients visiting the emergency department. *J Pediatr* 2021;233:249–254 e1; doi: 10.1016/j.jpeds.2021.01.057
- Snedden TR, Pierpoint LA, Currie DW, et al. Postconcussion academic support in children who attend a primary care provider follow-up visit after presenting to the emergency department. *J Pediatr* 2019;209:168–175; doi: 10.1016/j.jpeds.2019.01.041
- Babcock L, Byczkowski T, Wade SL, et al. Predicting postconcussion syndrome after mild traumatic brain injury in children and adolescents who present to the emergency department. *JAMA Pediatr* 2013;167(2):156–161; doi: 10.1001/jamapediatrics.2013.434
- Wallace J, Covassin T, Moran R. Racial disparities in concussion knowledge and symptom recognition in American adolescent athletes. *J Racial Ethn Health Disparities* 2018;5(1):221–228; doi: 10.1007/s40615-017-0361-1
- Wallace J, Beidler E, Register-Mihalik JK, et al. Evaluating concussion nondisclosure in college athletes using a health disparities framework and appreciation for social determinants of health. *J Athl Train* 2022;57(1):16–24; doi: 10.4085/1062-6050-0054.21
- Sungwon K, Daniel PC, John OS. Youth sport parents' familiarity and perceptions of concussion legislation. *J Concussion* 2022;6:1–8; doi: 10.1177/20597002221075007
- Lumba-Brown A, Yeates KO, Sarmiento K, et al. Centers for Disease Control and Prevention guideline on the diagnosis and management of mild traumatic brain injury among children. *JAMA Pediatr* 2018;172(11):e182853; doi: 10.1001/jamapediatrics.2018.2853
- Lumba-Brown A, Yeates KO, Sarmiento K, et al. Diagnosis and management of mild traumatic brain injury in children: a systematic review. *JAMA Pediatr* 2018;172(11):e182847; doi: 10.1001/jamapediatrics.2018.2847
- Choe MC, Gregory AJ, Haegerich TM. What pediatricians need to know about the CDC guideline on the diagnosis and management of mTBI. *Front Pediatr* 2018;6:249; doi: 10.3389/fped.2018.00249
- Sarmiento K, Waltzman D, Lumba-Brown A, et al. CDC guideline on mild traumatic brain injury in children: important practice takeaways for sports medicine providers. *Clin J Sport Med* 2020;30(6):612–615; doi: 10.1097/JSM.0000000000000704
- Register-Mihalik JK, Sarmiento K, Vander Vegt CB, et al. Considerations for athletic trainers: a review of guidance on mild traumatic brain injury among children from the Centers for Disease Control and Prevention and the National Athletic Trainers' Association. *J Athl Train* 2019;54(1):12–20; doi: 10.4085/1062-6050-451-18

38. Mucha A, DeWitt J, Greenspan AI. The CDC guideline on the diagnosis and management of mild traumatic brain injury among children: what physical therapists need to know. *Phys Ther* 2019;99(10):1278–1280; doi: 10.1093/ptj/pzz085
39. Agrawal D, Gowda NK, Bal CS, et al. Is medial temporal injury responsible for pediatric postconcussion syndrome? A prospective controlled study with single-photon emission computerized tomography. *J Neurosurg* 2005;102(2 Suppl):167–171; doi: 10.3171/jns.2005.102.2.0167
40. Babikian T, Satz P, Zaucha K, et al. The UCLA longitudinal study of neurocognitive outcomes following mild pediatric traumatic brain injury. *J Int Neuropsychol Soc* 2011;17(5):886–895; doi: 10.1017/S1355617711000907
41. Babikian T, McArthur D, Asarnow RF. Predictors of 1-month and 1-year neurocognitive functioning from the UCLA longitudinal mild, uncomplicated, pediatric traumatic brain injury study. *J Int Neuropsychol Soc* 2013;19(2):145–154; doi: 10.1017/S135561771200104X
42. Barlow KM, Crawford S, Stevenson A, et al. Epidemiology of postconcussion syndrome in pediatric mild traumatic brain injury. *Pediatrics* 2010;126(2):e374–e381; doi: 10.1542/peds.2009-0925
43. Berger RP, Pierce MC, Wisniewski SR, et al. Serum S100B concentrations are increased after closed head injury in children: a preliminary study. *J Neurotrauma* 2002;19(11):1405–1409; doi: 10.1089/089771502320914633
44. Blume HK, Vavilala MS, Jaffe KM, et al. Headache after pediatric traumatic brain injury: a cohort study. *Pediatrics* 2012;129(1):e31–e39; doi: 10.1542/peds.2011-1742
45. Bouvier D, Fournier M, Dauphin JB, et al. Serum S100B determination in the management of pediatric mild traumatic brain injury. *Clin Chem* 2012;58(7):1116–1122; doi: 10.1373/clinchem.2011.180828
46. Castile L, Collins CL, McIlvain NM, et al. The epidemiology of new versus recurrent sports concussions among high school athletes, 2005–2010. *Br J Sports Med* 2012;46(8):603–610; doi: 10.1136/bjsports-2011-090115
47. Chrisman SP, Rivara FP, Schiff MA, et al. Risk factors for concussive symptoms 1 week or longer in high school athletes. *Brain Inj* 2013;27(1):1–9; doi: 10.3109/02699052.2012.722251
48. Fay TB, Yeates KO, Taylor HG, et al. Cognitive reserve as a moderator of postconcussive symptoms in children with complicated and uncomplicated mild traumatic brain injury. *J Int Neuropsychol Soc* 2010;16(1):94–105; doi: 10.1017/S1355617709991007
49. Gagnon I, Swaine B, Friedman D, et al. Visuomotor response time in children with a mild traumatic brain injury. *J Head Trauma Rehabil* 2004;19(5):391–404; doi: 10.1097/00001199-200409000-00004
50. Geyer C, Ulrich A, Grafe G, et al. Diagnostic value of S100B and neuron-specific enolase in mild pediatric traumatic brain injury. *J Neurosurg Pediatr* 2009;4(4):339–344; doi: 10.3171/2009.5.PEDS08481
51. Grubenhoff JA, Kirkwood M, Gao D, et al. Evaluation of the standardized assessment of concussion in a pediatric emergency department. *Pediatrics* 2010;126(4):688–695; doi: 10.1542/peds.2009-2804
52. Hessen E, Anderson V, Nestvold K. MMPI-2 profiles 23 years after paediatric mild traumatic brain injury. *Brain Inj* 2008;22(1):39–50; doi: 10.1080/02699050701846179
53. Levin HS, Hanten G, Roberson G, et al. Prediction of cognitive sequelae based on abnormal computed tomography findings in children following mild traumatic brain injury. *J Neurosurg Pediatr* 2008;1(6):461–470; doi: 10.3171/PED/2008/1/6/461
54. Lovell MR, Collins MW, Iverson GL, et al. Recovery from mild concussion in high school athletes. *J Neurosurg* 2003;98(2):296–301; doi: 10.3171/jns.2003.98.2.0296
55. Lumba-Brown A, Harley J, Lucio S, et al. Hypertonic saline as a therapy for pediatric concussive pain: a randomized controlled trial of symptom treatment in the emergency department. *Pediatr Emerg Care* 2014;30(3):139–145; doi: 10.1097/PEC.0000000000000084
56. Massagli TL, Fann JR, Burington BE, et al. Psychiatric illness after mild traumatic brain injury in children. *Arch Phys Med Rehabil* 2004;85(9):1428–1434; doi: 10.1016/j.apmr.2003.12.036
57. Max JE, Schachar RJ, Landis J, et al. Psychiatric disorders in children and adolescents in the first six months after mild traumatic brain injury. *J Neuropsychiatry Clin Neurosci* 2013;25(3):187–197; doi: 10.1176/appi.neuropsych.12010011
58. Max JE, Pardo D, Hanten G, et al. Psychiatric disorders in children and adolescents six-to-twelve months after mild traumatic brain injury. *J Neuropsychiatry Clin Neurosci* 2013;25(4):272–282; doi: 10.1176/appi.neuropsych.12040078
59. Moran LM, Taylor HG, Ganesalingam K, et al. Apolipoprotein E4 as a predictor of outcomes in pediatric mild traumatic brain injury. *J Neurotrauma* 2009;26(9):1489–1495; doi: 10.1089/neu.2008.0767
60. Mucha A, Collins MW, Elbin RJ, et al. A brief Vestibular/Ocular Motor Screening (VOMS) assessment to evaluate concussions: preliminary findings. *Am J Sports Med* 2014;42(10):2479–2486; doi: 10.1177/0363546514543775
61. O'Connor SS, Zatzick DF, Wang J, et al. Association between posttraumatic stress, depression, and functional impairments in adolescents 24 months after traumatic brain injury. *J Trauma Stress* 2012;25(3):264–271; doi: 10.1002/jts.21704
62. Olsson KA, Lloyd OT, Lebrocq RM, et al. Predictors of child post-concussion symptoms at 6 and 18 months following mild traumatic brain injury. *Brain Inj* 2013;27(2):145–157; doi: 10.3109/02699052.2012.729286
63. Papoutsis J, Stargatt R, Catroppa C. Long-term executive functioning outcomes for complicated and uncomplicated mild traumatic brain injury sustained in early childhood. *Dev Neuropsychol* 2014;39(8):638–645; doi: 10.1080/87565641.2014.979926
64. Ponsford J, Willmott C, Rothwell A, et al. Impact of early intervention on outcome after mild traumatic brain injury in children. *Pediatrics* 2001;108(6):1297–1303; doi: 10.1542/peds.108.6.1297
65. Ponsford J, Willmott C, Rothwell A, et al. Cognitive and behavioral outcome following mild traumatic head injury in children. *J Head Trauma Rehabil* 1999;14(4):360–372; doi: 10.1097/00001199-199908000-00005
66. Rivara FP, Koepsell TD, Wang J, et al. Disability 3, 12, and 24 months after traumatic brain injury among children and adolescents. *Pediatrics* 2011;128(5):e1129–1138; doi: 10.1542/peds.2011-0840
67. Schatz P, Pardini JE, Lovell MR, et al. Sensitivity and specificity of the IMPACT Test Battery for concussion in athletes. *Arch Clin Neuropsychol* 2006;21(1):91–99; doi: 10.1016/j.acn.2005.08.001
68. Smyth K, Sandhu SS, Crawford S, et al. The role of serotonin receptor alleles and environmental stressors in the development of post-concussive symptoms after pediatric mild traumatic brain injury. *Dev Med Child Neurol* 2014;56(1):73–77; doi: 10.1111/dmcn.12263
69. Taylor HG, Orchinik LJ, Minich N, et al. Symptoms of persistent behavior problems in children with mild traumatic brain injury. *J Head Trauma Rehabil* 2015;30(5):302–310; doi: 10.1097/HTR.0000000000000106
70. Teasdale TW, Engberg AW. Cognitive dysfunction in young men following head injury in childhood and adolescence: a population study. *J Neurol Neurosurg Psychiatry* 2003;74(7):933–936; doi: 10.1136/jnnp.74.7.933
71. Thomas DG, Apps JN, Hoffmann RG, et al. Benefits of strict rest after acute concussion: a randomized controlled trial. *Pediatrics* 2015;135(2):213; doi: 10.1542/peds.2014-0966
72. van der Veek EM, Oosterhoff M, Vos PE, et al. The juvenile head trauma syndrome: a trauma triggered migraine? *Neuropediatrics* 2015;46(2):116–122; doi: 10.1055/s-0035-1547344
73. Yeates KO, Luria J, Bartkowski H, et al. Postconcussive symptoms in children with mild closed head injuries. *J Head Trauma Rehabil* 1999;14(4):337–350; doi: 10.1097/00001199-199908000-00003
74. Zonfrillo MR, Durbin DR, Koepsell TD, et al. Prevalence of and risk factors for poor functioning after isolated mild traumatic brain injury in children. *J Neurotrauma* 2014;31(8):722–727; doi: 10.1089/neu.2013.3088
75. Zuckerman SL, Solomon GS, Forbes JA, et al. Response to acute concussive injury in soccer players: is gender a modifying factor? *J Neurosurg Pediatr* 2012;10(6):504–510; doi: 10.3171/2012.8.PEDS12139
76. Agrawal D, Gowda NK, Bal CS, et al. Is medial temporal injury responsible for pediatric postconcussion syndrome? A prospective controlled study with single-photon emission computerized tomography. *J Neurosurg Pediatr* 2005;102(2):167–171; doi: 10.3171/ped.2005.102.2.0167
77. Lumba-Brown A, Harley J, Lucio S, et al. Hypertonic saline as a therapy for pediatric concussive pain: a randomized controlled trial of symptom treatment in the emergency department. *Pediatr Emerg Care* 2014;30(3):139–145; doi: 10.1097/PEC.0000000000000084
78. Levin HS, Hanten G, Luerssen TG, et al. Prediction of cognitive sequelae based on abnormal computed tomography findings in children following mild traumatic brain injury. *J Neurosurg Pediatrics* 2008;1(6):461–470; doi: 10.3171/PED/2008/1/6/461
79. Fay TB, Yeates KO, Taylor HG, et al. Cognitive reserve as a moderator of postconcussive symptoms in children with complicated and uncomplicated mild traumatic brain injury. *J Int Neuropsychol Soc* 2010;16(1):94–105; doi: 10.1017/S1355617709991007

80. Hugentobler JA, Quatman-Yates C, Evanson NK, et al. Active rehabilitation after childhood and adolescent mild traumatic brain injury: a narrative review and clinical practice implications. *Curr Phys Med Rehabil Rep* 2019;7(1):15–22; doi: 10.1007/s40141-019-0207-3
81. Quatman-Yates C, Cupp A, Gunsch C, et al. Physical rehabilitation interventions for post-mTBI symptoms lasting greater than 2 weeks: systematic review. *Phys Ther* 2016;96(11):1753–1763; doi: 10.2522/ptj.20150557
82. Vuu S, Barr CJ, Killington M, et al. Physical exercise for people with mild traumatic brain injury: a systematic review of randomized controlled trials. *NeuroRehabilitation* 2022;51(2):185–200; doi: 10.3233/NRE-220044
83. Silverberg ND, Duhaime A-C, Iaccarino MA. Mild traumatic brain injury in 2019–2020. *JAMA* 2020;323(2):177–178; doi: 10.1001/jama.2019.18134
84. Pate J, Cummins I, Mooney J, et al. Socioeconomic and demographic considerations of pediatric concussion recovery. *J Clin Neurosci* 2022;100:94–99; doi: <https://doi.org/10.1016/j.jocn.2022.04.008>
85. Stewart M, Makwarimba E, Barnfather A, et al. Researching reducing health disparities: mixed-methods approaches. *Soc Sci Med* 2008;66(6):1406–1417; doi: 10.1016/j.socscimed.2007.11.021
86. LaPlant MB, Hess DJ. A review of racial/ethnic disparities in pediatric trauma care, treatment, and outcomes. *J Trauma Acute Care Surg* 2019 Mar;86(3):540–550; doi: 10.1097/TA.0000000000002160
87. Equator Network. Enhancing the Quality and Transparency of Health Research. Available from: www.equator-network.org
88. Revenson TA, Zoccola PM. New instructions to authors emphasize open science, transparency, full reporting of sociodemographic characteristics of the sample, and avoidance of piecemeal publication. *Ann Behav Med* 2022;56(5):415–417; doi: 10.1093/abm/kaac013
89. Kogan MD, Schuster MA, Yu SM, et al. Routine assessment of family and community health risks: parent views and what they receive. *Pediatrics* 2004;113(6 Suppl):1934–1943
90. Berman RS, Patel MR, Belamarich PF, et al. Screening for poverty and poverty-related social determinants of health. *Pediatr Rev* 2018;39(5):235–246; doi: 10.1542/pir.2017-0123
91. Herrera CN, Brochier A, Pellicer M, et al. Implementing social determinants of health screening at community health centers: clinician and staff perspectives. *J Prim Care Community Health* 2019;10:2150132719887260; doi: 10.1177/2150132719887260
92. Gottlieb L, Hessler D, Long D, et al. A randomized trial on screening for social determinants of health: the iScreen study. *Pediatrics* 2014;134(6):e1611–1618; doi: 10.1542/peds.2014-1439
93. Morone J. An integrative review of social determinants of health assessment and screening tools used in pediatrics. *J Pediatr Nurs* 2017;37:22–28; doi: 10.1016/j.pedn.2017.08.022
94. American Academy of Pediatrics. Social Determinants of Health Screening Resources. Available from: www.aap.org/en/patient-care/screening-technical-assistance-and-resource-center/screening-resource-library/social-determinants-of-health
95. PhenX Toolkit. Social Determinants of Health Project. Available from: www.phenxtoolkit.org/collections/sdoh
96. Cook NE, Gaudet Iii CE, Van Patten R, et al. Clinical outcome following sport-related concussion among children and adolescents with a history of prior concussion: a systematic review. *J Neurotrauma* 2022;39(17-18):1146–1158; doi: 10.1089/neu.2022.0078
97. Gornall A, Takagi M, Morawakage T, et al. Mental health after paediatric concussion: a systematic review and meta-analysis. *Br J Sports Med* 2021;55(18):1048–1058; doi: 10.1136/bjsports-2020-103548
98. Djukic S, Phillips NL, Lah S. Sleep outcomes in pediatric mild traumatic brain injury: a systematic review and meta-analysis of prevalence and contributing factors. *Brain Inj* 2022;36(12-14):1289–1322; doi: 10.1080/02699052.2022.2140198
99. Chadwick L, Sharma MJ, Madigan S, et al. Classification criteria and rates of persistent postconcussive symptoms in children: a systematic review and meta-analysis. *J Pediatr* 2022;246:131–137 e2; doi: 10.1016/j.jpeds.2022.03.039
100. van Ierssel J, Pennock KF, Sampson M, et al. Which psychosocial factors are associated with return to sport following concussion? A systematic review. *J Sport Health Sci* 2022;11(4):438–449; doi: 10.1016/j.jshs.2022.01.001
101. Goreth MB, Palokas M. Association between premorbid neuropsychological conditions and pediatric mild traumatic brain injury/concussion recovery time and symptom severity: a systematic review. *JBI Database System Rev Implement Rep* 2019;17(7):1464–1493; doi: 10.11124/JBISRIR-2017-004008
102. Leddy JJ, Haider MN, Ellis MJ, et al. Early subthreshold aerobic exercise for sport-related concussion: a randomized clinical trial. *JAMA Pediatr* 2019;173(4):319–325; doi: 10.1001/jamapediatrics.2018.4397
103. Leddy JJ, Master CL, Mannix R, et al. Early targeted heart rate aerobic exercise versus placebo stretching for sport-related concussion in adolescents: a randomised controlled trial. *Lancet Child Adolesc Health* 2021;5(11):792–799; doi: 10.1016/S2352-4642(21)00267-4
104. Chan C, Iverson GL, Purtzki J, et al. Safety of active rehabilitation for persistent symptoms after pediatric sport-related concussion: a randomized controlled trial. *Arch Phys Med Rehabil* 2018;99(2):242–249; doi: 10.1016/j.apmr.2017.09.108