

Short Communication

Datana drexelii (Lepidoptera: Notododontidae) Occurrence and Larval Survival on Highbush Blueberry Cultivars

Alex K. Baranowski,^{1,3,*} Steven R. Alm,² and Evan L. Preisser¹

¹Department of Biological Sciences, University of Rhode Island, Woodward Hall, 9 East Alumni Avenue, Kingston, RI 02881,

²Department of Plant Sciences and Entomology, University of Rhode Island, Kingston, RI 02881, and ³Corresponding author, e-mail: alexbaran74@gmail.com

Subject Editor: Cesar Rodriguez-Saona

Received 2 January 2020; Editorial decision 3 March 2020

Abstract

Plant genotype influences plant suitability to herbivores; domesticated plants selected for properties such as high fruit yield may be particularly vulnerable to herbivory. Cultivated strains of highbush blueberry, *Vaccinium corymbosum* L. can be high-quality hosts for larvae of the gregariously feeding notodontid *Datana drexelii* (Hy. Edwards). We conducted an experiment assessing *D. drexelii* larval survival and pupal weight when fed foliage from five blueberry cultivars: ‘Bluecrop’, ‘Bluetta’, ‘Blueray’, ‘Lateblue’, and ‘Jersey’. We complemented this experimental work with repeated bush-level surveys of a managed blueberry patch for naturally occurring *D. drexelii* larval clusters. Larval survival and pupal weight were significantly higher on ‘Lateblue’ foliage than from the ‘Bluecrop’, ‘Bluetta’, and ‘Jersey’ cultivars. The blueberry patch surveys found more *D. drexelii* larval clusters on ‘Bluehaven’, ‘Collins’, and ‘Darrow’ bushes than on the cultivars ‘Earliblue’ and ‘Jersey’. The low *D. drexelii* occurrence and performance on the ‘Jersey’ cultivar suggests that this variety may be appropriate for areas where this pest is common; conversely, their high occurrence on ‘Bluehaven’, ‘Collins’, and ‘Darrow’ suggests that these cultivars may be particularly vulnerable. Cultivar-level variation in herbivore vulnerability highlights how understanding plant–pest interactions can help manage agricultural species.

Key words: herbivory, preference, performance, defoliator

Herbivore fitness is influenced by host plant phenotype. Although wild plants experience strong selection for herbivore tolerance and/or resistance, domesticated plants are subjected to different pressures. Selection for high fruit yield in domesticated plants, for example, can reduce plant defense against herbivores (Sanchez-Hernandez et al. 2006, Turcotte et al. 2014, Hernandez-Cumplido et al. 2018). Larvae of *Lymantria dispar* L. (Lepidoptera: Erebidae) grow more quickly and have lower mortality when reared on domesticated versus wild-type *Vaccinium corymbosum* L. (Hernandez-Cumplido et al. 2018). Wild-type tomatoes (*Solanum lycopersicum* L.) produce more phenolic compounds than domesticated ones (Sanchez-Hernandez et al. 2006), and growth rate of the moth *Manduca sexta* L. (Lepidoptera: Sphingidae) is negatively correlated with such phenolics (Stamp and Yang 1996, Yang and Stamp 1996).

Vaccinium corymbosum (hereafter ‘blueberry’) is a deciduous ericaceous plant native to North America grown commercially for its fruits. As with other agricultural plants, blueberry has multiple cultivars that have been selected for yield, flavor, or pest/disease resistance (Lobos and Hancock 2015, Clift et al. 2017, Rodriguez-Saona

et al. 2019). Cultivar-related differences in herbivore growth and mortality have been recorded in lepidopteran species such as *Streblote panda* (Hübner, 1820) (Lepidoptera: Lasiocampidae) (Calvo and Molina 2010), and tephritid flies such as *Bactrocera dorsalis* (Hendel), *Ceratitidis capitata* (Wiedemann, 1824) (Follett et al. 2011), and *Rhagoletis mendax* (Curran) (Liburd et al. 1998).

Members of the notodontid genus *Datana* (Walker, 1855) are defoliating pests of agricultural, silvicultural, and horticultural crops (Cutler and Harris 1979, Harris 1983). *Datana drexelii* (Hy. Edwards) (hereafter ‘*Datana*’) is a native defoliating pest of ericaceous plants in the genera *Vaccinium* L. and *Gaylussacia* (Kunth). Females lay clusters of up to 200 eggs on suitable host plants; their gregarious nature, combined with the fact that larvae can reach 6 cm in length, makes it an especially destructive pest (Wagner 2005). While this insect does not directly attack fruit, its defoliation of blueberry bushes reduces the following year’s flowering and subsequent fruit crop (Lyrene 1992, Williamson and Miller 2000). We reared *Datana* larvae on different blueberry cultivars and measured their survival to pupation and pupal weight. In addition, we repeatedly

surveyed a patch planted with multiple blueberry cultivars for naturally occurring clusters of *Datana* larvae. Together, the data reveal substantial differences in *Datana* preference for and occurrence on different blueberry cultivars.

Materials and Methods

Performance Assay

In June 2019, we mated adults from a lab colony of *D. drexelii*, reared on wild-type *V. corymbosum*, in an outdoor emergence cage at the University of Rhode Island's East Farm research facility (Kingston, RI). We collected their eggs and assigned five each to 946-ml polypropylene cups (Pactiv LLC, Lake Forest, IL). Each cup was randomly assigned to one of five blueberry cultivars: 'Bluecrop', 'Blueray', 'Bluetta', 'Jersey', or 'Lateblue'. There were 22–23 replicate cups per cultivar. Eggs in a given cup generally hatched on the same day; in three cups 1 d, and in one cup 2 d, elapsed between the emergence of the first and last hatchling. Host foliage from the appropriate cultivar was added to the cup immediately following emergence of the first hatchling. Larvae received four-leaf sections of foliage from current year's growth (as indicated by soft, green bark); no other leaf position standardization was done. Prior to adding foliage to each container, each piece was dipped in a 2% bleach (=0.1% NaOCl) solution and allowed to air-dry; this measure was taken to decrease the threat posed by pathogenic fungi and bacteria (Trivedy et al. 2011). Foliage was replaced every 3 d or as needed to ensure a constant food supply.

Four days after the last hatchling in a given cup eclosed, we weighed all hatchlings together and counted the number of larvae and unhatched eggs. The total number of hatched larvae was our starting number of larvae for a cup, regardless of how many eggs hatched. We used this data to calculate posthatching survival. Larvae were subsequently counted and weighed together each week; we recorded the date each larva entered the prepupal phase. Prepupae were left in cups until all larvae in a cup reached such a state or died. When all prepupae had either died or become pupae, each pupa was sexed, weighed, and then held in a 6-liter polypropylene bin (Sterilite Corp, Townsend, MA) of moist coconut coir for overwintering.

Occurrence Assay

In summer 2019, we conducted a 6-wk *Datana* survey of an East Farm blueberry patch enclosed in bird-proof netting that did not exclude insects. The patch consisted of 240 bushes arranged in eight rows of 30 bushes. The cultivars represented (numbers of bushes in parentheses) were 'Bluecrop' (25), 'Bluegold' (5), 'Bluehaven' (15), 'Bluejay' (15), 'Blueray' (15), 'Bluetta' (15), 'Chandler' (5) 'Collins' (20), 'Darrow' (20), 'Earliblue' (30), 'Herbert' (15), 'Jersey' (15), 'Lateblue' (15), 'Northland' (15), and 'Reka' (15). Cultivars were arranged in five-bush groups within a given row.

Between 16 July and 26 August, we conducted 15 total censuses (with as many as 9 d and as few as 1 d between censuses) for *Datana* larval clusters. We walked on both sides of each bush and scanned for larval clusters. We spent a minimum of 30 s per bush and longer if necessary and recorded the number of larval clusters on each bush before removing them from the bush. Following the final census, we measured the height and maximum width of each bush.

Statistical Analysis

For the performance assay, mean pupal weight and percent survival (average per cup) to pupation were analyzed using analysis of

covariance (ANCOVA), with 'cultivar' as the main effect and 'hatch date' and, because we were concerned about sex-mediated performance differences, 'number of female pupae per cup' as covariates. We excluded 15 cups in which only a single larva hatched, leaving a total of 98 cups (=replicates). When the ANCOVA revealed a significant main effect, we used Tukey's HSD tests ($\alpha = 0.05$) to differentiate between treatment. We also conducted an overall linear correlation analysis between weight and survival across all cultivars.

For the occurrence assay, we summed the total number of larval clusters counted per bush over the 15 censuses. Prior to analysis, we removed data from two cultivars, 'Bluegold' and 'Chandler', only represented by a single five-bush cluster within the patch; all other cultivars were each represented by between three and six five-bush clusters. During our surveys, we noticed that small (~0.5 m in height) recently planted bushes had virtually no larval clusters regardless of their cultivar. We addressed this bias by excluding bushes below the 10th percentile in height (0.7 m) from the analysis; this excluded 27 bushes from five cultivars but only two of 108 larval clusters. We used GLMM (Poisson distribution with log-link function) to analyze the 203-bush data set. The model was generated with the main effect 'cultivar' and the blocking variables 'row' and 'column' as random effects; 'bush height' was included as a covariate. The GLMM was initially run using both blocking variables; the nonsignificant blocking variable 'row' was then removed and the resulting GLMM rerun. When the GLMM revealed a significant main effect of 'cultivar', we used likelihood-ratio chi-square tests ($\alpha = 0.05$; controlled for type 1 errors due to multiple comparisons) to differentiate between treatments.

All analyses were performed using JMP 9.0.0 (SAS 2010).

Results

Performance Assay

Larvae reared on 'Lateblue' pupated at nearly three times the weight of larvae reared on 'Blueray', 'Bluetta', and 'Jersey' (0.375 g vs 0.127 g, respectively; $F_{4,91} = 3.18$, $P = 0.017$; Fig. 1A). Survival to pupation was also higher on 'Lateblue' than on 'Bluecrop', 'Bluetta', and 'Jersey' (16.9% vs 5.1%; $F_{4,91} = 3.62$, $P = 0.009$; Fig. 1B). Hatch date affected survival, with later-hatching larvae having higher mortality ($F_{1,91} = 7.84$, $P = 0.006$). The number of female pupae per cup was correlated with both weight at and survival to pupation (both $P < 0.001$). There was a significant cultivar-level correlation between mean pupal weight and mean survival to pupation ($R^2 = 0.84$, $F_{1,3} = 15.6$, $P = 0.029$).

Occurrence Assay

We found a total of 108 *Datana* larval clusters over the 6-wk course of the survey. The distribution of larval clusters over time was as follows: 14 on 16 July, 10 on 18 July, 35 on 22 July, one on 23 July, five on 25 July, three on 26 July, one on 29 July, two on 30 July, two on 31 July, 11 on 1 August, 12 on 2 August, one on 6 August, two on 11 August, eight on 20 August, and one on 26 August. Cultivars differed in *Datana* colonization (L-R $\chi^2 = 28.01$ with 12 df, $P = 0.006$; Fig. 2), with 'Bluehaven', 'Collins', and 'Darrow' having more *Datana* clusters (1.00 per bush, 0.75 per bush, and 0.85 per bush, respectively) than either 'Jersey' or 'Earliblue' (0.13 and 0.07 per bush, respectively). Neither bush height (L-R $\chi^2 = 0.52$ with 1 df, $P = 0.47$) nor column (L-R $\chi^2 = 39.3$ with 29 df, $P = 0.096$) affected *Datana* colonization.




Fig. 1. Mean (\pm SE) *Datana drexelii* pupal weight (A) and mean (\pm SE) *D. drexelii* survival to pupation (B) when reared on five different *Vaccinium corymbosum* cultivars. Bars with different uppercase letters are significantly different (Tukey's HSD at $\alpha = 0.05$).

Discussion

Datana larval performance in the lab did not overlap with field observations. Despite high larval performance on 'Lateblue', larval occurrence on it was not the highest on this cultivar in the




Fig. 2. Mean (\pm SE) *D. drexelii* larval clusters counted per bush for 13 *V. corymbosum* cultivars over the course of 6 wk and 15 censuses. Cultivars in dark blue were included in the performance assay (Fig. 1). Bars with different uppercase letters are significantly different (Tukey's HSD at $\alpha = 0.05$).

field survey. High *Datana* densities on 'Bluehaven', 'Collins', and 'Darrow', cultivars not included in our performance assay, suggest they may be particularly suitable to this pest. In contrast, both occurrence and performance were significantly lower for 'Earliblue' and 'Jersey' than other tested cultivars (Figs. 1 and 2). This implies that 'Earliblue' and 'Jersey' may have antixenotic and antibiotic effects on *Datana*.

Mortality in the performance assay occurred mostly within a week of hatching, when larvae were small and inconspicuous. Our high early larval mortality across treatments could indicate that either cultivated blueberry is unsuitable for this species (comparison to wild-type blueberry is needed to determine this), or perhaps that the unnaturally low early instar densities could be reducing the feeding ability, and thus survival, of hatchlings (D. L. Wagner, personal communication). It is also possible that the quality of the cut blueberry may diminish more quickly than the foliage is replaced (within 24 h instead of 3 d), malnourishing larvae. If similarly high levels of hatchling mortality also occurred in the field survey, we could have missed some oviposition events when all larvae died prior to reaching a detectable size. Because of this, the patterns in our field survey data likely reflect some combination of female oviposition preference and plant resistance to early-instar larval feeding. While most larval clusters contained a similar number (10–20 individuals) of small 2nd–3rd instar larvae, we failed to detect some clusters until they contained 4th–5th instar larvae. The laboratory-based oviposition choice tests necessary to isolate the role of female preference may be complicated by this species' habit of readily ovipositing on container walls and other artificial objects.

Our work could be extended to comparisons of *Datana* interactions with cultivated versus wild-type blueberry, as well as with other *Vaccinium* species. Selective breeding for pest resistance (Lobos and Hancock 2015) and the incorporation of several related *Vaccinium* species into *V. corymbosum* cultivars (Lobos and Hancock 2015) may alter the cultivar's suitability to *Datana*. Both 'Lateblue' and 'Jersey', cultivars on which larvae did the best and worst, are pure *V. corymbosum*, but 'Bluecrop' is 4% *Vaccinium angustifolium* (Aiton, 1789) and 'Bluetta' is 28% *V. angustifolium*. Some cultivars are only 42% *V. corymbosum* and contain genes from up to five other species (Lobos and Hancock 2015). Intrageneric variation in herbivore susceptibility has been described for other *Vaccinium* (Ieri et al. 2013) species as well as for genera ranging from *Asclepias* L. (Waterbury et al. 2019) to *Quercus* L. (Rieske and Dillaway 2008).

In summary, there were blueberry cultivar-related differences in occurrence and performance of this blueberry defoliator. This information could prove useful for cultivar selection in areas where this pest becomes a major problem, and highlights how understanding plant–pest interactions can help reduce the need for costly chemical or mechanical (removal of individual larval clusters) treatments.

Acknowledgments

D. Vadnais assisted with larval rearing, and O. Barsoian, D. Butler, C. Johnson, M. Requentina, and L. Varkonyi assisted with oviposition surveys. M. Goldsmith provided helpful comments on an earlier version of this manuscript.

References Cited

- Calvo, D., and J. M. Molina. 2010. Differences in foliage affect performance of the lappet moth, *Streblote panda*: implications for species fitness. *J. Insect Sci.* 10: 177.

- Clift, A. D., G. Murdoch, and S. Mansfield. 2017. Blueberry cultivars differ in susceptibility to the elephant weevil, *Orthorhinus cylindrirostris* (Coleoptera: Curculionidae). *J. Econ. Entomol.* 110: 2259–2262.
- Cutler, B. L., and M. K. Harris. 1979. Foliage consumption and damage by the walnut caterpillar on pecan in Texas USA. *J. Econ. Entomol.* 72: 315–318.
- Follett, P. A., F. T. Zee, R. T. Hamasaki, K. Hummer, and S. T. Nakamoto. 2011. Susceptibility of low-chill blueberry cultivars to Mediterranean fruit fly, oriental fruit fly, and melon fly (Diptera: Tephritidae). *J. Econ. Entomol.* 104: 566–570.
- Harris, M. K. 1983. Outbreak biology of walnut caterpillar *Datana integerrima* in Texas USA. *Southwest. Entomol.* 8: 231–240.
- Hernandez-Cumplido, J., M. M. Giusti, Y. Zhou, V. Kyryczenko-Roth, Y. H. Chen, and C. Rodriguez-Saona. 2018. Testing the ‘plant domestication-reduced defense’ hypothesis in blueberries: the role of herbivore identity. *Arthropod Plant Interact.* 12: 483–493.
- Ieri, F., S. Martini, M. Innocenti, and N. Mulinacci. 2013. Phenolic distribution in liquid preparations of *Vaccinium myrtillus* L. and *Vaccinium vitis idaea* L. *Phytochem. Anal.* 24: 467–475.
- Liburd, O. E., S. R. Alm, and R. A. Casagrande. 1998. Susceptibility of highbush blueberry cultivars to larval infestation by *Rhagoletis mendax* (Diptera: Tephritidae). *Environ. Entomol.* 27: 817–821.
- Lobos, G. A., and J. F. Hancock. 2015. Breeding blueberries for a changing global environment: a review. *Front. Plant Sci.* 6: 782.
- Lyrene, P. M. 1992. Early defoliation reduces flower bud counts on rabbiteye blueberry. *HortScience.* 27: 783–785.
- Rieske, L. K., and D. N. Dillaway. 2008. Response of two oak species to extensive defoliation: tree growth and vigor, phytochemistry, and herbivore suitability. *For. Ecol. Manage.* 256: 121–128.
- Rodriguez-Saona, C., C. Vincent, and R. Isaacs. 2019. Blueberry IPM: past successes and future challenges. *Annu. Rev. Entomol.* 64: 95–114.
- Sanchez-Hernandez, C., M. G. López, and J. P. Délano-Frier. 2006. Reduced levels of volatile emissions in jasmonate-deficient spr₂ tomato mutants favour oviposition by insect herbivores. *Plant. Cell Environ.* 29: 546–557.
- SAS. 2010. JMP user’s guide, version 9.0 computer program. SAS, Cary, NC.
- Stamp, N. E., and Y. Yang. 1996. Response of insect herbivores to multiple allelochemicals under different thermal regimes. *Ecology.* 77: 1088–1102.
- Trivedy, K., S. N. Kumar, N. Vinutha, and S. Qadri. 2011. In vitro testing of common disinfectants used in sericulture to control the growth of fungi in rearing houses. *Res. J. Microbiol.* 6: 439–465.
- Turcotte, M. M., N. E. Turley, and M. T. Johnson. 2014. The impact of domestication on resistance to two generalist herbivores across 29 independent domestication events. *New Phytol.* 204: 671–681.
- Wagner, D. L. 2005. Caterpillars of Eastern North America. Princeton University Press, Princeton, NJ.
- Waterbury, B., A. Potter, and L. K. Svancara. 2019. Monarch butterfly distribution and breeding ecology in Idaho and Washington. *Front. Ecol. Evol.* 7: 172.
- Williamson, J. G., and E. P. Miller. 2000. Early fall defoliation of southern highbush blueberry inhibits flower bud initiation and retards flower bud development. *HortScience.* 35: 505.
- Yang, Y., and N. E. Stamp. 1996. Simultaneous effects of temperature and multiple allelochemicals on the performance of a Solanaceae specialist caterpillar (*Manduca sexta*). *Ecoscience.* 3: 81–92.