

# Geochemistry, Geophysics, Geosystems

## RESEARCH ARTICLE

10.1029/2018GC007630

### Special Section:

Carbon degassing through volcanoes and active tectonic regions

### Key Points:

- New analyses (753) and compiled data (2,446) for volatiles CO<sub>2</sub>, H<sub>2</sub>O, F, S, and Cl are reported in a global suite of mid-ocean ridge basalts
- Estimated primary MORB CO<sub>2</sub> contents vary from 104 ppm to 1.9 wt% and are the result of variations in mantle composition
- High CO<sub>2</sub> fluxes at ridges correlate with high primary CO<sub>2</sub> contents and are due to the presence of subduction components in MORB sources

### Supporting Information:

- Supporting Information S1
- Data Set S1

### Correspondence to:

K. A. Kelley,  
kelley@uri.edu

### Citation:

Le Voyer, M., Hauri, E. H., Cottrell, E., Kelley, K. A., Salters, V. J. M., Langmuir, C. H., et al. (2019). Carbon fluxes and primary magma CO<sub>2</sub> contents along the global mid-ocean ridge system. *Geochemistry, Geophysics, Geosystems*, 20. <https://doi.org/10.1029/2018GC007630>

Received 18 APR 2018

Accepted 26 SEP 2018

Accepted article online 23 NOV 2018

## Carbon Fluxes and Primary Magma CO<sub>2</sub> Contents Along the Global Mid-Ocean Ridge System

Marion Le Voyer<sup>1,2</sup>, Erik H. Hauri<sup>1,3</sup> , Elizabeth Cottrell<sup>2</sup> , Katherine A. Kelley<sup>4</sup> , Vincent J. M. Salters<sup>5</sup> , Charles H. Langmuir<sup>6</sup> , David R. Hilton<sup>7,8</sup>, Peter H. Barry<sup>9</sup> , and Evelyn Füre<sup>10</sup> 

<sup>1</sup>Department of Terrestrial Magnetism, Carnegie Institution for Science, Washington, DC, USA, <sup>2</sup>Smithsonian Institution, National Museum of Natural History, Washington, DC, USA, <sup>3</sup>Deceased 5 September 2018, <sup>4</sup>Graduate School of Oceanography, University of Rhode Island, Narragansett, RI, USA, <sup>5</sup>National High Magnetic Field Laboratory and Department of Earth, Ocean and Atmospheric Sciences, Florida State University, Tallahassee, FL, USA, <sup>6</sup>Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA, USA, <sup>7</sup>Scripps Institution of Oceanography, University of California San Diego, San Diego, CA, USA, <sup>8</sup>Deceased 7 January 2018, <sup>9</sup>Department of Earth Sciences, University of Oxford, Oxford, UK, <sup>10</sup>Centre de Recherches Pétrographiques et Géochimiques, UMR7358, CNRS - Université de Lorraine, Vandoeuvre-lès-Nancy, France

**Abstract** The concentration of carbon in primary mid-ocean ridge basalts (MORBs), and the associated fluxes of CO<sub>2</sub> outgassed at ocean ridges, is examined through new data obtained by secondary ion mass spectrometry (SIMS) on 753 globally distributed MORB glasses. MORB glasses are typically 80–90% degassed of CO<sub>2</sub>. We thus use the limited range in CO<sub>2</sub>/Ba (81.3 ± 23) and CO<sub>2</sub>/Rb (991 ± 129), derived from undegassed MORB and MORB melt inclusions, to estimate primary CO<sub>2</sub> concentrations for ridges that have Ba and/or Rb data. When combined with quality-controlled volatile-element data from the literature ( $n = 2,446$ ), these data constrain a range of primary CO<sub>2</sub> abundances that vary from 104 ppm to 1.90 wt%. Segment-scale data reveal a range in MORB magma flux varying by a factor of 440 (from  $6.8 \times 10^5$  to  $3.0 \times 10^8$  m<sup>3</sup>/year) and an integrated global MORB magma flux of  $16.5 \pm 1.6$  km<sup>3</sup>/year. When combined with CO<sub>2</sub>/Ba and CO<sub>2</sub>/Rb-derived primary magma CO<sub>2</sub> abundances, the calculated segment-scale CO<sub>2</sub> fluxes vary by more than 3 orders of magnitude ( $3.3 \times 10^7$  to  $4.0 \times 10^{10}$  mol/year) and sum to an integrated global MORB CO<sub>2</sub> flux of  $1.32_{-0.85}^{+0.77} \times 10^{12}$  mol/year. Variations in ridge CO<sub>2</sub> fluxes have a muted effect on global climate; however, because the vast majority of CO<sub>2</sub> degassed at ridges is dissolved into seawater and enters the marine bicarbonate cycle. MORB degassing would thus only contribute to long-term variations in climate via degassing directly into the atmosphere in shallow-water areas or where the ridge system is exposed above sea level.

**Plain Language Summary** Estimated CO<sub>2</sub> contents of primary mid-ocean ridge basalts (MORB), calculated on a segment-by-segment basis, vary from 104 ppm to 1.9 wt%. CO<sub>2</sub>-enriched MORB are present in all ocean basins, in particular, in the Atlantic Ocean basin, which is younger and more likely to contain admixed material from recent subduction compared to the much older Pacific Ocean basin. CO<sub>2</sub> fluxes at individual ridge segments vary by 3 orders of magnitude due primarily to large variability in primary CO<sub>2</sub> content. This study provides the most detailed and accurate estimate to date of the integrated total flux of CO<sub>2</sub> from mid-ocean ridges of  $1.32_{-0.85}^{+0.77} \times 10^{12}$  mol/year.

## 1. Introduction

The 56,000-km length of the global mid-ocean ridge system represents the dominant locus of magma production and heat loss on Earth. Decades of deep-water oceanographic research have revealed that new seafloor is produced over a limited range of crustal thicknesses ( $6 \pm 1$  km, Behn & Grove, 2015; Klein & Langmuir, 1987; Van Avendonk et al., 2017; White et al., 1992) and at variable spreading rates that are well-described globally (e.g., Gale et al., 2013). As a result, magma production rates at mid-ocean ridges are well understood and the fluxes of nonvolatile elements emerging from the mantle at ridges are well constrained. These same research cruises have retrieved tens of thousands of individual submarine mid-ocean ridge basalt (MORB) samples, and the study of these samples has revealed that chemical variations in MORB magmas are produced by polybaric melting of mantle sources at variable temperatures, whereas local deviations from global trends are primarily due to variations in mantle composition (Gale et al., 2014; Kelley et al., 2013; Kinzler & Grove, 1992; Klein & Langmuir, 1987; Langmuir et al., 1992).

Carbon is present in MORBs in trace amounts as CO<sub>2</sub>-rich vesicles and as carbonate ions dissolved in quenched seafloor glass (Cartigny et al., 2008; Dixon & Stolper, 1995; Javoy & Pineau, 1991; Marty & Tolstikhin, 1998; Michael, 1995). The solubility of CO<sub>2</sub> in basaltic magma is low at hydrostatic pressures corresponding to seafloor water depths (e.g., Dixon et al., 1995; Jendrzewski et al., 1997; Shishkina et al., 2010), and thus nearly all magmas erupted at mid-ocean ridges have degassed the majority of their carbon. This flux of CO<sub>2</sub> into the oceans contributes to the global surface carbon cycle, and changes in basaltic magma production and associated degassing have been proposed as one of several important forcing mechanisms that have influenced past global climate variations (Burley & Katz, 2015; Robock, 2000; Shindell et al., 2003; Tolstoy, 2015; Zhong et al., 2010). At the same time, it is now widely recognized that most of the Earth's carbon is present in the Earth's deep interior, rather than at Earth's surface (Marty et al., 2013; Marty & Tolstikhin, 1998). Importantly, basaltic magmatism and subduction of oceanic crust represent the two largest outgassing and ingassing fluxes at the interface between the surface and deep-Earth carbon cycles.

These observations provide the motivation of the present study, in which we apply the rapid, precise, and accurate measurement of volatiles by secondary ion mass spectrometry (SIMS; Hauri et al., 2002) to characterize the extensive worldwide sample collection of MORBs to examine in detail the flux of CO<sub>2</sub> from the global ridge system. In this study we report new SIMS data on 753 geographically distributed MORB samples, and combine these data with a quality-controlled database of published data from an additional 2,446 samples (Figure 1). This combined data set permits a description of CO<sub>2</sub> fluxes, and estimated primary magma CO<sub>2</sub> contents, for 381 of the 458 global ridge segments with major element data, in addition to examination of the variability of primary MORB CO<sub>2</sub> contents, the role of carbon in the melting processes that produce MORB, and the role of the oceanic crust in the context of the Earth's deep carbon cycle.

## 2. Methods


Here we briefly describe the methods used in this study for sample preparation, SIMS analysis of volatile elements, major elements, and trace elements.

### 2.1. Sample Preparation

The majority of the basalt glasses analyzed in this study were obtained from the Rock and Ore collection of the Smithsonian Institution's National Museum of Natural History. The equatorial mid-Atlantic Ridge glasses were obtained from the Schilling Volcanic Glass collection at the University of Rhode Island (University of Rhode Island Graduate School of Oceanography Marine Geological Samples Laboratory, 2018). Additional glass samples were provided from the Pacific-Antarctic Ridge (PACANTARCTIC 1 and 2 cruises, Clog et al., 2013; Hamelin et al., 2010; Labidi et al., 2014; Moreira et al., 2008; Vlastelic et al., 1999, 2000), the Central Indian Ridge (GIMNAUT, CD127, and KNOX11RR cruises, Fűri et al., 2011; Murton et al., 2005; Barry, 2012), the Southwest Indian Ridge (PROTEA-5, MD34, AG22, and AG53 cruises, Arevalo & McDonough, 2010; B. Hamelin & Allègre, 1985; Janney et al., 2005; Le Roex et al., 1989; Mahoney et al., 1992), and the East Pacific Rise (CHEPR and PANORAMA-1 cruises, Langmuir, 2014; Salters et al., 2011). The major element, trace element, and isotopic ratio data for the glasses measured in this study were compiled from the literature (Clog et al., 2013; Cottrell & Kelley, 2011; C. Hamelin et al., 2010; Jenner & O'Neill, 2012; Labidi et al., 2014; Le Voyer et al., 2015; Moreira et al., 2008; Vlastelic et al., 2000, 1999; Nauret et al., 2006). We selected 1- to 3-mm glass chips that did not show any sign of alteration, crystallization, or devitrification. The chips were pressed into indium mounts together with secondary standards (ALV519-4-1 and VG2), and polished using silicon carbide papers, diamond paste, and finally 0.3- $\mu$ m alumina paste. We washed the mounts using successive baths of distilled water, ethanol, and acetone, then stored them for >48 hr in a vacuum oven at 70 °C, before applying a gold coat.

### 2.2. Volatile Elements

CO<sub>2</sub> and other volatile elements, along with phosphorus, were measured by SIMS at the Department of Terrestrial Magnetism (DTM), Carnegie Institution of Washington, following the method of Hauri et al. (2002) but with the modification of measuring <sup>16</sup>OH (rather than <sup>1</sup>H) for determination of H<sub>2</sub>O contents. These new data are reported in supporting information Table S1. The analyses were performed during 13 analytical sessions from March 2012 to October 2016, for a combined analytical time of 59 days. For 12 of the 13 sessions, we used a Cs<sup>+</sup>, 15 ± 2 nA primary beam accelerated to 10 kV, with charge compensation



**Figure 1.** Locations of newly measured mid-ocean ridge basalt samples ( $n = 753$ ) are shown as red circles, with locations of previously published samples ( $n = 2,446$ ) shown as gray circles.

provided by an electron flood gun. For one session we used a  $10 \pm 1$  nA primary beam instead, which did not create any differences in the results. For 11 of the 13 sessions we applied a  $25\text{-}\mu\text{m}^2$  raster on the beam, which resulted in a crater size of  $\sim 35\ \mu\text{m}^2$ . For two sessions we used a  $20\text{-}\mu\text{m}^2$  raster size, and no differences in the results were observed. The vacuum was in the  $10^{-9}$  to  $10^{-10}$  torr range during each analysis. We used the smallest field aperture (#4) in order to mask the edge of the crater and extract data from a central  $10\text{-}\mu\text{m}$ -diameter area in the center of the sputter crater. The mass spectrometer was set to a mass resolution sufficient to separate  $^{17}\text{O}$  from  $^{16}\text{OH}$ ,  $^{18}\text{OH}$  from  $^{19}\text{F}$ , and  $^{16}\text{O}_2$  from  $^{32}\text{S}$  (Mass Resolving Power  $\sim 6,000$ ).

**Table 1**  
Concentrations and Reproducibility of Volatile and Phosphorus Contents of Smithsonian Basalt Glass Standard VG2 Measured During the Course of This Study

	CO <sub>2</sub>	H <sub>2</sub> O	F	P	S	Cl
Unit	ppm	wt%	ppm	ppm	ppm	ppm
Number of analyses ( <i>n</i> )	76	74	76	77	75	76
Average over <i>n</i> analyses	187	0.28	211	970	1,486	259
2σ	16	0.02	15	43	53	17
2σ (%)	8	7	7	5	4	7

After 300 s of presputtering, we successively collected signal for 10 s on mass <sup>12</sup>C, 5 s on masses <sup>16</sup>OH, <sup>19</sup>F, <sup>30</sup>Si (reference mass), <sup>31</sup>P, <sup>32</sup>S, and <sup>35</sup>Cl. The waiting time before signal collection was 2 s before <sup>12</sup>C and 1 s before all other masses. This cycle was repeated 5 times. A set of basaltic standards and blanks (see Hauri et al., 2002) were used to perform the calibration (linear regression forced through the origin) and to assess detection levels (average detection levels over the 13 sessions: 5 ppm for H<sub>2</sub>O, 2 ppm for CO<sub>2</sub>, 0.2 ppm for S, Cl, and P, and 0.1 ppm for F). The analytical error (2σ over the five analytical cycles) was typically 4% for CO<sub>2</sub>, 2% for others. Each sample was measured three times, and the reproducibility over the three analyses was typically better than 10% for CO<sub>2</sub> and 5% for

other elements. Standard ALV519-4-1 was mounted together with samples in each indium mount, and measured every nine analyses, in order to correct for potential instrumental drift within each session, potential offsets between sessions, and to assess long-term reproducibility (better than 5%, 2RSD, for all volatiles, over 2 years). Standard VG-2 was also mounted together with samples in each indium mount to assess the reliability of this correction, and results are reported in Table 1. The reproducibility on VG2 was better than 8% for all volatiles and phosphorus (2σ over 2 years).

### 3. Published Data Quality Control

Data from PetDB ([www.earthchem.org/petdb](http://www.earthchem.org/petdb)) for MORB glasses and melt inclusions, consisting of samples containing data for at least one volatile (CO<sub>2</sub>, H<sub>2</sub>O, F, S, Cl) were downloaded in late 2017; the data are compiled, along with the newly reported data from this study, in supporting information Table S1, and the references for the original data are included in the supporting information Table S5. All corresponding data (major elements, trace elements, isotopes) were included in this download. Concentrations were recalculated as weight percent for H<sub>2</sub>O, and parts per million by weight for CO<sub>2</sub>, F, S, and Cl for data reported in other units or as oxides (e.g., C data were converted to CO<sub>2</sub> ppm; SO<sub>2</sub> and SO<sub>3</sub> data were converted to S ppm), and the database was checked for obvious typographic errors by comparison with the original references. Samples with CO<sub>2</sub> and/or H<sub>2</sub>O were retained in the database if the samples were determined by SIMS, Fourier-transform infrared spectroscopy, or post-1995 vacuum extraction of the volatile fraction dissolved in the glass; vesicle-only analyses were discarded due to the inability to use the data to calculate a glass concentration. Loss on ignition data for H<sub>2</sub>O were also discarded, as were four analyses of melt inclusions from Helo et al. (2011) with >3,900 ppm CO<sub>2</sub> that were judged to have been contaminated by carbon coat during SIMS analysis. Samples with F data were retained only if they were determined by SIMS or ion-selective electrode methods (Bach et al., 1994; Schilling et al., 1983); electron probe microanalysis (EPMA) for F is subject to interference from overlapping Fe peaks in basaltic glasses, and these data were thus rejected from the database. Samples with S or Cl data were retained if they were determined by SIMS or EPMA.

We also compiled previously published Fe<sup>3+</sup>/ΣFe ratios of MORB glasses for this study. For this compilation, we report Fe-speciation data collected using micro X-ray absorption near-edge structure (μXANES) spectroscopy, with all data collected at beamline X26A of the National Synchrotron Light Source, Brookhaven National Laboratory, following methods described by Cottrell et al. (2009) and Zhang et al. (2018). The Zhang et al. (2018) study assessed the effects of recoil-free fraction on the room temperature Mössbauer spectra, against which the XANES centroid energy was calibrated by Cottrell et al. (2009). Room temperature Mössbauer spectra overestimate the Fe<sup>3+</sup>/ΣFe ratios of basaltic glasses, and the updated calibration of Zhang et al. (2018) accounts for this effect. The Fe<sup>3+</sup>/ΣFe ratios compiled here include those reported by Cottrell and Kelley (2011) and Le Voyer et al. (2015), both updated using the new calibration of Zhang et al. (2018).

Interlaboratory bias between the DTM SIMS method and published data sets is essentially nonexistent, for several reasons. SIMS calibrations for H<sub>2</sub>O and CO<sub>2</sub> rely on the use of an extensive suite of MORB and back-arc basin glasses previously analyzed by vacuum manometry and/or Fourier-transform infrared spectroscopy (references in Hauri et al., 2002) originating from laboratories that provided the vast majority of previously published data archived in PetDB. The SIMS method for S and Cl is more precise and has lower detection limits than EPMA. The precision of bias factors between data sets is limited by the reproducibility of the EPMA data and is thus indeterminate. For F the SIMS calibration, which relies on only four glasses (Hauri


et al., 2002), was checked against the data sets of Rowe and Schilling (1979), Schilling et al. (1983), and Bach et al. (1994) by SIMS analysis of samples in common with these studies, which used a halogen-selective electrode wet chemical method. We found excellent agreement ( $\pm 8\%$   $2\sigma$ ) with the data set of Bach et al. (1994) but much poorer agreement with the other, significantly older, data sets that used the same method for F analysis. Thus, the comparison for F found no interlaboratory bias but does show clear examples of variable extents of reproducibility of other methods compared with SIMS, which likely permeate much of the published data for fluorine and, to a lesser extent, sulfur and chlorine. In general, interlaboratory comparisons agree to better than 10% for CO<sub>2</sub>, H<sub>2</sub>O, and F, and by a somewhat worse and variable degree for S and Cl.

The supporting information contains the data used in this paper and is available through the EarthChem Library (<https://doi.org/10.1594/IEDA/111195>). Table S1 contains the results of new analyses of MORB glasses, together with associated data on major and trace elements, and isotopic compositions. In Table S2, these new data are compiled along with the quality-controlled published data in a single table. Table S3 is a compilation of data (volatiles, major elements, trace elements, isotope compositions of host glass) for MORB melt inclusions, for comparison to the submarine glass data. Table S4 is a modification of the segment-averaged tables from Gale et al. (2013, 2014) that includes data for segment metrics (length, spreading rate, depth, etc.) as well as averaged data for major elements, trace elements, volatiles, and isotopic compositions; this table also includes estimated primary magma CO<sub>2</sub> concentrations, and calculated CO<sub>2</sub> fluxes on a per-segment basis, with estimated uncertainties shown in yellow highlight. Table S5 contains an alphabetic list of all the publications that contained the original data that were used in the data compilations of all of the data tables.

#### 4. Results


The samples in this compiled volatile data set were assigned to individual ridge segments based on a comparison of their sampling coordinates with the global spreading center segment catalog of Gale et al. (2013; Figure 1). In constructing geochemical averages of global spreading segments, Gale et al. (2013) included in their compilation samples within 10 km of the ridge axis. At the slowest spreading rates, a distance of 10 km off-axis corresponds to an age of approximately 1.5 million years, and so in our study individual samples were categorized as *on-axis* if their apparent age was less than or equal to 1.5 million years. Samples older or farther than this were categorized as either *off-axis* or as *transform* samples depending on their location relative to the nearest ridge axis or transform fault. Segment average volatile concentrations were then calculated for all segments that contained at least one on-axis sample analyzed for H<sub>2</sub>O, CO<sub>2</sub>, F, S, and/or Cl. The PetDB data download also contained a small amount of new major element, trace element, and isotope data published after the compilations of Gale et al. (2013) and Class and Lehnert (2012), and this permitted the calculation of several new segment averages for major elements, trace elements, and isotopes to augment the database. For consistency, these updated segment averages were calculated using the same methods described in Gale et al. (2013); however, it should be mentioned that only samples with volatile data were used to calculate these additional new segment averages. Finally, we recalculated the segment-averaged major and trace element data for segment EPRR31 to account for two additional samples from this segment analyzed by Cottrell and Kelley (2011); samples from this segment have a bimodal distribution of depleted ( $n = 3$ ) and enriched ( $n = 3$ ) MORB. By doing this, the segment averages for all data (major elements, trace elements, volatiles) all come from the same six samples.

The MORB glasses measured in this study range from 2 to 420 ppm CO<sub>2</sub>, 0.04 to 1.37 wt% H<sub>2</sub>O, 41 to 1,131 ppm F, 121 to 2,380 ppm S, and 1.7 to 5,540 ppm Cl. These new volatile data extend the range of MORB H<sub>2</sub>O, F, and Cl abundances, while the data for CO<sub>2</sub> and S cover most of the range defined by previously published data for MORB (up to 600 ppm CO<sub>2</sub> and 3,300 ppm S). H<sub>2</sub>O and F are broadly correlated in the global data (Figure 2a), with an average H<sub>2</sub>O/F ratio of 14.3 but with a large range displayed by individual samples (0.56 to 216). Sulfur is correlated with total Fe (FeO\*) (Figure 2b), reflecting the dependence of sulfide solubility on iron content in mafic melts (Keith et al., 2017; O'Neill & Mavrogenes, 2002). Chlorine is generally not well correlated with H<sub>2</sub>O, but a lower bound to the global data is apparent with a Cl/H<sub>2</sub>O ratio in the range of 0.01 to 0.05. Most of the samples with higher Cl/H<sub>2</sub>O ratios have likely undergone assimilation of Cl-rich seawater-derived components (Le Roux et al., 2006; Le Voyer et al., 2015; Michael & Schilling, 1989). There does not appear to be any relationship of the volatiles with the oxidation state of Fe in MORBs examined in this study.



**Figure 2.** (a) Fluorine versus H<sub>2</sub>O and (b) FeO\* versus S in mid-ocean ridge basalt glasses from this study. Fluorine and H<sub>2</sub>O are not degassed at seafloor pressures typical of mid-ocean ridge basalt eruptions. The correlation of S with FeO\* reflects the control of FeO content on the sulfur solubility in basaltic melts, which are typically saturated in a Fe-Ni-Cu monosulfide melt (Jenner et al., 2015).

Segment averages for the volatile data were corrected to MgO = 8 wt% and to compositions in equilibrium with Fo<sub>90</sub> olivine, in order to compare with the fractionation-normalized database of Gale et al. (2014). This correction for crystal fractionation cannot account for the irreversible loss of CO<sub>2</sub> by degassing; however, we will estimate primary magma CO<sub>2</sub> contents using Ba and Rb as proxies, in a subsequent section (below). The scarcity of available volatile data at any given ridge segment, coupled with potential CO<sub>2</sub>, H<sub>2</sub>O, or S loss by magmatic outgassing, resulted in poorly defined liquid lines of descent (LLDs) for the volatile data. Thus the 8% MgO corrections for the volatile segment averages were done by calculating a fractionation factor from the ratio of the segment average P<sub>2</sub>O<sub>5</sub>(8) divided by the segment average P<sub>2</sub>O<sub>5</sub> (uncorrected) using the data given in Gale et al. (2013, 2014). P<sub>2</sub>O<sub>5</sub> was chosen because it typically forms better-correlated and more linear LLDs than K<sub>2</sub>O; nevertheless, for segments lacking P<sub>2</sub>O<sub>5</sub> data the K<sub>2</sub>O(8) and K<sub>2</sub>O data were used. An identical procedure was used to further correct the volatile segment averages to Fo<sub>90</sub> values, by determining a fractionation factor from the ratio of K<sub>2</sub>O(90) to K<sub>2</sub>O(8) for each segment reported by Gale et al. (2014). Uncertainties from numerous sources were quantified throughout the examination of the data. Analytical uncertainties for CO<sub>2</sub> and other elements are typically minor (a few percent relative) compared to the compositional heterogeneity within individual ridge segments; the standard deviations of segment averages were quantified in Gale et al. (2013) while standard deviations for compositions corrected to 8 wt% MgO were quantified in Gale et al. (2014). For the purposes of calculating segment magma fluxes, we have assigned a ±4 km uncertainty on ridge length, ±3 mm/year uncertainty in spreading rates, and a variable uncertainty on crustal thickness (ranging from 3% to 50%) determined by the uncertainty of a regression of estimated errors in crustal thickness versus absolute thickness determined seismically (Behn & Grove, 2015; Van Avendonk et al., 2017).



**Figure 3.** Histogram of the pressure of vapor saturation ( $P_{\text{sat}}$ ) normalized to the pressure of sampling ( $P_{\text{sample}}$ ) for the mid-ocean ridge basalt glasses measured in this study. The distribution is strongly peaked near unity but displays a tail to high ratios (oversaturation in  $\text{CO}_2\text{-H}_2\text{O}$  vapor) due to rapid magma ascent, during which vesicle growth is limited by the diffusion of carbon in the melt.

The uncertainty that is most difficult to quantify is our assumption of single average values for the ratios  $\text{CO}_2/\text{Ba}$  ( $81.3 \pm 28\%$ ) and  $\text{CO}_2/\text{Rb}$  ( $991 \pm 13\%$ ) for MORB; these values are the averages of the population averages of six groups of  $\text{CO}_2$ -undersaturated MORB glasses and melt inclusions (Hauri et al., 2017). To the extent that normal MORB (N-MORB) can be characterized as a mixture of depleted MORB (D-MORB) and enriched MORB (E-MORB; e.g., Donnelly et al., 2004), this assumption implies that D-MORB and E-MORB sources have the same  $\text{CO}_2/\text{Ba}$  and  $\text{CO}_2/\text{Rb}$  ratios. Two studies support this assumption. Shimizu et al. (2016) estimated  $\text{CO}_2$  contents in Pacific E-MORB from their Cl contents and the constant  $\text{CO}_2/\text{Cl}$  ratio in Siqueiros Fracture Zone D-MORB melt inclusions, yielding  $\text{CO}_2/\text{Ba}$  ratios of  $105 \pm 61$  for Pacific D-MORB and  $134 \pm 23$  for Pacific EMORB. Similarly, in the Atlantic the D-MORB melt inclusion suite of Le Voyer et al. (2017) has  $\text{CO}_2/\text{Ba}$  of  $97 \pm 10$ , very close to the  $\text{CO}_2/\text{Ba}$  of  $106 \pm 34$  of the highly enriched E-MORB popping rocks from  $14^\circ\text{N}$  on the Mid-Atlantic ridge (MAR; Cartigny et al., 2008; Hauri et al., 2017). Despite the possibility that this assumption may eventually be contradicted, it presently provides a data-supported basis from which to estimate the primitive, pre-eruptive  $\text{CO}_2$  content for degassed MORB. We emphasize here that a key underpinning of our estimations of undegassed global MORB  $\text{CO}_2$  contents is the assumed constancy of  $\text{CO}_2/\text{Ba}$  and/or  $\text{CO}_2/\text{Rb}$  across the full compositional spectrum of MORBs; uncertainties on the  $\text{CO}_2$  concentrations estimated here will increase if these ratios are later shown to vary. We calculate unde-

gassed MORB  $\text{CO}_2$  concentrations by multiplying these ratios by the Ba and/or Rb concentrations measured in the basalts. In our calculations of primary  $\text{CO}_2$  content, segment-scale  $\text{CO}_2$  fluxes, and overall ridge flux we have combined all of these sources of error in quadrature, resulting in the stated uncertainties listed with the data in the yellow-highlighted columns of Table S4.

## 5. Degassing of $\text{CO}_2$ From MORB

MORBs are erupted on the seafloor under the hydrostatic pressure of the overlying water column. The solubility of  $\text{CO}_2$  in MORB is strongly dependent on pressure (Dixon & Stolper, 1995), which would predict that MORB  $\text{CO}_2$  concentrations should correlate with the pressure of eruption on the seafloor. Yet MORBs often have dissolved  $\text{CO}_2$  contents that exceed the equilibrium solubility predicted at the eruption pressure. In most cases, this oversaturation is due to rapid melt ascent during eruption, during which diffusion of carbon in the melt is too slow to maintain equilibrium between the dissolved  $\text{CO}_3^{2-}$  in the melt and the  $\text{CO}_2$  vapor within the vesicles (Chavrit et al., 2012; Javoy & Pineau, 1991).

The vapor solubility model of Dixon et al. (1995) has long been used to constrain vapor saturation pressure ( $P_{\text{sat}}$ ) in MORBs because these authors determined  $\text{CO}_2$  and  $\text{H}_2\text{O}$  solubility specifically on MORB melt compositions. When we use measured  $\text{CO}_2$  and  $\text{H}_2\text{O}$  contents in MORBs to calculate  $P_{\text{sat}}$  using the Dixon et al. (1995) model, and compare these with the pressures of MORB sample collection ( $P_{\text{sample}}$ ), the distribution of the  $P_{\text{sat}}/P_{\text{sample}}$  ratio is strongly peaked near unity (median = 1.37) but is clearly skewed toward high  $P_{\text{sat}}/P_{\text{sample}}$  ratios (Figure 3). This skew is likely due to vapor oversaturation in MORBs driven by high magma ascent rates, as discussed above. At the same time, there is a sharp cut-off below unity, with few samples displaying  $P_{\text{sat}}/P_{\text{sample}}$  significantly  $< 1$ .

A similar comparison of  $P_{\text{sample}}$  with  $P_{\text{sat}}$  predicted by other experimental solubility studies provides a rough approximation of the accuracy of these solubility models. The model of Papale et al. (2006) produces  $P_{\text{sat}}/P_{\text{sample}}$  ratios that are 60% lower on average compared with Dixon et al. (1995), and suggests that most MORB are vapor undersaturated. Vesicle abundances observed in MORB, however, contradict this prediction (Chavrit et al., 2012), indicating that the Papale et al. (2006) model likely underestimates vapor saturation pressures in MORB. The Papale et al. (2006) model was calibrated over a very wide range in melt compositions, whereas the Dixon et al. (1995) model was calibrated specifically on MORB


compositions at pressures  $<1$  kbar, so disagreement between these models is perhaps not surprising. The solubility model of Iacono-Marziano et al. (2012) provides values of  $P_{\text{sat}}$  for MORB that are within 2% of those calculated from the Dixon et al. (1995) model. More recently, the  $\text{CO}_2$ - $\text{H}_2\text{O}$  solubility model of Ghiorso and Gualda (2015) results in  $P_{\text{sat}}/P_{\text{sample}}$  ratios that are 20% lower on average compared with Dixon et al. (1995), with a peak of the  $P_{\text{sat}}/P_{\text{sample}}$  ratio distribution closer to unity (median 1.10) and a lower limit of the ratio of  $\sim 0.6$ . The Ghiorso and Gualda (2015) model thus also predicts more samples than Dixon et al. (1995) to be vapor undersaturated. Both models provide reasonable estimates of vapor saturation pressures, but we prefer the model of Dixon et al. (1995) because it is specific to MORB and it predicts that fewer MORBs are vapor undersaturated, which is consistent with the presence of vesicles in the vast majority of MORBs.

The ratios of volatile to nonvolatile elements have been useful to examine processes of loss or gain of volatile elements, such as degassing, dehydration, and shallow-level assimilation that do not affect nonvolatile elements that otherwise share similar mineral-melt partitioning behavior during partial melting and fractional crystallization. Further constraints on the extent of degassing of individual MORB samples can thus be obtained by examination of the ratios of  $\text{CO}_2/\text{Rb}$  and  $\text{CO}_2/\text{Ba}$ , which exhibit a narrow range in genuine vapor-undersaturated MORB glasses and melt inclusions. Rosenthal et al. (2015), Michael and Graham (2015), and Hauri et al. (2017) recently reviewed the data for vapor-undersaturated MORB; in this compilation we use average ratios  $\text{CO}_2/\text{Rb}$  ( $991 \pm 26\%$ ) and  $\text{CO}_2/\text{Ba}$  ( $81.3 \pm 56\%$ ) to represent undegassed MORBs, which are calculated as the average of the population averages for six groups of vapor-undersaturated samples given in Table 1 of Hauri et al. (2017).  $\text{CO}_2/\text{Rb}$  and  $\text{CO}_2/\text{Ba}$  ratios are much more uniform than  $\text{CO}_2/\text{Nb}$  or  $\text{CO}_2/\text{Th}$  ratios in undegassed MORBs. The ratio of the measured  $\text{CO}_2/\text{Ba}$  ratio in degassed MORBs to the average  $\text{CO}_2/\text{Ba}$  ratio for undegassed MORB gives a measure of the amount of  $\text{CO}_2$  lost by degassing. The extent of degassing of typical MORB samples estimated in this way demonstrates that a majority of samples degassed  $>70\%$  of their initial inventory of  $\text{CO}_2$  (Figure 4). Approximately 3% of MORB samples, however, exhibit  $\text{CO}_2/\text{Rb}$  ratios close to the undegassed ratio, and most of these samples have been identified previously (Hauri et al., 2017; Le Voyer et al., 2017; Michael & Graham, 2015; Saal et al., 2002; Shimizu et al., 2016). Based on this analysis, emplacement of MORB on the seafloor typically results in eruptive degassing into the oceans of  $\geq 70\%$  of the primary carbon carried by MORB magmas.

The  $\text{CO}_2/\text{Ba}$  ratios of melt inclusions from mid-ocean ridges also bear importantly on the carbon budget of MORB magmas. The olivine-hosted melt inclusion data of Saal et al. (2002; Siqueiros FZ, East Pacific Rise), Le Voyer et al. (2017; equatorial mid-Atlantic Ridge), and Hauri et al. (2017; northern Iceland), for example, all display excellent correlations of  $\text{CO}_2$  with Ba (and with Rb, where measured), demonstrating the undegassed nature of these melt inclusion populations. An additional 113 melt inclusions have been measured for  $\text{CO}_2$  and Ba from the Gakkel Ridge (Shaw et al., 2010; Wanless et al., 2014), the Juan de Fuca Ridge (Helo et al., 2011; Wanless & Shaw, 2012), and from the Lucky Strike segment of the mid-Atlantic Ridge (Wanless et al., 2015). Of these 113 melt inclusions, 22 exhibit  $\text{CO}_2/\text{Ba}$  ratios within 20% of the average undegassed ratio of 81.3 (Figure 4b). The majority have likely experienced  $\text{CO}_2$  degassing either prior to melt inclusion entrapment or into secondary vapor bubbles that often form within melt inclusions during post-entrapment cooling (e.g., Moore et al., 2015) but are difficult to measure quantitatively. More data could be readily obtained from MORB melt inclusions through concentrated efforts to quantify the effect of vapor bubbles within melt inclusions (e.g., Aster et al., 2016) and by targeting ridge segments that erupt samples on the seafloor that have experienced little to no degassing (e.g., Le Voyer et al., 2017).

Matthews et al. (2017) present an alternative interpretation of certain suites of MORBs with  $\text{CO}_2$  that is well correlated with highly incompatible trace elements. Such suites are often interpreted as undegassed with respect to  $\text{CO}_2$ , because the degassing process fractionates  $\text{CO}_2$  from other highly incompatible elements. Matthews et al. (2017) instead envisages a process by which magmas emerging from the melting regime stall at a depth where enriched magma batches become oversaturated in  $\text{CO}_2$ , leading to degassing. This degassing is followed by limited mixing at shallower levels, which can produce correlated  $\text{CO}_2/\text{Ba}$  ratios in MORBs that are lower than the original mantle source  $\text{CO}_2/\text{Ba}$  ratio. Yet this process is limited in its ability to explain  $\text{CO}_2/\text{Ba}$  ratios in highly depleted MORB, many of which arrive on the seafloor with low  $\text{CO}_2$  contents but also






**Figure 4.** (a) CO<sub>2</sub> versus Ba for suites of vapor-undersaturated melt inclusions and mid-ocean ridge basalt (MORB; squares, summarized in Hauri et al., 2017) together with data from globally distributed MORB (crosses). (b) Same diagram as panel (a) showing only MORB glasses and MORB melt inclusions from Shaw et al. (2010), Wanless & Shaw (2012), Helo et al. (2011), and Wanless et al. (2014, 2015). The heavy black line represents the average CO<sub>2</sub>/Ba (81.3) derived from the CO<sub>2</sub>/Ba ratio of undersaturated samples. Red dashed lines with labels indicate the extent of CO<sub>2</sub> degassing.

vapor undersaturated. Such magmas would need to degas CO<sub>2</sub> at pressures significantly lower than their seafloor collection pressures (e.g., Siqueiros data of Saal et al. (2002); Quebrada-Discovery-Gofar Transform data of Shimizu et al. (2016); the vapor-undersaturated data of Michael and Graham (2015)), or at pressures within the eruptive Layer 2 of the oceanic crust (equatorial MAR data of Le Voyer et al. (2017)). For the more intermediate-enriched data sets (Borgarhraun, Iceland, and Top 10% MORB data sets of Hauri et al., 2017; 14°N MAR, Cartigny et al., 2008) the highest CO<sub>2</sub> contents vary widely, and each sample suite would need to have stalled and degassed at a very specific depth (e.g., not just the Moho depth) in order to produce such similar CO<sub>2</sub>/Ba ratios as reported in Hauri et al. (2017). We thus conclude that, although the processes in Matthews et al. (2017) are plausible, they are unlikely to have operated in the samples that currently constrain the CO<sub>2</sub>/Ba ratio of MORB sources.

## 6. Variations in Primary MORB CO<sub>2</sub> Contents

The restricted ranges in CO<sub>2</sub>/Rb and CO<sub>2</sub>/Ba ratios of undegassed MORB permit an estimation of the initial (undegassed) CO<sub>2</sub> concentration in MORB primary magmas, given an estimate of their primary Rb and/or Ba abundances. The studies of Gale et al. (2013, 2014) provide such estimates based on the compositions of MORB magmas averaged by ridge segment, and corrected for shallow-level crystal fractionation to MgO = 8 wt% and also further corrected to be in equilibrium with Fo<sub>90</sub> olivine. The data compilations of



**Figure 5.** Segment-averaged estimated primary  $\text{CO}_2$  versus (a) segment-averaged primary Th and (b) segment-averaged primary  $\text{K}_2\text{O}$  for global mid-ocean ridge basalt;  $\text{CO}_2$  is estimated from the segment-averaged Ba content multiplied by a constant  $\text{CO}_2/\text{Ba}$  (81.3).


Gale et al. (2014) and this study provide primary MORB magma estimates for 403 segments that contain data for Ba and/or Rb, out of the 711 global spreading ridge segments identified by Gale et al. (2013). These segment-averaged concentrations were then multiplied by the undegassed-average ratios of  $\text{CO}_2/\text{Rb}$  (991) and  $\text{CO}_2/\text{Ba}$  (81.3) to calculate segment-averaged primary  $\text{CO}_2$  concentrations. MORB primary  $\text{CO}_2$  thus calculated varies over a factor of 180, from 104 ppm to 1.90 wt%. These calculated primary  $\text{CO}_2$  contents further enable calculation of  $\text{CO}_2/\text{Nb}$  and  $\text{CO}_2/\text{Th}$  ratios for all segments that have segment-averaged Nb and Th data. These latter ratios are used to compare with prior studies that used  $\text{CO}_2/\text{Nb}$  ratios (e.g., Cartigny et al., 2008; Saal et al., 2002), while Th is another highly incompatible element that has a mantle melt partition coefficient similar to  $\text{CO}_2$  (Rosenthal et al., 2015), and there exists Th data of sufficient quality in MORB and MORB melt inclusions to make a useful comparison (Hauri et al., 2017).

Primary MORB  $\text{CO}_2$  abundances can be compared with other indices of depletion and enrichment in MORB such as the  $\text{K}_2\text{O}/\text{TiO}_2$  ratio, which is low in depleted MORB and high in enriched MORB. The lowest calculated primary MORB  $\text{CO}_2$  abundance is 104 ppm  $\text{CO}_2$  at a depleted segment of the Galapagos Spreading Center (segment GALA16 with  $\text{K}_2\text{O}/\text{TiO}_2$  of 0.05), and the highest is 1.90 wt%  $\text{CO}_2$  at a highly enriched segment of the Southwest Indian Ridge (segment SWIR9 with  $\text{K}_2\text{O}/\text{TiO}_2 = 0.66$ ). The distribution of MORB primary  $\text{CO}_2$  contents is log-normal with a median value of 2,100 ppm. In general, the segment-averaged primary  $\text{CO}_2$  contents, as derived from reconstructed primary Ba and Rb concentrations, correlate well with segment-averaged primary Th and  $\text{K}_2\text{O}$  contents (Figure 5). Such correlations are most reasonably interpreted as signatures of variation in mantle source composition, rather than variable mantle temperature or spreading rate. In this view, variations in the primary MORB  $\text{CO}_2$  contents determined here would also reflect mantle source variations.

Segment-averaged  $\text{CO}_2/\text{Nb}$  ratios calculated above vary by more than a factor of 10, from a low of 152 at the very depleted segment GALA16 on the Galapagos Spreading Center, to a high of 1,760 at enriched segment SWIR22 on the Southwest Indian Ridge. This range in  $\text{CO}_2/\text{Nb}$  contrasts with the much more limited range in  $\text{CO}_2/\text{Nb}$  measured in undegassed MORB glasses or melt inclusion populations (230–730, Cartigny et al., 2008; Hauri et al., 2017; Saal et al., 2002). We emphasize again that the  $\text{CO}_2/\text{Nb}$  ratios constrained here effectively reflect variations in Ba/Nb or Rb/Nb in MORB because of the assumptions adopted in the calculation of  $\text{CO}_2$  concentrations. Yet the calculated segment-averaged  $\text{CO}_2/\text{Nb}$  ratios display rough trends with segment-averaged Sr and Nd isotopes (Figure 6) that are consistent with the relationships observed by Hauri et al. (2017) and further support the inference that primary  $\text{CO}_2$  contents of MORB, inasmuch as these are inseparable from Ba and Rb, are a function of mantle composition. It remains to be seen if this large range in estimated  $\text{CO}_2/\text{Nb}$  ratios is reflected in actual sample measurements, but the data in this study provide a first-order prediction and help identify specific ridge segments to focus on to test the accuracy of these estimates. The variations are consistent with the dichotomy of  $\text{CO}_2/\text{Nb}$  and  $\text{CO}_2/\text{Th}$  ratios between Atlantic and Pacific Ocean basins, which could be a result of a greater amount of recycled subduction zone components in the Atlantic Ocean basin compared with the relatively older Pacific Ocean basin (Dixon et al., 2017; Hauri et al., 2017).

## 7. The Flux of $\text{CO}_2$ From the Global Ridge System

Crustal thickness at mid-ocean ridges is known to vary from a few kilometers to over 20 km at northern Iceland (Menke, 1999), controlled by a combination of the local magma production rate and the spreading rate. The study of Behn and Grove (2015) summarized seismically determined measurements of crustal thickness at ridge axes, where it varies from 2 km at the slow-spreading Gakkel Ridge to 30 km at central Iceland; additional data for crust <20 Ma old was taken from Van Avendonk et al. (2017). Seismically



**Figure 6.** Heat map diagram showing relationship between segment-averaged calculated mid-ocean ridge basalt  $\text{CO}_2/\text{Nb}$  ratios and (a) Sr isotopes and (b) Nd isotopes.  $\text{CO}_2/\text{Nb}$  ratios were calculated using segment-averaged  $\text{CO}_2(8)$  contents estimated from segment-averaged Ba(8) (Gale et al., 2014) multiplied by  $\text{CO}_2/\text{Ba}$  of 81.3; this estimated primary  $\text{CO}_2$  content is divided by segment-averaged Nb(8) to calculate primary  $\text{CO}_2/\text{Nb}$ . Yellow colors indicate increasing density of data points. Isotope ratios are segment averages from Gale et al. (2013). The pink squares are the average values for six populations of vapor-undersaturated glasses and melt inclusions summarized by Hauri et al. (2017).


constraints. This estimate lies within the range of  $\text{CO}_2$  fluxes estimated by Resing et al. (2004) from  $\text{CO}_2$  and  $^3\text{He}$  fluxes into the water column above ridges, and by Barry et al. (2014) in their study of Iceland  $\text{CO}_2$  fluxes. Our estimate overlaps the lower average flux found by Saal et al. (2002;  $0.93 \pm 0.28 \times 10^{12}$  mol/year), who based their estimate on the  $\text{CO}_2$  content of the depleted Siqueiros Fracture Zone melt inclusions. Our estimate is lower than the estimates of Marty and Tolstikhin (1998), Cartigny et al. (2008), Michael and Graham (2015), Le Voyer et al. (2017), and Hauri et al. (2017) based on a variety of approaches, but these prior studies utilize much more limited data sets than we have employed here. Our estimate is substantially higher than the study of Chavrit et al. (2014) who considered vesicle abundances in MORB; their underestimation is likely biased by  $\text{CO}_2$  contained in vesicles that were lost from samples by degassing. In general there is a good correlation of segment-scale  $\text{CO}_2$  flux with the segment-averaged primary  $\text{CO}_2$  content (Figure 9), while the correlation of  $\text{CO}_2$  flux with magma flux is much weaker.

The MORB flux of  $\text{CO}_2$  is nearly quantitatively delivered to the overlying water column, during eruptive degassing and later by hydrothermal circulation, where it becomes part of the marine bicarbonate cycle. Short-term global effects of large variations in the ridge  $\text{CO}_2$  flux are, however, muted by the  $\sim 100,000$ -

determined crustal thickness correlates well with mean ridge depth (Figure 7), reflecting isostatic compensation of MORB crust at the ridge. These constraints permit an estimation of the crustal thickness at all of the 711 segments defined by Gale et al. (2013) based on the axial depth. We then calculate a segment-averaged magma production rate by multiplying the crustal thickness by the mean segment spreading rate, which we then use to calculate the fluxes of  $\text{CO}_2$  (and other elements) on a segment-by-segment basis. There is one exception to the relationship between ridge depth and crustal thickness at the northernmost three segments of the Alarcón Ridge, where the ridge crest shallows as it begins to intersect the incipient continental rift in the Gulf of California; here we have chosen to assign an average crustal thickness of 6.4 km to these three ridge segments (EPRR1, EPRR2, and EPRR3).

Carbon dioxide fluxes, normalized by ridge length, vary by a factor of  $>100$  because they are the products of local primary magma  $\text{CO}_2$  contents and local magma production rates, which are uncorrelated. Fluxes of  $\text{CO}_2$  vary from a low of  $1.52 \times 10^6$  mol/year/km at segment SWIR17 (which is both depleted and slow-spreading) to a high of  $4.74 \times 10^8$  mol/year/km at segment JUAN1 (which has high primary  $\text{CO}_2$  and moderate spreading rate). High  $\text{CO}_2$  fluxes are independent of spreading rate, occurring at both slow- and fast-spreading ridge segments. High  $\text{CO}_2$  fluxes also occur on segments both near to and far from hotspots, and there is a clear tendency for estimated  $\text{CO}_2$  fluxes to increase toward hotspots (Figure 8), in concert with enrichment in highly incompatible elements (i.e., Ba, Rb). Provided that the mechanism of mantle enrichment for hotspots does not fractionate  $\text{CO}_2$  from Ba or Rb, then this observation supports our perspective that variations in the MORB primary magma  $\text{CO}_2$  content are mostly driven by variations in mantle composition rather than magma production dynamics or mantle temperature.

When summed over the length of the global mid-ocean ridge system, we calculate the global MORB magma production rate to be  $16.5 \text{ km}^3/\text{year}$ , significantly lower than the widely cited  $21 \text{ km}^3/\text{year}$  estimated by Crisp (1984); this difference in magma production rate alone accounts for 27% of our total flux calculation. The total flux of  $\text{CO}_2$ , summed over the widely varying fluxes at individual ridge segments (Figure 9), is  $1.32_{-0.85}^{+0.77} \times 10^{12}$  mol/year. Importantly, for individual ridge segments with no Ba or Rb data from which to constrain  $\text{CO}_2$ , we assume an average magmatic  $\text{CO}_2$  content calculated from the log-normal mean of those segments with Ba or Rb




**Figure 7.** Correlation between seismically determined ridge crest crustal thickness (Behn & Grove, 2015; Van Avendonk et al., 2017) and ridge depth (Gale et al., 2013). This relationship allows for an estimation of crustal thickness at each ridge segment.

year residence time of carbon dissolved in seawater (Broecker & Peng, 1986) and thus variations in the ridge  $\text{CO}_2$  flux are likely to influence short-term surface climate conditions only if large fluxes bypass the water column in areas where the ridge is shallow or exposed above sea level. On the longer timescale, the ridge  $\text{CO}_2$  flux is dwarfed by the deposition of carbonate sediments and organic carbon on the seafloor and formation of carbonate alteration veins within the crustal and mantle sections of the oceanic lithosphere, making the seafloor a net sink for carbon by a large factor (Alt & Shanks, 2003; Alt & Teagle, 1999; Hayes & Waldbauer, 2006; Kelemen & Manning, 2015). Much (but not all) of this carbon is present in forms, and at locations, where it can be returned to the deep crust and/or mantle in subduction zones.


## 8. Influence of Carbon on Mantle Melting

Several studies have demonstrated that the presence of carbonate in the mantle can strongly influence the depth of melt initiation beneath ridges (Dasgupta & Hirschmann, 2006, 2007; Dasgupta et al., 2013; Keller et al., 2017; Stagno et al., 2013), as incipient melting of carbonate may represent the first step in a polybaric process that ultimately results in eruption of MORB. To first order, the 2-order-of-magnitude range in primary MORB  $\text{CO}_2$  content must reflect a large variation in the carbon content of the MORB source mantle. Le Voyer et al. (2017) hypothesized how changes in the carbon content of the mantle could result in geophysical observations illuminating the depth beneath ridges where low, but possibly interconnected, melt fractions (near 0.05%; Minarik & Watson, 1995) could first be detected at constant potential temperature. Specifically, they showed that changing the  $\text{CO}_2$  content of relatively oxidized and anhydrous mantle from 20 ppm to 140 ppm increased the depth where interconnected melt ( $F \sim 0.05\%$ ) would exist from 75 km to 275 km. If a combination of seafloor seismic and electromagnetic methods could elucidate clearly such low melt fractions (Eilon & Abers, 2017; Gaillard et al., 2008; Sifré et al., 2014), then a test of this hypothesis might be possible by comparing the geophysical signals at ridge segments characterized by widely different primary magma  $\text{CO}_2$  contents as identified here.



**Figure 8.** Heat map diagram of segment-averaged  $\text{CO}_2$  flux (log scale, mol/year/km) versus distance to nearest hotspot (Gale et al., 2013); brighter color (yellow) scales with increasing density of data points. Individual segments with high primary  $\text{CO}_2$  contents are observed both near to and far from hotspots, but the data density suggest an increase of  $\text{CO}_2$  approaching hotspots at distances  $< 1,400$  km from the nearest hotspot.

This test is complicated, however, if carbonate is not the stable phase of carbon in the mantle at the depth of incipient melting. While direct analyses of MORBs and their peridotitic residues suggest that mantle oxygen fugacity ( $f_{\text{O}_2}$ ) is near the quartz-fayalite-magnetite buffer, where carbonate would be the stable phase (Bezos & Humler, 2005; Cottrell & Kelley, 2011; Birner et al., 2018), oxygen fugacity ( $f_{\text{O}_2}$ ) may decrease with depth in the convecting oceanic mantle, as observed in the cratonic xenolith record and supported by theory and experiment (Eggler & Baker, 1982; O'Neill et al., 1993; Ballhaus, 1995; Frost & McCammon, 2008; Rohrbach et al., 2007; Stagno and Frost, 2010; Stagno et al., 2013). It has been suggested that this decrease in mantle  $f_{\text{O}_2}$  would reduce carbonate to graphite, diamond, or metal alloy, and this would prevent carbon from fluxing the mantle. In this scenario, the existence of an interconnected melt network would not occur until upwelling mantle became oxidized enough to stabilize carbonate ( $\sim$  QFM-2) or reached the onset of silicate melting, (e.g., Stagno et al., 2013). However, even in the limit where the mantle is too reduced to stabilize carbonate, carbon would dissolve in silicate melts and become exhausted within the first 1–2% of melting for typical carbon concentrations that characterize the MORB mantle (20–200 ppm C; Le Voyer et al., 2017; Rosenthal et al., 2015; Saal et al., 2002). Thus, whether the mantle is reduced or oxidized, and regardless of whether carbon actively fluxes mantle melting or not, we expect carbon to behave like a highly incompatible element and produce the correlations



**Figure 9.** (a) Map of segment-averaged CO<sub>2</sub> fluxes, and (b) estimated primary magma CO<sub>2</sub> concentrations along the global mid-ocean ridge system. Both quantities are derived from segment-averaged Ba and Rb abundances and average CO<sub>2</sub>/Ba and CO<sub>2</sub>/Rb ratios (see text). For segments without Ba or Rb data, CO<sub>2</sub> fluxes in (a) were calculated from the global average CO<sub>2</sub> flux normalized by ridge length (mol/year/km) derived from segments with Ba or Rb data, multiplied by segment length, spreading rate, and crustal thickness (estimated from the relationship in Figure 7).

of C with Ba, Rb, Th, and Nb that characterize suites of undegassed melt inclusions for typical mantle carbon concentrations (cf. Hauri et al., 2017). That we observe carbon's incompatible behavior up to much higher carbon concentrations supports other inferences, based on MORB and peridotite analyses, that graphite is not stable at shallow depths beneath the ridge.

## 9. Conclusions

We have determined the concentration of CO<sub>2</sub> and other volatiles in 753 MORB glasses by SIMS from all the major ocean basins and, when combined with data from prior studies, these data provide global coverage of carbon fluxes from mid-ocean ridges. Assuming constant ratios of CO<sub>2</sub>/Ba and CO<sub>2</sub>/Rb, we estimate primary magma CO<sub>2</sub> concentrations that vary from 104 ppm to 1.90 wt%, MORB CO<sub>2</sub> fluxes that range from  $3.3 \times 10^7$  to  $4.0 \times 10^{10}$  mol/year at the scale of individual ridge segments, and a global integrated CO<sub>2</sub> flux of  $1.32_{-0.85}^{+0.77} \times 10^{12}$  mol/year, overlapping the global CO<sub>2</sub> flux estimated from depleted MORB ( $0.93 \pm 0.28 \times 10^{12}$  mol/year, Saal et al., 2002) but considerably lower than the flux estimated from E-MORB popping rock at 14°N on the mid-Atlantic Ridge ( $15 \times 10^{12}$  mol/year, Javoy & Pineau, 1991). The accuracy of the CO<sub>2</sub> fluxes determined here is due to consideration of the heterogeneity of trace element abundances in MORB along the global ridge system, with the assumption that our global CO<sub>2</sub>/Ba and CO<sub>2</sub>/Rb ratios are both representative and homogeneous in the MORB mantle.

The primary magma abundances, and segment-scale fluxes, of mid-ocean ridge CO<sub>2</sub> are heterogeneously distributed both along the ridge system and between major ocean basins. Areas of both high flux and high primary CO<sub>2</sub> are predicted at hotspots (Iceland, Azores, Bouvet, Afar, Reunion, Easter) and regionally along the Gakkel Ridge (Arctic). On the East Pacific Rise, these fluxes are correlated with deep areas of elevated mantle temperatures (Dalton et al., 2014) and high magma production rates. High primary CO<sub>2</sub> contents are, however, mainly predicted for areas of enriched mantle composition, which may be associated with long-term enrichment of the mantle as measured by radiogenic isotopes.

Large variations in primary MORB CO<sub>2</sub> imply similar large variations in mantle carbon content beneath ridges. If the upper mantle is sufficiently oxidized, these variations would result in large changes in the depth at which interconnected melt ( $F \sim 0.05\%$ ) would exist (from 75 to 275 km). Such large depth differences might be resolvable using geophysical methods by comparing ridge segments characterized by vastly different CO<sub>2</sub> contents as we have described here.

## Acknowledgments

We would like to dedicate this paper to our colleague and mentor, Erik Hauri. As co-authors, it was both an honor and a challenge to step into his shoes in order to finish this work after his passing. With a passion for scientific achievement, an always positive attitude, and an everlasting care for peers, Erik was a cornerstone of the Deep Carbon Observatory Initiative. His legacy work on carbon in the mantle will inspire the next generation of scientists to press farther. Many thanks to Jianhua Wang for expert assistance with temperamental instrumentation in the Carnegie SIMS lab, and to Tim Gooding and the Smithsonian Department of Mineral Sciences for assistance with sample preparation. We thank the geochemistry group at IPGP, University of Paris for sharing data and samples from the Pacific-Antarctic Ridge. We wish to thank the captain, crew, and scientific parties of the cruises PACANTARCTIC 1 and 2 cruises (PAR), the GIMNAUT, CD127, and KNOX11RR cruises (CIR), the PROTEA-5, MD34, AG22, and AG53 (SWIR), and the CHEPR and PANORAMA-1 cruises (EPR). We also acknowledge support from the U. California Ship Funds, and NSF grant OCE-0726573 (DRH). National Science Foundation grant OCE-1555523 provides support for the curation and distribution of geological samples at the University of Rhode Island. This is a contribution to the Deep Carbon Observatory.

## References

- Alt, J. C., & Shanks, W. C. (2003). Serpentinization of abyssal peridotites from the MARK area, mid-Atlantic ridge: Sulfur geochemistry and reaction modeling. *Geochimica et Cosmochimica Acta*, 67(4), 641–653. [https://doi.org/10.1016/S0016-7037\(02\)01142-0](https://doi.org/10.1016/S0016-7037(02)01142-0)
- Alt, J. C., & Teagle, D. (1999). The uptake of carbon during alteration of ocean crust. *Geochimica et Cosmochimica Acta*, 63(10), 1527–1535. [https://doi.org/10.1016/S0016-7037\(99\)00123-4](https://doi.org/10.1016/S0016-7037(99)00123-4)
- Arevalo, R. Jr., & McDonough, W. F. (2010). Chemical variations and regional diversity observed in MORB. *Chemical Geology*, 271(1-2), 70–85. <https://doi.org/10.1016/j.chemgeo.2009.12.013>
- Aster, E. M., Wallace, P. J., Moore, L. R., Watkins, J., Gazel, E., & Bodnar, R. J. (2016). Reconstructing CO<sub>2</sub> concentrations in basaltic melt inclusions using Raman analysis of vapor bubbles. *Journal of Volcanology and Geothermal Research*, 323, 148–162. <https://doi.org/10.1016/j.jvolgeores.2016.04.028>
- Bach, W., Hegner, E., Erzinger, J., & Satir, M. (1994). Chemical and isotopic variations along the superfast spreading East Pacific rise from 6°S to 30°S. *Contributions to Mineralogy and Petrology*, 116(4), 365–380. <https://doi.org/10.1007/BF00310905>
- Ballhaus, C. (1995). Is the upper mantle metal-saturated? *Earth and Planetary Science Letters*, 132(1–4), 75–86. [https://doi.org/10.1016/0012-821X\(95\)00047-G](https://doi.org/10.1016/0012-821X(95)00047-G)
- Barry, P. H. (2012). Laboratory and field-based instrumentation developments and noble gas-stable isotope systematics of Rungwe Volcanic Province, Iceland and the Central Indian Ridge (Doctoral dissertation). Location: Scripps Institution of Oceanography, University of California San Diego.
- Barry, P. H., Hilton, D. R., Fúri, E., Halldórsson, S. A., & Grönvold, K. (2014). Carbon isotope and abundance systematics, and CO<sub>2</sub> fluxes from Icelandic geothermal gases, fluids and subglacial basalts. *Geochimica et Cosmochimica Acta*, 134, 74–99. <https://doi.org/10.1016/j.gca.2014.02.038>
- Behn, M. D., & Grove, T. L. (2015). Melting systematics in mid-ocean ridge basalts: Application of a plagioclase-spinel melting model to global variations in major element chemistry and crustal thickness. *Journal of Geophysical Research: Solid Earth*, 120, 4863–4886. <https://doi.org/10.1002/2015JB011885>
- Bézos, A., & Humler, E. (2005). The Fe+3/ΣFe ratios of MORB glasses and their implications for mantle melting. *Geochimica et Cosmochimica Acta*, 63(3), 711–725. <https://doi.org/10.1016/j.gca.2004.07.026>
- Birner, S. K., Cottrell, E., Warren, J. M., Kelley, K. A., & Davis, F. A. (2018). Peridotites and basalts reveal broad congruence between two independent records of mantle  $f_{O_2}$  despite local redox heterogeneity. *Earth and Planetary Science Letters*, 494, 172–189. <https://doi.org/10.1016/j.epsl.2018.04.035>
- Broecker, W. S., & Peng, T.-H. (1986). Carbon cycle: 1985 glacial to interglacial changes in the operation of the global carbon cycle. *Radiocarbon*, 28(2A), 309–327. <https://doi.org/10.1017/S0033822200007414>
- Burley, J. M. A., & Katz, R. F. (2015). Variations in mid-ocean ridge CO<sub>2</sub> emissions driven by glacial cycles. *Earth and Planetary Science Letters*, 426, 246–258. <https://doi.org/10.1016/j.epsl.2015.06.031>
- Cartigny, P., Pineau, F., Aubaud, C., & Javoy, M. (2008). Towards a consistent mantle carbon flux estimate: Insights from volatile systematics (H<sub>2</sub>O/Ce, δD, CO<sub>2</sub>/Nb) in the North Atlantic mantle (14°N and 34°N). *Earth and Planetary Science Letters*, 265(3-4), 672–685. <https://doi.org/10.1016/j.epsl.2007.11.011>

- Chavrit, D., Humler, E., & Grasset, O. (2014). Mapping modern CO<sub>2</sub> fluxes and mantle carbon content all along the mid-ocean ridge system. *Earth and Planetary Science Letters*, 387, 229–239. <https://doi.org/10.1016/j.epsl.2013.11.036>
- Chavrit, D., Humler, E., Morizet, Y., & Laporte, D. (2012). Influence of magma ascent rate on carbon dioxide degassing at oceanic ridges: Message in a bubble. *Earth and Planetary Science Letters*, 357–358, 376–385. <https://doi.org/10.1016/j.epsl.2012.09.042>
- Class, C., & Lehnert, K. (2012). PetDB expert MORB (mid-ocean ridge basalt) compilation. *EarthChem Library*, 1–2. <https://doi.org/10.1594/IEDA/100060>
- Clog, M., Aubaud, C., Cartigny, P., & Dosso, L. (2013). The hydrogen isotopic composition and water content of southern Pacific MORB: A reassessment of the D/H ratio of the depleted mantle reservoir. *Earth and Planetary Science Letters*, 381, 156–165. <https://doi.org/10.1016/j.epsl.2013.08.043>
- Cottrell, E., & Kelley, K. A. (2011). The oxidation state of Fe in MORB glasses and the oxygen fugacity of the upper mantle. *Earth and Planetary Science Letters*, 305(3–4), 270–282. <https://doi.org/10.1016/j.epsl.2011.03.014>
- Cottrell, E., Kelley, K. A., Lanzirotti, A., & Fischer, R. A. (2009). High-precision determination of iron oxidation state in silicate glasses using XANES. *Chemical Geology*, 268(3–4), 167–179. <https://doi.org/10.1016/j.chemgeo.2009.08.008>
- Crisp, J. A. (1984). Rates of magma emplacement and volcanic output. *Journal of Volcanology and Geothermal Research*, 20(3–4), 177–211. [https://doi.org/10.1016/0377-0273\(84\)90039-8](https://doi.org/10.1016/0377-0273(84)90039-8)
- Dalton, C. A., Langmuir, C. H., & Gale, A. (2014). Geophysical and geochemical evidence for deep temperature variations beneath mid-ocean ridges. *Science*, 344, 80–83. <https://doi.org/10.1126/science.1249466>
- Dasgupta, R., & Hirschmann, M. M. (2006). Melting in the Earth's deep upper mantle caused by carbon dioxide. *Nature*, 440, 659–662. <https://doi.org/10.1038/nature04612>
- Dasgupta, R., & Hirschmann, M. M. (2007). Effect of variable carbonate concentration on the solidus of mantle peridotite. *American Mineralogist*, 92(2–3), 370–379. <https://doi.org/10.2138/am.2007.2201>
- Dasgupta, R., Mallik, A., Tsuno, K., Withers, A. C., Hirth, G., & Hirschmann, M. M. (2013). Carbon-dioxide-rich silicate melt in the Earth's upper mantle. *Nature*, 493(7431), 211–215. <https://doi.org/10.1038/nature11731>
- Dixon, J. E., Bindeman, I. N., Kingsley, R. H., Simons, K. K., LeRoux, P. J., Hajewski, T. R., et al. (2017). Light stable isotopic compositions of enriched mantle sources: Resolving the dehydration paradox. *Geochemistry, Geophysics, Geosystems*, 18, 3801–3839. <https://doi.org/10.1002/2016GC006743>
- Dixon, J. E., & Stolper, E. M. (1995). An experimental study of water and carbon dioxide solubilities in mid-ocean ridge basaltic liquids. Part II: Applications to degassing. *Journal of Petrology*, 36, 1633–1646.
- Dixon, J. E., Stolper, E. M., & Holloway, J. R. (1995). An experimental study of water and carbon dioxide solubilities in mid-ocean ridge basaltic liquids. Part I: Calibration and solubility models. *Journal of Petrology*, 36, 1607–1631.
- Donnelly, K. E., Goldstein, S. L., Langmuir, C. H., & Spiegelman, M. (2004). Origin of enriched ocean ridge basalts and implications for mantle dynamics. *Earth and Planetary Science Letters*, 226(3–4), 347–366. <https://doi.org/10.1016/j.epsl.2004.07.019>
- Eggler, D. H., & Baker, D. R. (1982). Reduced volatiles in the system C-O-H: Implications to mantle melting, fluid formation and diamond genesis. In S. Akimoto & M. H. Manghnani (Eds.), *High-Pressure Research in Geophysics* (pp. 237–250). Tokyo: Center for Academic Publications. [https://doi.org/10.1007/978-94-009-7867-6\\_19](https://doi.org/10.1007/978-94-009-7867-6_19)
- Eilon, Z. C., & Abers, G. A. (2017). High seismic attenuation at a mid-ocean ridge reveals the distribution of deep melt. *Science Advances*, 3(5), e1602829. <https://doi.org/10.1126/sciadv.1602829>
- Frost, D. J., & McCammon, C. A. (2008). The redox state of Earth's mantle. *Annual Review of Earth and Planetary Sciences*, 36, 389–420. <https://doi.org/10.1146/Annurev.Earth.36.031207.124322>
- Füri, E., Hilton, D. R., Murton, B. J., Hemond, C., Dymert, J., & Day, J. M. D. (2011). Helium isotope variations between Réunion Island and the Central Indian Ridge (17°–21°S): New evidence for ridge-hot spot interaction. *Journal of Geophysical Research*, 116, B02207. <https://doi.org/10.1029/2010JB007609>
- Gaillard, F., Malki, M., Iacono-Marziano, G., Pichavant, M., & Scaillet, B. (2008). Carbonatite melts and electrical conductivity in the asthenosphere. *Science*, 322(5906), 1363–1365. <https://doi.org/10.1126/science.1164446>
- Gale, A., Escrig, S., Gier, E. J., Langmuir, C. H., & Goldstein, S. L. (2011). Enriched basalts at segment centers: The Lucky Strike (37°17'N) and Menez Gwen (37°50'N) segments of the Mid-Atlantic Ridge. *Geochemistry, Geophysics, Geosystems*, 12, Q06016. <https://doi.org/10.1029/2010GC003446>
- Gale, A., Dalton, C. A., Langmuir, C. H., Su, Y., & Schilling, J.-G. (2013). The mean composition of ocean ridge basalts. *Geochemistry, Geophysics, Geosystems*, 14, 489–518. <https://doi.org/10.1029/2012GC004334>
- Gale, A., Laubier, M., Escrig, S., & Langmuir, C. H. (2013). Constraints on melting processes and plume-ridge interaction from comprehensive study of the FAMOUS and north FAMOUS segments, Mid-Atlantic Ridge. *Earth and Planetary Science Letters*, 365, 209–220.
- Gale, A., Langmuir, C. H., & Dalton, C. A. (2014). The global systematics of ocean ridge basalts and their origin. *Journal of Petrology*, 55(6), 1051–1082. <https://doi.org/10.1093/petrology/egu017>
- Ghiorso, M. S., & Gualda, G. A. R. (2015). An H<sub>2</sub>O–CO<sub>2</sub> mixed fluid saturation model compatible with rhyolite-MELTS. *Contributions to Mineralogy and Petrology*, 169(6), 1–30. <https://doi.org/10.1007/s00410-015-1141-8>
- Hamelin, B., & Allègre, C. J. (1985). Large-scale regional units in the depleted upper mantle revealed by an isotope study of the southwest Indian ridge. *Nature*, 315(6016), 196–199. <https://doi.org/10.1038/315196a0>
- Hamelin, C., Dosso, L., Hanan, B., Barrat, J.-A., & Ondréas, H. (2010). Sr-Nd-Hf isotopes along the Pacific Antarctic Ridge from 41 to 53°S. *Geophysical Research Letters*, 37, L10303. <https://doi.org/10.1029/2010GL042979>
- Hauri, E., Wang, J., Dixon, J. E., King, P. L., & Mandeville, C. (2002). SIMS analysis of volatiles in silicate glasses 1. Calibration, matrix effects and comparisons with FTIR. *Chemical Geology*, 183(1–4), 99–114. [https://doi.org/10.1016/S0009-2541\(01\)00375-8](https://doi.org/10.1016/S0009-2541(01)00375-8)
- Hauri, E. H., Maclennan, J., McKenzie, D., Gronvold, K., Oskarsson, N., & Shimizu, N. (2017). CO<sub>2</sub> content beneath northern Iceland and the variability of mantle carbon. *Geology*, 46(1), 55–58. <https://doi.org/10.1130/G39413.1>
- Hayes, J. M., & Waldbauer, J. R. (2006). The carbon cycle and associated redox processes through time. *Philosophical Transactions of the Royal Society, B: Biological Sciences*, 59(7), 1217–1232. [https://doi.org/10.1016/0016-7037\(95\)00038-2](https://doi.org/10.1016/0016-7037(95)00038-2)
- Helo, C., Longpre, M. A., Shimizu, N., Clague, D. A., & Stix, J. (2011). Explosive eruptions at mid-ocean ridges driven by CO<sub>2</sub>-rich magmas. *Nature Geoscience*, 4(4), 260–263. <https://doi.org/10.1038/NGEO1104>
- Iacono-Marziano, G., Morizet, Y., Trong, E. L., & Gaillard, F. (2012). New experimental data and semi-empirical parameterization of H<sub>2</sub>O–CO<sub>2</sub> solubility in mafic melts. *Geochimica et Cosmochimica Acta*, 97, 1–23. <https://doi.org/10.1016/j.gca.2012.08.035>
- Janney, P. E., Le Roex, A. P., & Carlson, R. W. (2005). Hafnium isotope and trace element constraints on the nature of mantle heterogeneity beneath the central southwest Indian ridge (13°E to 47°E). *Journal of Petrology*, 46(12), 2427–2464. <https://doi.org/10.1093/petrology/egi060>

- Javoy, M., & Pineau, F. (1991). The volatiles record of a popping rock from the mid-Atlantic ridge at 14°N - chemical and isotopic composition of gas trapped in the vesicles. *Earth and Planetary Science Letters*, 107(3-4), 598–611. [https://doi.org/10.1016/0012-821X\(91\)90104-P](https://doi.org/10.1016/0012-821X(91)90104-P)
- Jendrzewski, N., Trull, T. W., Pineau, F., & Javoy, M. (1997). Carbon solubility in mid-ocean ridge basaltic melt at low pressures (250-1950 bar). *Chemical Geology*, 138(1-2), 81–92. [https://doi.org/10.1016/S0009-2541\(96\)00176-3](https://doi.org/10.1016/S0009-2541(96)00176-3)
- Jenner, F. E., Hauri, E. H., Bullock, E. S., König, S., Arculus, R. J., Mavrogenes, J. A., et al. (2015). The competing effects of sulfide saturation versus degassing on the behavior of the chalcophile elements during the differentiation of hydrous melts. *Geochemistry, Geophysics, Geosystems*, 16, 1490–1507. <https://doi.org/10.1002/2014GC005670>
- Jenner, F. E., & O'Neill, H. S. C. (2012). Analysis of 60 elements in 616 ocean floor basaltic glasses. *Geochemistry, Geophysics, Geosystems*, 13, Q02005. <https://doi.org/10.1029/2011GC004009>
- Keith, M., Haase, K. M., Klemm, R., Schwarz-Schampera, U., & Franke, H. (2017). Systematic variations in magmatic sulphide chemistry from mid-ocean ridges, back-arc basins and island arcs. *Chemical Geology*, 451, 67–77. <https://doi.org/10.1016/j.chemgeo.2016.12.028>
- Kelemen, P. B., & Manning, C. E. (2015). Reevaluating carbon fluxes in subduction zones, what goes down, mostly comes up. *Proceedings of the National Academy of Sciences*, 112(30), E3997–E4006. <https://doi.org/10.1073/pnas.1507889112>
- Keller, T., Katz, R. F., & Hirschmann, M. M. (2017). Volatiles beneath mid-ocean ridges: Deep melting, channelised transport, focusing, and metasomatism. *Earth and Planetary Science Letters*, 464, 55–68. <https://doi.org/10.1016/j.epsl.2017.02.006>
- Kelley, K. A., Kingsley, R., & Schilling, J.-G. (2013). Composition of plume-influenced mid-ocean ridge lavas and glasses from the mid-Atlantic Ridge, East Pacific rise, Galapagos spreading center, and Gulf of Aden. *Geochemistry, Geophysics, Geosystems*, 14, 223–242. <https://doi.org/10.1029/2012GC004415>
- Kinzler, R. J., & Grove, T. L. (1992). Primary magmas of mid-ocean ridge basalts 1. Experiments and methods. *Journal of Geophysical Research*, 97(B5), 6885–6906. <https://doi.org/10.1029/91JB02840>
- Klein, E. M., & Langmuir, C. H. (1987). Global correlations of ocean ridge basalt chemistry with axial depth and crustal thickness. *Journal of Geophysical Research*, 92(B8), 8089–8115. <https://doi.org/10.1029/JB092iB08p08089>
- Labidi, J., Cartigny, P., Hamelin, C., Moreira, M., & Dosso, L. (2014). Sulfur isotope budget (<sup>32</sup>S, <sup>33</sup>S, <sup>34</sup>S and <sup>36</sup>S) in Pacific-Antarctic ridge basalts: A record of mantle source heterogeneity and hydrothermal sulfide assimilation. *Geochimica et Cosmochimica Acta*, 133, 47–67. <https://doi.org/10.1016/j.gca.2014.02.023>
- Langmuir, C. H. (2014). *Major and trace element composition of basalt glasses from the CHEPR Cruise, East Pacific Rise*. Interdisciplinary Earth Data Alliance (IEDA). <https://doi.org/10.1594/IEDA/100441>
- Langmuir, C. H., Klein, E. M., & Plank, T. (1992). Petrological systematics of mid-ocean ridge basalts: Constraints on melt generation beneath ocean ridges. In J. P. Morgan, D. K. Blackman, & J. M. Sinton (Eds.), *Mantle flow and melt generation at mid-ocean ridges, geophysics monogram series* (Vol. 71, pp. 183–280). Washington, DC: American Geophysical Union. <https://doi.org/10.1029/GM071p0183>
- Le Roex, A. P., Dick, H., & Fisher, R. L. (1989). Petrology and geochemistry of MORB from 25°E to 46°E along the southwest Indian Ridge—Evidence for contrasting styles of mantle enrichment. *Journal of Petrology*, 30(4), 947–986. <https://doi.org/10.1093/petrology/30.4.947>
- Le Roux, P. J., Shirey, S. B., Hauri, E. H., & Perfit, M. R. (2006). The effects of variable sources, processes and contaminants on the composition of northern EPR MORB (8–10°N and 12–14°N): Evidence from volatiles (H<sub>2</sub>O, CO<sub>2</sub>, S) and halogens (F, Cl). *Earth and Planetary Science Letters*, 251(3-4), 209–231. <https://doi.org/10.1016/j.epsl.2006.09.012>
- Le Roux, P. J., Le Roex, A. P., & Schilling, J.-G. (2002a). Crystallization processes beneath the southern Mid-Atlantic Ridge (40–55°S): Evidence for high-pressure initiation of crystallization. *Contributions to Mineralogy and Petrology*, 142, 582–602.
- Le Roux, P. J., Le Roex, A. P., & Schilling, J.-G. (2002b). MORB melting processes beneath the southern Mid-Atlantic Ridge (40–55°S): A role for pyroxenite and residual garnet. *Contributions to Mineralogy and Petrology*, 144, 206–229.
- Le Voyer, M., Cottrell, E., Kelley, K. A., Brounce, M., & Hauri, E. H. (2015). The effect of primary versus secondary processes on the volatile content of MORB glasses: An example from the equatorial mid-Atlantic ridge (5°N–3°S). *Journal of Geophysical Research: Solid Earth*, 120, 25–144. <https://doi.org/10.1002/2014JB011160>
- Le Voyer, M., Kelley, K. A., Cottrell, E., & Hauri, E. H. (2017). Heterogeneity in mantle carbon content from CO<sub>2</sub>-undersaturated basalts. *Nature Communications*, 8, 1–8. <https://doi.org/10.1038/ncomms14062>
- Mahoney, J. J., Le Roex, A. P., Peng, Z., Fisher, R. L., & Natland, J. H. (1992). Southwestern limits of Indian Ocean ridge mantle and the origin of low <sup>206</sup>Pb/<sup>204</sup>Pb mid-ocean ridge basalt: Isotope systematics of the central southwest Indian ridge (17°–50°E). *Journal of Geophysical Research*, 97(B13), 19,771–19,790. <https://doi.org/10.1029/92JB01424>
- Marty, B., Alexander, C. M. O'D., & Raymond, S. N. (2013). Primordial origins of Earth's carbon. *Reviews in Mineralogy and Geochemistry*, 75(1), 149–181. <https://doi.org/10.2138/rmg.2013.75.6>
- Marty, B., & Tolstikhin, I. N. (1998). CO<sub>2</sub> fluxes from mid-ocean ridges, arcs and plumes. *Chemical Geology*, 145(3-4), 233–248. [https://doi.org/10.1016/S0009-2541\(97\)00145-9](https://doi.org/10.1016/S0009-2541(97)00145-9)
- Matthews, S., Shorttle, O., Rudge, J. F., & MacLennan, J. (2017). Constraining mantle carbon: CO<sub>2</sub>-trace element systematics in basalts and the roles of magma mixing and degassing. *Earth and Planetary Science Letters*, 480, 1–14. <https://doi.org/10.1016/j.epsl.2017.09.047>
- Menke, W. (1999). Crustal isostasy indicates anomalous densities beneath Iceland. *Geophysical Research Letters*, 26(9), 1215–1218. <https://doi.org/10.1029/1999GL900202>
- Michael, P. (1995). Regionally distinctive sources of depleted MORB: Evidence from trace elements and H<sub>2</sub>O. *Earth and Planetary Science Letters*, 131(3-4), 301–320. [https://doi.org/10.1016/0012-821X\(95\)00023-6](https://doi.org/10.1016/0012-821X(95)00023-6)
- Michael, P. J., & Graham, D. W. (2015). The behavior and concentration of CO<sub>2</sub> in the suboceanic mantle: Inferences from undegassed ocean ridge and ocean island basalts. *Lithos*, 236-237, 338–351. <https://doi.org/10.1016/j.lithos.2015.08.020>
- Michael, P. J., & Schilling, J.-G. (1989). Chlorine in mid-ocean ridge magmas: Evidence for assimilation of seawater-influenced components. *Geochimica et Cosmochimica Acta*, 53(12), 3131–3143. [https://doi.org/10.1016/0016-7037\(89\)90094-X](https://doi.org/10.1016/0016-7037(89)90094-X)
- Minarik, W. G., & Watson, E. B. (1995). Interconnectivity of carbonate melt at low melt fraction. *Earth and Planetary Science Letters*, 133(3-4), 423–437. [https://doi.org/10.1016/0012-821X\(95\)00085-Q](https://doi.org/10.1016/0012-821X(95)00085-Q)
- Moore, L. R., Gazel, E., Tuohy, R., Lloyd, A. S., Esposito, R., Steele-MacLennan, M., et al. (2015). Bubbles matter: An assessment of the contribution of vapor bubbles to melt inclusion volatile budgets. *American Mineralogist*, 100(4), 806–823. <https://doi.org/10.2138/am-2015-5036>
- Moreira, M. A., Dosso, L., & Ondreas, H. (2008). Helium isotopes on the Pacific-Antarctic ridge (52.5–41.5 °S). *Geophysical Research Letters*, 35, L10306. <https://doi.org/10.1029/2008GL033286>
- Murton, B. J., Tindle, A. G., Milton, J. A., & Sauter, D. (2005). Heterogeneity in southern Central Indian Ridge MORB: Implications for ridge-hot spot interaction. *Geochemistry, Geophysics, Geosystems*, 6, Q03E20. <https://doi.org/10.1029/2004GC000798>



- Nauret, F., Abouchami, W., Galer, S. J. G., Hofmann, A. W., Hémond, C., Chauvel, C., & Dymant, J. (2006). Correlated trace element-Pb isotope enrichments in Indian MORB along 18–20°S, Central Indian Ridge. *Earth and Planetary Science Letters*, 245(1-2), 137–152. <https://doi.org/10.1016/j.epsl.2006.03.015>
- O'Neill, H. S. C., & Mavrogenes, J. (2002). The sulfide capacity and the sulfur content at sulfide saturation of silicate melts at 1400 °C and 1 bar. *Journal of Petrology*, 43(6), 1049–1087. <https://doi.org/10.1093/petrology/43.6.1049>
- O'Neill, H. S. C., Rubie, D. C., Canil, D., Geiger, C. A., Ross, C. R., Seifert, F., & Woodland, A. B. (1993). Ferric iron in the upper mantle and in transition zone assemblages: Implications for relative oxygen fugacities in the mantle. In E. Takahashi, R. Jeanloz, & D. C. Rubie (Eds.), *Evolution of earth and planets, geophysical monograph series* (Vol. 74, pp. 73–88). Washington, DC: International Union of Geodesy and Geophysics/American Geophysical Union.
- Papale, P., Moretti, R., & Barbato, D. (2006). The compositional dependence of the saturation surface of H<sub>2</sub>O+CO<sub>2</sub> fluids in silicate melts. *Chemical Geology*, 229(1-3), 78–95. <https://doi.org/10.1016/j.chemgeo.2006.01.013>
- Resing, J. A., Lupton, J. E., Feely, R. A., & Lilley, M. D. (2004). CO<sub>2</sub> and <sup>3</sup>He in hydrothermal plumes: Implications for mid-ocean ridge CO<sub>2</sub> flux. *Earth and Planetary Science Letters*, 226(3-4), 449–464. <https://doi.org/10.1016/j.epsl.2004.07.028>
- Robock, A. (2000). Volcanic eruptions and climate. *Reviews of Geophysics*, 38(2), 191–219. <https://doi.org/10.1029/1998RG000054>
- Rohrbach, A., Ballhaus, C., Golla-Schindler, U., Ulmer, P., Kamenetsky, V. S., & Kuzmin, D. V. (2007). Metal saturation in the upper mantle. *Nature*, 449(7161), 456–458. <https://doi.org/10.1038/Nature06183>
- Rosenthal, A., Hauri, E. H., & Hirschmann, M. M. (2015). Experimental determination of C, F and H partitioning between mantle minerals and carbonated basalt; CO<sub>2</sub>/Ba and CO<sub>2</sub>/Nb systematics of partial melting, and the CO<sub>2</sub> contents of basaltic source regions. *Earth and Planetary Science Letters*, 412, 77–87. <https://doi.org/10.1016/j.epsl.2014.11.044>
- Rowe, E. C., & Schilling, J. G. (1979). Fluorine in Iceland and Reykjanes ridge basalts. *Nature*, 279(5708), 33–37. <https://doi.org/10.1038/279033a0>
- Saal, A. E., Hauri, E. H., Langmuir, C. H., & Perfit, M. R. (2002). Vapour undersaturation in primitive mid-ocean-ridge basalt and the volatile content of Earth's upper mantle. *Nature*, 419(6906), 451–455. <https://doi.org/10.1038/nature01073>
- Salters, V. J. M., Mallick, S., Hart, S. R., Langmuir, C. E., & Stracke, A. (2011). Domains of depleted mantle: New evidence from hafnium and neodymium isotopes. *Geochemistry, Geophysics, Geosystems*, 12, Q08001. <https://doi.org/10.1029/2011GC003617>
- Schilling, J. G., Zajac, M., Evans, R., Johnston, T., White, W., Devine, J. D., & Kingsley, R. (1983). Petrologic and geochemical variations along the mid-Atlantic ridge from 29°N to 73°N. *American Journal of Science*, 283(6), 510–586. <https://doi.org/10.2475/ajs.283.6.510>
- Shaw, A. M., Behn, M. D., Humphris, S. E., Sohn, R. A., & Gregg, P. M. (2010). Deep pooling of low degree melts and volatile fluxes at the 85°E segment of the Gakkel ridge: Evidence from olivine-hosted melt inclusions and glasses. *Earth and Planetary Science Letters*, 289(3-4), 311–322. <https://doi.org/10.1016/j.epsl.2009.11.018>
- Shimizu, K., Saal, A. E., Myers, C. E., Nagle, A. N., Hauri, E. H., Forsyth, D. W., et al. (2016). Two-component mantle melting-mixing model for the generation of mid-ocean ridge basalts: Implications for the volatile content of the Pacific upper mantle. *Geochimica et Cosmochimica Acta*, 176, 44–80. <https://doi.org/10.1016/j.gca.2015.10.033>
- Shindell, D. T., Schmidt, G. A., Miller, R. L., & Mann, M. E. (2003). Volcanic and solar forcing of climate change during the preindustrial era. *Journal of Climate*, 16(24), 4094–4107. [https://doi.org/10.1175/1520-0442\(2003\)016<4094:VASFOC>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<4094:VASFOC>2.0.CO;2)
- Shishkina, T. A., Botcharnikov, R. E., Holtz, F., Almeev, R. R., & Portnyagin, M. V. (2010). Solubility of H<sub>2</sub>O- and CO<sub>2</sub>-bearing fluids in tholeiitic basalts at pressures up to 500MPa. *Chemical Geology*, 277(1-2), 115–125. <https://doi.org/10.1016/j.chemgeo.2010.07.014>
- Sifré, D., Gardés, E., Massuyeau, M., Hashim, L., Hier-Majumder, S., & Gaillard, F. (2014). Electrical conductivity during incipient melting in the oceanic low-velocity zone. *Nature*, 509(7498), 81–85. <https://doi.org/10.1038/nature13245>
- Stagno, V., & Frost, D. J. (2010). Carbon speciation in the asthenosphere: Experimental measurements of the redox conditions at which carbonate-bearing melts coexist with graphite or diamond in peridotite assemblages. *Earth and Planetary Science Letters*, 300(1-2), 72–84. <https://doi.org/10.1016/j.epsl.2010.09.038>
- Stagno, V., Ojwang, D. O., McCammon, C. A., & Frost, D. J. (2013). The oxidation state of the mantle and the extraction of carbon from Earth's interior. *Nature*, 493(7430), 84–88. <https://doi.org/10.1038/nature11679>
- Tolstoy, M. (2015). Mid-ocean ridge eruptions as a climate valve. *Geophysical Research Letters*, 42, 1346–1351. <https://doi.org/10.1002/2014GL063015>
- University of Rhode Island (URI) Graduate School of Oceanography (GSO) Marine Geological Samples Laboratory (MGSL) (2018). Archive of URI geosample data and information. NOAA National Centers for Environmental Information. <https://doi.org/10.7289/V5JQZ0W>
- Van Avendonk, H. J., Davis, J. K., Harding, J. L., & Lawver, L. A. (2017). Decrease in oceanic crustal thickness since the breakup of Pangaea. *Nature Geoscience*, 10(1), 58–61. <https://doi.org/10.1038/ngeo2849>
- Vlastelic, I., Aslanian, D., Dosso, L., Bougault, H., Olivet, J. L., & Geli, L. (1999). Large-scale chemical and thermal division of the Pacific mantle. *Nature*, 399(6734), 345–350. <https://doi.org/10.1038/20664>
- Vlastelic, I., Dosso, L., Bougault, H., Aslanian, D., Geli, L., Etoubleau, J., et al. (2000). Chemical systematics of an intermediate spreading ridge: The Pacific-Antarctic ridge between 56 °S and 66 °S. *Journal of Geophysical Research*, 105(B2), 2915–2936. <https://doi.org/10.1029/1999JB900234>
- Wanless, V. D., Behn, M. D., Shaw, A. M., & Plank, T. (2014). Variations in melting dynamics and mantle compositions along the eastern volcanic zone of the Gakkel Ridge: Insights from olivine-hosted melt inclusions. *Contributions to Mineralogy and Petrology*, 167(5), 1005–1027. <https://doi.org/10.1007/s00410-014-1005-7>
- Wanless, V. D., & Shaw, A. M. (2012). Lower crustal crystallization and melt evolution at mid-ocean ridges. *Nature Geoscience*, 5(9), 651–655. <https://doi.org/10.1038/ngeo1552>
- Wanless, V. D., Shaw, A. M., Behn, M. D., Soule, S. A., Escartin, J., & Hamelin, C. (2015). Magmatic plumbing at Lucky Strike volcano based on olivine-hosted melt inclusion compositions. *Geochemistry, Geophysics, Geosystems*, 16, 126–147. <https://doi.org/10.1002/2014GC005517>
- White, R. S., McKenzie, D., & O'Nions, R. K. (1992). Oceanic crustal thickness from seismic measurements and rare earth element inversions. *Journal of Geophysical Research*, 97(B13), 19,683–19,715. <https://doi.org/10.1029/92JB01749>
- Zhang, H. L., Cottrell, E., Solheid, P. A., Kelley, K. A., & Hirschmann, M. M. (2018). Determination of Fe<sup>3+</sup>/ΣFe of XANES basaltic glass standards by Mössbauer spectroscopy and its application to the oxidation state of iron in MORB. *Chemical Geology*, 479, 166–175. <https://doi.org/10.1016/j.chemgeo.2018.01.006>
- Zhong, Y., Miller, G. H., Otto-Bliesner, B. L., Holland, M. M., Bailey, D. A., Schneider, D. P., & Geirsdottir, A. (2010). Centennial-scale climate change from decadal-paced explosive volcanism: A coupled sea ice-ocean mechanism. *Climate Dynamics*, 37(11-12), 2373–2387. <https://doi.org/10.1007/s00382-010-0967-z>

## References from the supporting information

- Agar, S. M., Casey, J. F., & Kempton, P. D. (1997). Textural, geochemical, and isotopic variations in gabbroic shear zones from the MARK area. In J. A. Karson, M. Cannat, D. J. Miller, & D. Elthon (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 153, pp. 99–121). College Station, Texas: Ocean Drilling Program.
- Aggrey, K. E., Muenow, D. W., & Batiza, R. (1988). Volatile abundances in basaltic glasses from seamounts flanking the East Pacific Rise at 21°N and 12–14°N. *Geochimica et Cosmochimica Acta*, 52, 2115–2119.
- Aggrey, K. E., Muenow, D. W., & Sinton, J. M. (1988). Volatile abundances in submarine glasses from the north Fiji and Lau back-arc basins. *Geochimica et Cosmochimica Acta*, 52, 2501–2506.
- Agranier, A., Blichert-Toft, J., Graham, D. W., Debaille, V., Schiano, P., & Albarede, F. (2005). The spectra of isotopic heterogeneities along the Mid-Atlantic Ridge. *Earth and Planetary Science Letters*, 238, 96–109.
- Allan, J. F. (1995). Data report: Trace element geochemistry of Leg 142 basalts by instrumental neutron activation analysis. In R. Batiza, M. A. Storms, & J. F. Allan (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 142, pp. 87–89). College Station, TX: Ocean Drilling Program.
- Allan, J. F., Batiza, R., & Lonsdale, P. F. (1987). Petrology and chemistry of lavas from seamounts flanking the East Pacific Rise axis, 21°N: Implications concerning the mantle source composition for both seamount and adjacent EPR lavas. In B. H. Keating, P. Fryer, R. Batiza, & G. W. Boechler (Eds.), *Seamounts, Islands, and Atolls, Geophysical Monograph Series* (Vol. 43, pp. 255–282). Washington, DC: American Geophysical Union.
- Allan, J. F., Batiza, R., Perfit, M. R., Fornari, D. J., & Sack, R. O. (1989). Petrology of lavas from the Lamont Seamount chain and adjacent East Pacific Rise, 10°N. *Journal of Petroleum*, 30, 1245–1298.
- Allan, J. F., Batiza, R., & Sack, R. O. (1994). Geochemical characteristics of Cocos Plate Seamount lavas. *Contributions to Mineralogy and Petrology*, 116, 47–61.
- Allan, J. F., Falloon, T. J., Pedersen, R. B., Lakkapragada, B. S., Natland, J. H., & Malpas, J. (1996). Petrology of selected Leg 147 basaltic lavas and dikes. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 147, pp. 173–186). College Station, Texas: Ocean Drilling Program.
- Altherr, R., Henjes-Kunst, F., & Baumann, A. (1990). Asthenosphere versus lithosphere as possible sources for basaltic magmas erupted during formation of the red sea: Constraints from Sr, Pb and Nd isotopes. *Earth and Planetary Science Letters*, 96, 269–286.
- Altherr, R., Henjes-Kunst, F., Puchelt, H., & Baumann, A. (1988). Volcanic activity in the Red Sea axial trough—evidence for a large mantle diapir? *Tectonophysics*, 150, 121–133.
- Anderson, R. N., Clague, D. A., Klitgord, K. D., Marshall, M., & Nishimori, R. K. (1975). Magnetic and petrologic variations along the Galapagos Spreading Center and their relation to the Galapagos melting anomaly. *Geological Society of America Bulletin*, 86, 683–694.
- Anderson, R. N., Sparious, J., Weissel, J. K., & Hayes, D. E. (1980). The interrelation between variations in magnetic anomaly amplitudes and basaltic magnetization and chemistry along the southeast Indian Ridge. *Journal of Geophysical Research*, 85, 3883–3898. <https://doi.org/10.1029/JB085iB07p03883>
- Andres, M., Blichert-Toft, J., & Schilling, J.-G. (2002). Hafnium isotopes in basalts from the southern Mid-Atlantic Ridge from 40°S to 55°S: Discovery and Shona plume-ridge interactions and the role of recycled sediments. *Geochemistry, Geophysics, Geosystems*, 3(10), 8502. <https://doi.org/10.1029/2002GC000324>
- Andres, M., Blichert-Toft, J., & Schilling, J.-G. (2004). Nature of the depleted upper mantle beneath the Atlantic: Evidence from Hf isotopes in normal Mid-Ocean Ridge basalts from 79°N to 55°S. *Earth and Planetary Science Letters*, 225, 89–103.
- Antonini, P., Petrini, R., & Contin, G. (1998). A segment of sea-floor spreading in the central Red Sea: Basalts from the Nereus Deep (23° 00' - 23° 20'N). *Journal of African Earth Sciences*, 27, 107–114.
- Arcyana (1977). Rocks collected by bathyscaph and diving saucer in the FAMOUS area of the Mid-Atlantic Rift Valley: Petrological diversity and structural setting. *Deep Sea Research*, 24, 565–589.
- Arevalo, R. Jr., & McDonough, W. F. (2008). Tungsten geochemistry and implications for understanding the Earth's interior. *Earth and Planetary Science Letters*, 272, 656–665.
- Arevalo, R. Jr., McDonough, W. F., & Luong, M. (2009). The K/U ratio of the silicate Earth: Insights into mantle composition, structure and thermal evolution. *Earth and Planetary Science Letters*, 278, 361–369.
- Aumento, F. (1968). The Mid-Atlantic Ridge near 45°N II. Basalts from the area of Confederation Peak. *Canadian Journal of Earth Sciences*, 5, 1–21.
- Aumento, F., & Kempe, D. R. C. (1977). Geochemistry, normative mineralogy, and differentiation trends of basalt glasses from DSDP Leg 37. *Initial Reports of the Deep Sea Drilling Project*, 37, 729–734.
- van Avendonk, H. J., Davis, J. K., Harding, J. L., & Lawver, L. A. (2017). Decrease in oceanic crustal thickness since the breakup of Pangaea. *Nature Geoscience*, 10(1), 58–61.
- Bach, W., Alt, J. C., Niu, Y., Humphris, S. E., Erzinger, J., & Dick, H. J. B. (2001). The geochemical consequences of late-stage low-grade alteration of lower ocean crust at the SW Indian Ridge: Results from ODP Hole 735B (Leg 176). *Geochimica et Cosmochimica Acta*, 65, 3267–3287.
- Bach, W., & Erzinger, J. (1995). Volatile components in basalts and basaltic glasses from the EPR at 9°30'N. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 142, pp. 23–29). College Station, Texas: Ocean Drilling Program.
- Bach, W., Erzinger, J., Dosso, L., Bollinger, C., Bougault, H., Etoubleau, J., & Sauerwein, J. (1996). Unusually large Nb-Ta depletions in North Chile Ridge basalts at 36°50' to 38°56'S: Major element, trace element, and isotopic data. *Earth and Planetary Science Letters*, 142, 223–240.
- Bach, W., Hegner, E., Erzinger, J., & Satir, M. (1994). Chemical and isotopic variations along the superfast spreading East Pacific Rise from 6 to 30° S. *Contributions to Mineralogy and Petrology*, 116, 365–380.
- Barker, A. K., Coogan, L. A., & Gillis, K. M. (2010). Insights into the behaviour of sulphur in Mid-Ocean Ridge axial hydrothermal systems from the composition of the sheeted dyke complex at Pito Deep. *Chemical Geology*, 275, 105–115.
- Barker, A. K., Coogan, L. A., Gillis, K. M., & Weis, D. (2008). Strontium isotope constraints on fluid flow in the sheeted dike complex of fast spreading crust: Pervasive fluid flow at Pito Deep. *Geochemistry, Geophysics, Geosystems*, 9, Q06010. <https://doi.org/10.1029/2007GC001901>
- Barnes, J. D., & Straub, S. (2010). Chlorine stable isotope variations in Izu Bonin tephra: Implications for serpentinite subduction. *Chemical Geology*, 272, 62–74.
- Barrat, J. A., Jahn, B. M., Fourcade, S., & Joron, J.-L. (1993). Magma genesis in an ongoing rifting zone: The Tadjoura Gulf (Afar area). *Geochimica et Cosmochimica Acta*, 57, 2291–2302.
- Barth, G. A., Kastens, K. A., & Klein, E. M. (1994). The origin of bathymetric highs at ridge-transform intersections: Multi-disciplinary case study at the Clipperton fracture zone. *Marine Geophysical Research*, 16, 1–50.

- Batiza, R., Allan, J. F., Anderson, D. M., Bach, W., Bostroem, K., Brophy, J. G., et al. (1993). Site 864. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 142, pp. 55–99). College Station, Texas: Ocean Drilling Program.
- Batiza, R., Allan, J. F., Bach, W., Bostroem, K., Brophy, J. G., Fryer, G. J., et al. (1995). Petrology, geochemistry, and petrogenesis of Leg 142 basalts - synthesis of results. In R. Batiza, M. A. Storms, & J. F. Allan (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 142, pp. 3–8). College Station, Texas: Ocean Drilling Program.
- Batiza, R., & Johnson, J. R. (1980). Trace element and isotopic evidence for magma mixing in alkalic and transitional basalts near the East Pacific Rise at 8°N. *Initial Reports of the Deep Sea Drilling Project*, 54, 63–69.
- Batiza, R., Niu, Y., Karsten, J. L., Boger, W., Potts, E., Norby, L., & Butler, R. (1996). Steady and non-steady state magma chambers below the East Pacific Rise. *Geophysical Research Letters*, 23(3), 221–224. <https://doi.org/10.1029/95GL00016>
- Batiza, R., Oestrike, R., & Futa, K. (1982). Chemical and isotopic diversity in basalts dredged from the East Pacific Rise at 10°S, the Fossil Galapagos Rise and the Nazca Plate. *Marine Geology*, 49, 115–132.
- Batiza, R., Rosendahl, B. R., & Fisher, R. L. (1977). Evolution of oceanic crust. 3. Petrology and chemistry of basalts from the East Pacific Rise and the Siqueiros Transform Fault. *Journal of Geophysical Research*, 82(2), 265–276. <https://doi.org/10.1029/JB082i002p00265>
- Batiza, R., & Vanko, D. A. (1984). Petrology of young Pacific seamounts. *Journal of Geophysical Research*, 89, 11,235–11,260. <https://doi.org/10.1029/JB089iB13p11235>
- Beier, C., Haase, K. M., Abouchami, W., Krienitz, M. S., & Hauff, F. (2008). Magma genesis by rifting of oceanic lithosphere above anomalous mantle: Terceira Rift, Azores. *Geochemistry, Geophysics, Geosystems*, 9, Q12013. <https://doi.org/10.1029/2008GC002112>
- Ben Othman, D., & Allegre, C. J. (1990). U-Th isotopic systematics at 13°N East Pacific Ridge segment. *Earth and Planetary Science Letters*, 98, 129–137.
- Bender, J. F., Langmuir, C. H., & Hanson, G. N. (1984). Petrogenesis of basalt glasses from the Tamayo region, East Pacific Rise. *Journal of Petroleum*, 25, 213–254.
- Bezard, R., Fischer-Goedde, M., Hamelin, C., Brennecka, G., & Kleine, T. (2016). The effects of magmatic processes and crustal recycling on the molybdenum stable isotopic composition of Mid-Ocean Ridge basalts. *Earth and Planetary Science Letters*, 453, 171–181.
- Bezos, A., & Humler, E. (2005). The Fe<sup>3+</sup>/ΣFe ratios of MORB glasses and their implications for mantle melting. *Geochimica et Cosmochimica Acta*, 69, 711–725.
- Blichert-Toft, J., Agrani, A., Andres, M., Kingsley, R. H., Schilling, J.-G., & Albarede, F. (2005). Geochemical segmentation of the Mid-Atlantic Ridge north of Iceland and ridge-hot spot interaction in the North Atlantic. *Geochemistry, Geophysics, Geosystems*, 6, Q01E19. <https://doi.org/10.1029/2004GC000788>
- Bonatti, E., Colantoni, P., Della Vedova, B., & Taviani, M. (1984). Geology of the Red Sea transitional region (22°N to 25°N). *Oceanologica Acta*, 7, 385–398.
- Bonatti, E., Ferrara, G., & Honnorez, J. (1970). Equatorial Mid-Atlantic Ridge: Petrological and Sr-isotopic evidence for an alpine-type rock assemblage. *Earth and Planetary Science Letters*, 9, 24–256.
- Bonatti, E., Honnorez, J., & Ferrara, G. (1971). Peridotite-gabbro-basalt complex from the equatorial Mid-Atlantic Ridge. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 268, 385–402.
- Bonatti, E., Ottonello, G., & Hamlyn, P. R. (1986). Peridotites from the Island of Zabargad (St. John), Red Sea: Petrology and geochemistry. *Journal of Geophysical Research*, 91, 599–631. <https://doi.org/10.1029/JB091iB01p00599>
- Bostroem, K., & Bach, W. (1995). Data report: Chemical analyses of basaltic rocks: An interlaboratory comparison. In R. Batiza, M. A. Storms, & J. F. Allan (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 142, pp. 75–81). College Station, TX: Ocean Drilling Program.
- Bougault, H., Cambon, P., Corre, O., Joron, J.-L., & Treuil, M. (1979). Evidence for variability of magmatic processes and upper mantle heterogeneity in the Axial region of the Mid-Atlantic Ridge near 22° and 36°N. *Tectonophysics*, 55, 11–34.
- Bougault, H., Dmitriev, L. V., Schilling, J.-G., Sobolev, A. V., Joron, J.-L., & Needham, H. D. (1988). Mantle heterogeneity from trace elements: MAR triple junction near 14°N. *Earth and Planetary Science Letters*, 88, 27–36.
- Bougault, H., & Hekinian, R. (1974). Rift Valley in the Atlantic Ocean near 36°50'N: Petrology and geochemistry of basaltic rocks. *Earth and Planetary Science Letters*, 24, 249–261.
- Bougault, H., Joron, J.-L., & Treuil, M. (1984). Mantle heterogeneity: Along strike variations of hygromagmaphile elements between Azores and 10°N. *Proceedings of the 27th International Geological Congress*, 6, 213–244.
- Bougault, H., Joron, J.-L., Treuil, M., & Maury, R. C. (1985). Local versus regional mantle heterogeneities: Evidence from hygromagmaphile elements. *Initial Reports of the Deep Sea Drilling Project*, 82, 459–482.
- Bougault, H., & Treuil, M. (1980). Mid-Atlantic Ridge: Zero-age geochemical variations between Azores and 22°N. *Nature*, 286, 209–212.
- Bourdon, B., Goldstein, S. J., Bourles, D., Murrell, M. T., & Langmuir, C. H. (2000). Evidence from 10Be and U series disequilibria on the possible contamination of Mid-Ocean Ridge basalt glasses by sedimentary material. *Geochemistry, Geophysics, Geosystems*, 1(8), 1029. <https://doi.org/10.1029/2000GC000047>
- Bourdon, B., Langmuir, C. H., & Zindler, A. (1996). Ridge-hotspot interaction along the Mid-Atlantic Ridge between 37°30' and 40°30': The U-Th disequilibrium evidence. *Earth and Planetary Science Letters*, 142, 175–189.
- Brandl, P. A., Regelous, M., Beier, C., & Haase, K. M. (2013). High mantle temperatures following rifting caused by continental insulation. *Nature Geoscience*, 6(5), 391–394.
- Brevart, O., Dupre, B., & Allegre, C. J. (1981). Metallogensis at spreading centers: Lead isotope systematics for sulfides, manganese-rich crusts, basalts and sediments from the Cyamex and Alvin areas (East Pacific Rise). *Economic Geology*, 76, 1205–1210.
- Briais, A., Ondreas, H., Klingelhoefer, F., Dosso, L., Hamelin, C., & Guillou, H. (2009). Origin of volcanism on the flanks of the Pacific-Antarctic ridge between 41°30'S and 52°S. *Geochemistry, Geophysics, Geosystems*, 10, Q09001. <https://doi.org/10.1029/2008GC002350>
- Brooks, C. K., Jakobsson, S. P., & Campsie, J. (1974). Dredged basaltic rocks from the seaward extensions of the Rykjanes and Snaefellsnes volcanic zones, Iceland. *Earth and Planetary Science Letters*, 22, 320–327.
- Brueckner, H. K., Zindler, A., Seyler, M., & Bonatti, E. (1988). Zabargad and the isotopic evolution of the sub-Red Sea mantle and crust. *Tectonophysics*, 150, 163–176.
- Bryan, W. B. (1979). Regional variation and petrogenesis of basalt glasses from the FAMOUS area, Mid-Atlantic Ridge. *Journal of Petroleum*, 20, 293–325.
- Bryan, W. B., Frey, F. A., & Thompson, G. (1977). Oldest Atlantic seafloor. *Contributions to Mineralogy and Petrology*, 64, 223–242.
- Bryan, W. B., & Moore, J. G. (1977). Compositional variations of young basalts in the Mid-Atlantic Ridge rift valley near lat 36°49'N. *Geological Society of America Bulletin*, 88, 556–570.
- Bryan, W. B., Thompson, G., Frey, F. A., Dickey, J. S. Jr., & Roy, S. (1977). Petrology and geochemistry of basement rocks recovered on Leg 37, DSDP. *Initial Reports of the Deep Sea Drilling Project*, 37, 695–704.

- Bryan, W. B., Thompson, G., & Ludden, J. N. (1981). Compositional variation in normal MORB from 22°–25°N: Mid-Atlantic Ridge and Kane Fracture Zone. *Journal of Geophysical Research*, *86*, 11,815–11,836. <https://doi.org/10.1029/JB086iB12p11815>
- Bryan, W. B., Thompson, G., & Michael, P. J. (1979). Compositional variation in a steady-state zoned magma chamber: Mid-Atlantic Ridge at 36°50'N. *Tectonophysics*, *55*, 63–85.
- Burnard, P. (1999). The bubble-by-bubble volatile evolution of two Mid-Ocean Ridge basalts. *Earth and Planetary Science Letters*, *174*, 199–211.
- Burnard, P., Graham, D. W., & Farley, K. A. (2002). Mechanisms of magmatic gas loss along the Southeast Indian Ridge and the Amsterdam-St. Paul Plateau. *Earth and Planetary Science Letters*, *203*, 131–148.
- Burnard, P., Graham, D. W., & Farley, K. A. (2004). Fractionation of noble gases (He, Ar) during MORB mantle melting: A case study on the Southeast Indian Ridge. *Earth and Planetary Science Letters*, *227*, 457–472.
- Burnard, P., Harrison, D., Turner, G., & Nesbitt, R. W. (2003). Degassing and contamination of noble gases in Mid-Atlantic Ridge basalts. *Geochemistry, Geophysics, Geosystems*, *4*(1), 1002. <https://doi.org/10.1029/2002GC000326>
- Busigny, V., Laverne, C., & Bonifacie, M. (2005). Nitrogen content and isotopic composition of oceanic crust at a superfast spreading ridge: A profile in altered basalts from ODP site 1256, Leg 206. *Geochemistry, Geophysics, Geosystems*, *6*, Q12001. <https://doi.org/10.1029/2005GC001020>
- Busigny, V., Laverne, C., & Bonifacie, M. (2005). Nitrogen content and isotopic composition of oceanic crust at a superfast spreading ridge: A profile in altered basalts from ODP site 1256, Leg 206. *Geochemistry, Geophysics, Geosystems*, *6*, Q12001. <https://doi.org/10.1029/2005GC001020>
- Byerly, G. R. (1980). The nature of differentiation trends in some volcanic rocks from the Galapagos Spreading Center. *Journal of Geophysical Research*, *85*, 3797–3810. <https://doi.org/10.1029/JB085iB07p03797>
- Byerly, G. R., Melson, W. G., & Vogt, P. R. (1976). Rhyodacites, andesites, ferro-basalts and ocean tholeiites from the Galapagos Spreading Center. *Earth and Planetary Science Letters*, *30*, 215–221.
- Byerly, G. R., & Sinton, J. M. (1980). Compositional trends in natural basaltic glasses from Deep Sea Drilling Project Holes 417D and 418A1. *Initial Reports of the Deep Sea Drilling Project*, *51-53*, 957–971.
- Byers, C. D., Christie, D. M., Muenow, D. W., & Sinton, J. M. (1984). Volatile contents and ferric-ferrous ratios of basalt, ferrobasalt, andesite and rhyodacite glasses from the Galapagos 95.5°W propagating rift. *Geochimica et Cosmochimica Acta*, *48*, 2239–2245.
- Byers, C. D., Garcia, M. O., & Muenow, D. W. (1986). Volatiles in basaltic glasses from the East Pacific Rise at 21°N: Implications for MORB sources and submarine lava flow morphology. *Earth and Planetary Science Letters*, *79*, 9–20.
- Byers, C. D., Muenow, D. W., & Garcia, M. O. (1983). Volatiles in basalts and andesites from the Galapagos Spreading Center, 85° to 86°W. *Geochimica et Cosmochimica Acta*, *47*, 1551–1558.
- Campsie, J., Bailey, J. C., Rasmussen, M. H., & Dittmer, F. (1973). Chemistry of tholeiites from the Reykjanes Ridge and Charlie Gibbs fracture zone. *Nature*, *244*, 71–73.
- Carlson, R. W. (1980). Crust-mantle differentiation on the Earth and Moon: Evidence from isotopic studies for contrasting mechanisms and duration, (PhD thesis). San Diego, CA: California University.
- Carlson, R. W., Maccougall, J. D., & Lugmair, G. W. (1978). Differential Sm/Nd evolution in oceanic basalts. *Geophysical Research Letters*, *5*, 229–232. <https://doi.org/10.1029/GL005i004p00229>
- Cartigny, P., Jendrzewski, N., Pineau, F., Petit, E., & Javoy, M. (2001). Volatile (C, N, Ar) variability in MORB and the respective roles of mantle source heterogeneity and degassing; the case of the Southwest Indian Ridge. *Earth and Planetary Science Letters*, *194*, 241–257.
- Casey, J. F. (1997). Comparison of major- and trace-element geochemistry of abyssal peridotites and mafic plutonic rocks with basalts from the mark region of the Mid-Atlantic Ridge. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 153, pp. 181–241). College Station, Texas: Ocean Drilling Program.
- Castillo, P. R., & Batiza, R. (1989). Strontium, neodymium and lead isotope constraints on near-ridge seamount production beneath the South Atlantic. *Nature*, *342*, 262–265.
- Castillo, P. R., Klein, E. M., Bender, J. F., Langmuir, C. H., Shirey, S. B., Batiza, R., & White, W. M. (2000). Petrology and Sr, Nd, and Pb isotope geochemistry of Mid-Ocean Ridge basalt glasses from the 11°45' to 15° 00'N segment of the East Pacific Rise. *Geochemistry, Geophysics, Geosystems*, *1*(11), 1011. <https://doi.org/10.1029/1999GC000024>
- Castillo, P. R., Natland, J. H., Niu, Y., & Lonsdale, P. F. (1998). Sr, Nd and Pb isotopic variation along the Pacific-Antarctic risecrest, 53–57°S: Implications for the composition and dynamics of the South Pacific upper mantle. *Earth and Planetary Science Letters*, *154*, 109–125.
- Chadwick, J., Perfit, M. R., Ridley, W. I., Jonasson, I. R., Kamenov, G. D., Chadwick, W., et al. (2005). Magmatic effects of the Cobb hot spot on the Juan de Fuca Ridge. *Journal of Geophysical Research*, *110*, B03101. <https://doi.org/10.1029/2003JB002767>
- Chaussidon, M., & Jambon, A. (1994). Boron content and isotopic composition of oceanic basalts: Geochemical and cosmochemical implications. *Earth and Planetary Science Letters*, *121*, 277–291.
- Chaussidon, M., Sheppard, S. M. F., & Michard, A. (1991). Hydrogen, sulphur and neodymium isotope variations in the mantle beneath the EPR at 12°50'N. *Journal Geochemical Society*, *3*, 325–337.
- Chauvel, C., & Blichert-Toft, J. (2001). A hafnium isotope and trace element perspective on melting of the depleted mantle. *Earth and Planetary Science Letters*, *190*, 137–151.
- Chavrit, D., Burgess, R., Sumino, H., Teagle, D. A. H., Droop, G., Shimizu, A., & Ballentine, C. (2016). The contribution of hydrothermally altered ocean crust to the mantle halogen and noble gas cycles. *Geochimica et Cosmochimica Acta*, *183*, 106–124.
- Chen, Z.-Q., Zhou, H.-Y., Liu, Y., Yang, Q.-H., Li, J.-W., & Dick, H. J. B. (2012). Influence of igneous processes and serpentinization on geochemistry of the Logatchev Massif harzburgites (14°45'N, Mid-Atlantic Ridge) and comparison with global abyssal peridotites. *International Geology Review*, *55*, 115–130.
- Christie, D. M. (1984). Petrological effects of mid-oceanic rift propagation: The Galapagos spreading center at 95.5°W, Ph.D. Desertation, Honolulu, Hawaii: University of Hawaii.
- Christie, D. M., Carmichael, I., & Langmuir, C. H. (1986). Oxidation states of mid-ocean-ridge basalt glasses. *Earth and Planetary Science Letters*, *79*, 397–411.
- Christie, D. M., Pyle, D. G., & Sylvander, B. A. (2004). Major and trace element composition of basalts from the Southeast Indian Ridge (westward cruise, Leg 10). Earthchem Library.
- Christie, D. M., Werner, R., Hauff, F., & Hanan, B. B. (2005). Morphological and geochemical variations along the eastern Galapagos Spreading Center. *Geochemistry, Geophysics, Geosystems*, *6*, Q01006. <https://doi.org/10.1029/2004GC000714>
- Church, S. E., & Tatsumoto, M. (1975). Lead isotope relocations in oceanic ridge basalts from the Juan de Fuca-Gorda Ridge area, NE Pacific Ocean. *Contributions to Mineralogy and Petrology*, *53*, 253–279.

- Cipriani, A., Brueckner, H. K., Bonatti, E., & Brunelli, D. (2004). Oceanic Crust Generated By Elusive Parents: Sr And Nd Isotopes In Basalt-Peridotite Pairs From The Mid-Atlantic Ridge. *Geology*, 32(8), 657–660.
- Clague, D. A., Dreyer, B., Paduan, J. B., Chadwick, W., Caress, D., Portner, R., et al. (2013). Geologic history of the summit of Axial Seamount, Juan de Fuca Ridge. *Geochemistry, Geophysics, Geosystems*, 14, 4403–4443. <https://doi.org/10.1002/ggge.20240>
- Clague, D. A., Frey, F. A., Thompson, G., & Rindge, S. (1981). Minor and trace element geochemistry of volcanic rocks dredged from the Galapagos Spreading Center: Role of crystal fractionation and mantle heterogeneity. *Journal of Geophysical Research*, 86, 9469–9482. <https://doi.org/10.1029/JB086iB10p09469>
- Clague, D. A., Friesen, W. B., Quintero, P., Holmes, M., Morton, J. L., Bouse, R., et al. (1984). Preliminary geological, geophysical, and biological data from the Gorda Ridge. U.S. Geological Survey Open-File Report 84, 364–49.
- Clague, D. A., & Holmes, M. (1989). Geology, petrology and mineral potential of the Gorda Ridge. In D. W. Scholl, A. Grantz, & J. G. Vedder (Eds.), *Geology and Resource Potential of the Continental Margin of Western North America and Adjacent Ocean Basins - Beaufort Sea to Baja California*, Earth Science Series (Vol. 6, pp. 563–580). Houston, TX: Circum Pacific Council Energy Mineral Resources.
- Clague, D. A., Paduan, J. B., & Davis, A. S. (2009). Widespread strombolian eruptions of mid-ocean ridge basalts. *Journal of Volcanology and Geothermal Research*, 180, 171–188.
- Clague, D. A., Paduan, J. B., Duncan, R. A., Huard, J. J., Davis, A. S., Castillo, P. R., et al. (2009). Five million years of compositionally diverse, episodic volcanism: Construction of Davidson Seamount atop an abandoned spreading center. *Geochemistry, Geophysics, Geosystems*, 10, Q12009. <https://doi.org/10.1029/2009GC002665>
- Clarke, D. B., Cameron, B., Muecke, G. K., & Bates, J. L. (1989). Early tertiary basalts from the Labrador Sea floor and Davis Strait region. *Canadian Journal of Earth Sciences*, 26, 956–968.
- Clog, M., Aubad, C., Cartigny, P., & Dosso, L. (2013). The hydrogen isotopic composition and water content of southern Pacific MORB: A reassessment of the D/H ratio of the depleted mantle reservoir. *Earth and Planetary Science Letters*, 381, 156–165.
- Cohen, R. S., Evensen, N. M., Hamilton, P. J., & O'Nions, R. K. (1980). U-Pb, Sm-Nd and Rb-Sr systematics of Mid-Ocean Ridge basalt glasses. *Nature*, 283, 149–153.
- Cohen, R. S., & O'Nions, R. K. (1982). The lead, neodymium and strontium isotopic structure of ocean ridge basalts. *Journal of Petroleum*, 23, 299–324.
- Colin, A., Burnard, P., Graham, D. W., & Marrocchi, Y. (2011). Plume-ridge interaction along the Galapagos Spreading Center: Discerning between gas loss and source effects using neon isotopic compositions and  $4\text{He}-40\text{Ar}^*$ - $\text{CO}_2$  relative abundances. *Geochimica et Cosmochimica Acta*, 75, 1145–1160.
- Colin, A., Moreira, M., Gautheron, C., & Burnard, P. (2015). Constraints on the noble gas composition of the deep mantle by bubble-by-bubble analysis of a volcanic glass sample from Iceland. *Chemical Geology*, 417, 173–183. <https://doi.org/10.1016/j.chemgeo.2015.09.020>
- Condomines, M., Morand, P., & Allegre, C. J. (1981).  $^{230}\text{Th}$ - $^{238}\text{U}$  radioactive disequilibria in tholeiites from the FAMOUS zone (Mid-Atlantic Ridge,  $36^{\circ}50'\text{N}$ ): th and sr isotopic geochemistry. *Earth and Planetary Science Letters*, 55, 247–256.
- Coogan, L. A., Thompson, G. M., MacLeod, C. J., Dick, H. J. B., Edwards, S. J., Hosford Scheirer, A., & Barry, T. L. (2004). A combined basalt and peridotite perspective on 14 million years of melt generation at the Atlantis Bank segment of the Southwest Indian Ridge: Evidence for temporal changes in mantle dynamics? *Chemical Geology*, 207(1), 13–30.
- Cooper, K. M. (2007). Data report: Trace element analyses in whole-rock basement samples, site 1256, ODP Leg 206. In D. A. H. Teagle, D. S. Wilson, G. A., Acton, & D. A. Vanko (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 206, pp. 1–11). College Station, Texas: Ocean Drilling Program.
- Cooper, K. M., Goldstein, S. J., Sims, K. W. W., & Murrell, M. T. (2003). Uranium-series chronology of Gorda Ridge volcanism: New evidence from the 1996 eruption. *Earth and Planetary Science Letters*, 206, 459–475.
- Cooper, L. B., Plank, T., Arculus, R. J., Hauri, E. H., & Hall, P. S. (2010). High-Ca boninites from the active Tonga Arc. *Journal of Geophysical Research*, 115, B10206. <https://doi.org/10.1029/2009JB006367>
- Cordier, C., Caroff, M., Juteau, T., Fleutelot, C., Hémond, C., Drouin, M., et al. (2007). Bulk-rock geochemistry and plagioclase zoning in lavas exposed along the northern flank of the Western Blanco Depression (Northeast Pacific): Insight into open-system magma chamber processes. *Lithos*, 99(3-4), 289–311. <https://doi.org/10.1016/j.lithos.2007.06.009>
- Cottrell, E., & Kelley, K. A. (2011). The oxidation state of Fe in MORB glasses and the oxygen fugacity of the upper mantle. *Earth and Planetary Science Letters*, 305, 270–282.
- Cottrell, E., & Kelley, K. A. (2011). The oxidation state of Fe in MORB glasses and the oxygen fugacity of the upper mantle. *Earth and Planetary Science Letters*, 305, 270–282. <https://doi.org/10.1016/j.epsl.2011.03.014>
- Cottrell, E., & Kelley, K. A. (2013). Redox heterogeneity in Mid-Ocean Ridge basalts as a function of mantle source. *Science*, 340, 1314–1317.
- Cousens, B. L., Allan, J. F., Leybourne, M. I., Chase, R. L., & van Wagoner, N. A. (1995). Mixing of magmas from enriched and depleted mantle sources in the northeast Pacific: West Valley segment, Juan de Fuca Ridge. *Contributions to Mineralogy and Petrology*, 120, 337–357.
- Cousens, B. L., Chase, R. L., & Schilling, J.-G. (1984). Basalt geochemistry of the Explorer Ridge area, northeast Pacific Ocean. *Canadian Journal of Earth Sciences*, 31, 157–170.
- Craig, H., & Lupton, J. E. (1976). Primordial neon, helium, and hydrogen in oceanic basalts. *Earth and Planetary Science Letters*, 31, 369–385.
- Cushman, B., Sinton, J. M., Ito, G., & Dixon, J. E. (2004). Glass compositions, plume-ridge interaction, and hydrous melting along the Galapagos Spreading Center,  $990.5^{\circ}\text{W}$  to  $98^{\circ}\text{W}$ . *Geochemistry, Geophysics, Geosystems*, 5, Q08E17. <https://doi.org/10.1029/2004GC000709>
- Czamanske, G. K., & Moore, J. G. (1977). Composition and phase chemistry of sulfide globules from Mid-Atlantic Ridge rift valley near  $37^{\circ}\text{N}$  latitude. *Geological Society of America Bulletin*, 88, 587–599.
- Dale, C. W., Macpherson, C. G., Pearson, G., Hammond, S., & Arculus, R. J. (2012). Inter-element fractionation of highly siderophile elements in the Tonga Arc due to flux melting of a depleted source. *Geochimica et Cosmochimica Acta*, 89, 202–225.
- Dalrymple, G., Clague, D. A., Vallier, T. L., & Menard, H. (1987). Ar40/Ar39 age, petrology and tectonic significance of some seamounts in the Gulf of Alaska. B. H. Keating, P. Fryer, R. Batiza, & G. W. Boehlert, (Eds.), *AGU Geophysical Monograph* (Vol. 43). Seamounts, Islands, and Atolls.
- Danyushevsky, L. V., Eggins, S. M., Falloon, T. J., & Christie, D. M. (2000). H<sub>2</sub>O abundance in depleted to moderately enriched Mid-Ocean Ridge magmas; part I: Incompatible behaviour, implications for mantle storage, and origin of regional variations. *Journal of Petroleum*, 41, 1329–1364.
- Danyushevsky, L. V., Falloon, T. J., Sobolev, A. V., Crawford, A. J., Carroll, M., & Price, R. C. (1993). The H<sub>2</sub>O content of basalt glasses from Southwest Pacific back-arc basins. *Earth and Planetary Science Letters*, 117, 347–362.
- Danyushevsky, L. V., Perfit, M. R., Eggins, S. M., & Falloon, T. J. (2003). Crustal origin for coupled ultra-depleted and plagioclase signatures in MORB olivine-hosted melt inclusions: Evidence from the Siqueiros Transform Fault, East Pacific Rise. *Contributions to Mineralogy and Petrology*, 144, 619–637.

- David, K., Schiano, P., & Allegre, C. J. (2000). Assessment of the Zr/Hf fractionation in oceanic basalts and continental materials during petrogenetic processes. *Earth and Planetary Science Letters*, *178*, 285–301.
- Davis, A. S., & Clague, D. A. (1987). Geochemistry, mineralogy, and petrogenesis of basalt from the Gorda Ridge. *Journal of Geophysical Research*, *92*, 10,476–10,483.
- Davis, A. S., Clague, D. A., Cousens, B. L., Keaten, R., & Paduan, J. B. (2008). Geochemistry of basalt from the north Gorda segment of the Gorda Ridge: Evolution toward ultraslow spreading ridge lavas due to decreasing magma supply. *Geochemistry, Geophysics, Geosystems*, *9*, Q04004. <https://doi.org/10.1029/2007GC001775>
- Davis, A. S., Clague, D. A., & Friesen, W. B. (1993). Petrology and mineral chemistry of basalt from Escanaba Trough. In J. L. Morton, R. A. Zierenberg, & C. A. Reiss (Eds.), *Geologic, Hydrothermal, and Biologic Studies At Escanaba Trough, U.S. Geological Survey* (Vol. 2022, pp. 153–170). Offshore Northern California: Gorda Ridge.
- Davis, A. S., Clague, D. A., & Morton, J. L. (1990). Volcanic glass compositions from two spreading centers in Lau Basin, southwest Pacific Ocean. *Geologisches Jahrbuch*, *92*, 481–501.
- Davis, A. S., Clague, D. A., & White, W. M. (1998). Geochemistry of basalt from Escanaba Trough: Evidence for sediment contamination. *Journal of Petroleum*, *39*, 841–858.
- De Paolo, D. J., & Wasserburg, G. J. (1976). Inferences about magma sources and mantle structure from variations of  $^{143}\text{Nd}/^{144}\text{Nd}$ . *Geophysical Research Letters*, *3*, 743–746. <https://doi.org/10.1029/GL003i012p00743>
- Debaillé, V., Blichert-Toft, J., Agraniér, A., Doucelance, R., Schiano, P., & Albarede, F. (2006). Geochemical component relationships in MORB from the Mid-Atlantic Ridge, 22–35°N. *Earth and Planetary Science Letters*, *241*, 844–862.
- Delaney, J. R., Muenow, D. W., & Graham, D. G. (1978). Abundance and distribution of water, carbon and sulfur in the glassy rims of submarine pillow basalts. *Geochimica et Cosmochimica Acta*, *42*, 581–594.
- Deruelle, B., Dreibus, G., & Jambon, A. (1992). Iodine abundances in oceanic basalts: Implications for Earth dynamics. *Earth and Planetary Science Letters*, *108*, 217–227.
- Des Marais, D. J., & Moore, J. G. (1984). Carbon and its isotopes in mid-ocean basaltic glasses. *Earth and Planetary Science Letters*, *69*, 43–57.
- Detrick, R. S., Sinton, J. M., Ito, G., Canales, J. P., Behn, M., Blacic, T., et al. (2002). Correlated geophysical, geochemical, and volcanological manifestations of plume-ridge interaction along the Galapagos Spreading Center. *Geochemistry, Geophysics, Geosystems*, *3*(10), 8501. <https://doi.org/10.1029/2002GC000350>
- Devey, C. W., Garbe-Schönberg, D., Stoffers, P., Chauvel, C., & Mertz, D. F. (1994). Geochemical effects of dynamic melting beneath ridges: Reconciling major and trace element variations in Kolbeinsey (and global) Mid-Ocean Ridge basalt. *Journal of Geophysical Research*, *99*, 9077–9095. <https://doi.org/10.1029/93JB03364>
- Di Donato, G., Joron, J.-L., Treuil, M., & Loubet, M. (1990). Geochemistry of zero-age n-MORB from hole 648B, ODP Legs 106–109, M.A.R., 22°N. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 106–109, pp. 57–65). College Station, Texas: Ocean Drilling Program.
- Dickey, J. S. Jr., Frey, F. A., Hart, S. R., Watson, E. B., & Thompson, G. (1977). Geochemistry and petrology of dredged basalts from the Bouvet triple junction, South Atlantic. *Geochimica et Cosmochimica Acta*, *41*, 1105–1118.
- Dittmer, F., Fine, S., Rasmussen, M. H., Bailey, J. C., & Campsie, J. (1975). Dredged basalts from the Mid-Atlantic Ridge north of Iceland. *Nature*, *254*, 298–301.
- Dixon, J. E., Clague, D. A., & Eissen, J. P. (1986). Gabbroic xenoliths and host ferrobasalt from the southern Juan de Fuca Ridge. *Journal of Geophysical Research*, *91*(B3), 3795–3820. <https://doi.org/10.1029/JB091iB03p03795>
- Dixon, J. E., Stolper, E., & Delaney, J. R. (1988). Infrared spectroscopic measurements of CO<sub>2</sub> and H<sub>2</sub>O in Juan de Fuca Ridge basaltic glasses. *Earth and Planetary Science Letters*, *90*, 87–104.
- Dmitriev, L. V., Sobolev, A. V., Sushchevskaya, N. M., & Zapunny, S. A. (1985). Abyssal glasses, petrologic mapping of the oceanic floor and "geochemical Leg" 82. *Initial Reports of the Deep Sea Drilling Project*, *82*, 508–518.
- Donnelly, K. E., Goldstein, S. L., Langmuir, C. H., & Spiegelman, M. (2004). Origin of enriched ocean ridge basalts and implications for mantle dynamics. *Earth and Planetary Science Letters*, *226*, 347–366.
- Dosso, L., Bougault, H., Beuzart, P., Calvez, J. Y., & Joron, J.-L. (1988). The geochemical structure of the south-east Indian Ridge. *Earth and Planetary Science Letters*, *88*, 47–59.
- Dosso, L., Bougault, H., & Joron, J.-L. (1993). Geochemical morphology of the north Mid-Atlantic Ridge, 10°–24°N: Trace element - isotope complementarity. *Earth and Planetary Science Letters*, *120*, 443–462.
- Dosso, L., Bougault, H., Langmuir, C. H., Bollinger, C., Bonnier, O., & Etoubleau, J. (1999). The age and distribution of mantle heterogeneity along the Mid-Atlantic Ridge (31–41°N). *Earth and Planetary Science Letters*, *170*, 269–286.
- Dosso, L., Hanan, B. B., Bougault, H., Schilling, J.-G., & Joron, J.-L. (1991). Sr-Nd-Pb geochemical morphology between 10° and 17°N on the Mid-Atlantic Ridge: A new MORB isotope signature. *Earth and Planetary Science Letters*, *106*, 29–43.
- Douglass, J., Schilling, J.-G., & Fontignie, D. (1999). Plume-ridge interactions of the discovery and shona mantle plumes with the southern Mid-Atlantic Ridge (40–55°S). *Journal of Geophysical Research*, *104*(B2), 2941–2962. <https://doi.org/10.1029/98JB02642>
- Dreyer, B., Clague, D. A., & Gill, J. B. (2013). Petrological variability of recent magmatism at Axial Seamount, Juan de Fuca Ridge. *Geochemistry, Geophysics, Geosystems*, *14*, 4306–4333. <https://doi.org/10.1002/ggge.20239>
- Dril, S. I., Kuzmin, M. I., Tsipukova, S. S., & Zonenshain, L. P. (1997). Geochemistry of basalts from the western Woodlark, Lau and Manus basins: Implications for the petrogenesis and source rock compositions. *Marine Geology*, *142*, 57–83.
- Duncan, R. A., & Holmes, M. (1994). Radiometric dating of young MORB using the  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  incremental heating method. *Geophysical Research Letters*, *21*(18), 1927–1930. <https://doi.org/10.1029/94GL01375>
- Dungan, M. A., & Rhodes, J. M. (1978). Residual glasses and melt inclusions in basalts from DSDP Legs 45 and 46: Evidence for magma mixing. *Contributions to Mineralogy and Petrology*, *67*, 417–431.
- Dungan, M. A., Rhodes, J. M., Long, P. E., Blanchard, D. P., Brannon, J. C., & Rodgers, K. V. (1978). The petrology and geochemistry of basalts from Site 396 Legs 45 and 46 of the Deep Sea Drilling Project. *Initial Reports of the Deep Sea Drilling Project*, *46*, 89–113.
- Dupre, B., & Allegre, C. J. (1983). Pb-Sr isotope variation in Indian Ocean basalts and mixing phenomena. *Nature*, *303*, 142–146.
- Dupre, B., Lambret, B., Rousseau, D., & Allegre, C. J. (1981). Limitations on the scale of mantle heterogeneities under oceanic ridges lead and strontium isotope variations in the FAMOUS and CYAMEX areas. *Nature*, *294*, 552–554.
- Dymond, J., & Hogan, L. G. (1973). Noble gas abundance pattern in deep-sea basalts - Primordial gases from the mantle. *Earth and Planetary Science Letters*, *20*, 131.
- Eaby, J. S., Clague, D. A., & Delaney, J. R. (1984). Sr isotopic variations along the Juan de Fuca Ridge. *Journal of Geophysical Research*, *89*(B9), 7883–7890. <https://doi.org/10.1029/JB089iB09p07883>
- Eason, D., & Sinton, J. M. (2006). Origin of high-Al N-MORB by fractional crystallization in the upper mantle beneath the Galapagos Spreading Center. *Earth and Planetary Science Letters*, *252*, 423–436.

- Eiler, J. M., Schiano, P., Kitchen, N., & Stolper, E. (2000). Oxygen-isotope evidence for recycled crust in the sources of Mid-Ocean Ridge basalts. *Nature*, *403*, 530–534.
- Eissen, J. P. (1982). Petrologie comparee de basaltes de differents segments de zones d'accrétion oceanique à taux d'accrétion varies (Mer Rouge, Atlantique, Pacifique). Thesis Or Dissertation, Doctoral, Univ Strasbourg; Monographic.
- Eissen, J. P., Juteau, T., Joron, J.-L., Dupre, B., Humler, E., & Al'Mukhamedov, A. (1989). Petrology and geochemistry of basalts from the Red Sea Axial Rift at 18° North. *Journal of Petroleum*, *30*, 791–839.
- Eissen, J. P., Lefevre, C., Maillat, P., Morvan, G., & Nohara, M. (1991). Petrology and geochemistry of the central north Fiji Basin Spreading Center (southwest Pacific) between 16°S and 22°S. *Marine Geology*, *98*, 201–239.
- Elkins, L. J., Sims, K. W. W., Prytulak, J., Blichert-Toft, J., Elliott, T., Blusztajn, J., et al. (2014). Melt generation beneath Arctic Ridges: Implications from U decay series disequilibria in the Mohns, Lnipovich, and Gakkel Ridges. *Geochimica et Cosmochimica Acta*, *127*, 140–170.
- Elkins, L. J., Sims, K. W. W., Prytulak, J., Elliott, T., Mattielli, N., Blichert-Toft, J., et al. (2011). Understanding melt generation beneath the slow-spreading Kolbeinsey Ridge using <sup>238</sup>U, <sup>230</sup>Th, and <sup>231</sup>Pa excesses. *Geochimica et Cosmochimica Acta*, *75*, 6300–6329.
- Elthon, D., Ross, D. K., & Meen, J. K. (1995). Compositional variations of basaltic glasses from the Mid-Cayman Rise spreading center. *Journal of Geophysical Research*, *100*, 12,497–12,512. <https://doi.org/10.1029/94JB02777>
- Embley, R. W., Jonasson, I. R., Perfit, M. R., Franklin, J. M., Tivey, M. A., Malahoff, A., et al. (1988). Submersible investigation of an extinct hydrothermal system on the Galapagos Ridge: Sulfide mound, stockwork zone, and differentiated lavas. *The Canadian Mineralogist*, *26*, 517–539.
- Engel, A. E. J., & Engel, C. G. (1964a). Composition of basalts from the Mid-Atlantic Ridge. *Science*, *144*, 1330–1333.
- Engel, A. E. J., & Engel, C. G. (1964b). Igneous rocks of the East Pacific Rise. *Science*, *146*, 477–485.
- Engel, A. E. J., Engel, C. G., & Havens, R. G. (1965). Chemical characteristics of oceanic basalts and the upper mantle. *Geological Society of America Bulletin*, *76*, 719–734.
- Engel, C. G., & Fisher, R. L. (1969). Lherzolite, anorthosite, gabbro, and basalt dredged from the Mid-Indian Ocean Ridge. *Science*, *166*, 1136–1141.
- Engel, C. G., & Fisher, R. L. (1975). Granitic to ultramafic rock complexes of the Indian Ocean Ridge system, western Indian Ocean. *Geological Society of America Bulletin*, *96*, 1553–1578.
- Engel, C. G., Fisher, R. L., & Engel, A. E. J. (1965). Igneous rocks of the Indian Ocean floor. *Science*, *150*, 605–610.
- Erlank, A. J., & Kable, E. J. D. (1976). The significance of incompatible elements in Mid-Atlantic Ridge basalts from 45°N with particular reference to Zr/Nb. *Contributions to Mineralogy and Petrology*, *54*, 281–291.
- Escrig, S., Capmas, F., Dupre, B., & Al'Legre, C. J. (2004). Osmium isotopic constraints on the nature of the dupal anomaly from Indian Mid-Ocean-Ridge basalts. *Nature*, *431*, 59–63.
- Falloon, T. J., Malahoff, A., Zonenshain, L. P., & Bogdanov, Y. (1992). Petrology and geochemistry of back-arc basin basalts from Lau Basin spreading ridges at 15°, 18° and 19°S. *Mineralogy and Petrology*, *47*, 1–35.
- Ferguson, E. M., Klein, E. M., & Gilod, D. A. (1993). Fresh basalts from the Pacific-Antarctic Ridge extend the Pacific geochemical province. *Nature*, *366*, 330–333.
- Fine, G., & Stolper, E. (1985). Dissolved carbon dioxide in basaltic glasses: Concentrations and speciation. *Earth and Planetary Science Letters*, *76*, 263–278.
- Fisk, M. R., Bence, A. E., & Schilling, J.-G. (1982). Major element chemistry of Galapagos Rift Zone magmas and their phenocrysts. *Earth and Planetary Science Letters*, *61*, 171–189.
- Font, L., Murton, B. J., Roberts, S., & Tindle, A. G. (2007a). Variations in melt productivity and melting conditions along SWIR (70°E - 49°E): Evidence from olivine-hosted and plagioclase-hosted melt inclusions. *Journal of Petroleum*, *48*, 1471–1494.
- Font, L., Murton, B. J., Roberts, S., & Tindle, A. G. (2007b). Variations in melt productivity and melting conditions along swir (70 degrees e - 49 degrees east): Evidence from olivine-hosted and plagioclase-hosted melt inclusions. *Journal of Petroleum*, *48*, 1471–1494.
- Fontignie, D., & Schilling, J.-G. (1991). <sup>87</sup>Sr/<sup>86</sup>Sr and REE variations along the Easter Microplate boundaries (south Pacific): Application of multivariate statistical analyses to ridge segmentation. *Chemical Geology*, *89*, 209–241.
- Fontignie, D., & Schilling, J.-G. (1996). Mantle heterogeneities beneath the south Atlantic: a Nd-Sr-Pb isotope study along the Mid-Atlantic Ridge (3°S- 46°S). *Earth and Planetary Science Letters*, *142*, 209–221.
- Fornari, D. J., Perfit, M. R., Allan, J. F., Batiza, R., Haymon, R., Barone, A., et al. (1988). Geochemical and structural studies of the Lamont Seamounts: Seamounts as indicators of mantle processes. *Earth and Planetary Science Letters*, *89*, 63–83.
- Fornari, D. J., Perfit, M. R., Malahoff, A., & Embley, R. W. (1983). Geochemical studies of abyssal lavas recovered by DSRV Alvin from eastern Galapagos Rift, Inca Transform, and Ecuador Rift 1. Major element variations in natural glasses and spatial distribution of lavas. *Journal of Geophysical Research*, *88*(B12), 10,519–10,529. <https://doi.org/10.1029/JB088B12p10519>
- Fretzdorff, S., Haase, K. M., Leat, P. T., Livermore, R. A., Garbe-Schönberg, D., Fietzke, J., & Stoffers, P. (2003). <sup>230</sup>Th-<sup>238</sup>U disequilibrium in East Scotia backarc basalts: Implications for slab contributions. *Geology*, *31*, 693–696.
- Fretzdorff, S., Livermore, R. A., Devey, C. W., Leat, P. T., & Stoffers, P. (2002). Petrogenesis of the back-arc East Scotia Ridge, south Atlantic Ocean. *Journal of Petroleum*, *43*, 1435–1467.
- Freund, S., Beier, C., Krumm, S., & Haase, K. M. (2013). Oxygen isotope evidence for the formation of andesitic-dacitic magmas from the fast-spreading Pacific-Antarctic Rise by assimilation-fractional crystallisation. *Chemical Geology*, *347*, 271–283.
- Frey, F. A., Bryan, W. B., & Thompson, G. (1974). Atlantic Ocean floor: Geochemistry and petrology of basalts from Legs 2 and 3 of the Deep-Sea Drilling Project. *Journal of Geophysical Research*, *79*(35), 5507–5527. <https://doi.org/10.1029/JB079i035p05507>
- Frey, F. A., Dickey, J. S. Jr., Thompson, G., & Bryan, W. B. (Eds.). (1976). Eastern Indian ocean dsdp sites: correlations between petrography, geochemistry, and tectonic setting. Geological Society of America, Deep-Sea Drilling In The Indian Ocean.
- Frey, F. A., Dickey, J. S. Jr., Thompson, G., Bryan, W. B., & Davies, H. L. (1980). Evidence for heterogeneous primary MORB and mantle sources, NW Indian Ocean. *Contributions to Mineralogy and Petrology*, *74*, 387–402.
- Frey, F. A., Haskin, M. A., Poetz, J. A., & Haskin, L. A. (1968). Rare earth abundances in some basic rocks. *Journal of Geophysical Research*, *73*(18), 6085–6098. <https://doi.org/10.1029/JB073i018p06085>
- Frey, F. A., Walker, N., Stakes, D., Hart, S. R., & Nielsen, R. L. (1993). Geochemical characteristics of basaltic glasses from the AMAR and FAMOUS axial valleys, Mid-Atlantic Ridge (36° - 37°N): Petrogenetic implications. *Earth and Planetary Science Letters*, *115*, 117–136.
- Frisby, C., Bizimis, M., & Mallick, S. (2016a). Seawater-derived rare earth element addition to abyssal peridotites during serpentinization. *Lithos*, *248-251*, 432–454.
- Frisby, C., Bizimis, M., & Mallick, S. (2016b). Hf-Nd isotope decoupling in bulk abyssal peridotites due to serpentinization. *Chemical Geology*, *440*, 60–72. <https://doi.org/10.1016/j.chemgeo.2016.07.006>

- Fujii, T., Yamashita, T. I., Ishii, T., Tainosho, Y., Langmuir, C. H., Nakada, S., et al. (1995). Petrological sampling at the Rodriguez Triple Junction in the Indian Ocean. Kh93-3 Cruise Report, Ocean Research Institute, University Of Tokyo.
- Funkhouser, J. G., Fisher, D. E., & Bonatti, E. (1968). Excess argon in deep-sea rocks. *Earth and Planetary Science Letters*, *5*, 95–100.
- Furi, E., Hilton, D. R., Murton, B. J., Hemond, C., Dymont, J., & Day, J. M. D. (2011). Helium isotope variations between Réunion Island and the Central Indian Ridge (17°–21°S): New evidence for ridge–hot spot interaction. *Journal of Geophysical Research*, *116*, B02207. <https://doi.org/10.1029/2010JB007609>
- Gale, A., Escrig, S., Gier, E. J., Langmuir, C. H., & Goldstein, S. L. (2011). Enriched basalts at segment centers: The Lucky Strike (37°17'N) and Menez Gwen (37°50'N) segments of the Mid-Atlantic Ridge. *Geochemistry, Geophysics, Geosystems*, *12*, Q06016. <https://doi.org/10.1029/2010GC003446>
- Gannoun, A., Burton, K. W., Parkinson, I. J., Alard, O., Schiano, P., & Thomas, L. (2007). The scale and origin of the osmium isotope variations in Mid-Ocean Ridge basalts. *Earth and Planetary Science Letters*, *259*, 541–556.
- Gautheron, C., Moreira, M., Gerin, C., Tassan-Got, L., Bezos, A., & Humler, E. (2015). Constraints on the DUPAL anomaly from helium isotope systematics in the southwest Indian Mid-Ocean Ridge basalts. *Chemical Geology*, *417*, 163–172.
- Geist, D. J., Diefenbach, B., Fornari, D. J., Kurz, M. D., Harpp, K. S., & Blusztajn, J. (2008). Construction of the Galápagos platform by large submarine volcanic terraces. *Geochemistry, Geophysics, Geosystems*, *9*, Q03015. <https://doi.org/10.1029/2007GC001795>
- Geist, D. J., Naumann, T., Standish, J.-J., Kurz, M. D., Harpp, K. S., White, W. M., & Fornari, D. J. (2005). Wolf Volcano, Galapagos Archipelago: Melting and magmatic evolution at the margins of a mantle plume. *Journal of Petroleum*, *46*, 2197–2224. <https://doi.org/10.1093/petrology/egi052>
- Geldmacher, J., Hoefig, T., Hauff, F., Hoernle, K., Garbe-Schönberg, D., & Wilson, D. S. (2013). Influence of the Galapagos hotspot on the East Pacific Rise during Miocene superfast spreading. *Geology*, *41*, 183–186.
- Georgen, J., Kurz, M. D., Dick, H. J. B., & Lin, J. (2003). Low 3He/4He ratios in basalt glasses from the western Southwest Indian Ridge (10°–24°E). *Earth and Planetary Science Letters*, *206*, 509–528.
- Geshi, N., Umino, S., Kumagai, H., Sinton, J. M., White, S. M., Kisimoto, K., & Hilde, T. W. C. (2007). Discrete plumbing systems and heterogeneous magma sources of a 24 km<sup>2</sup> off-axis lava field on the western flank of East Pacific Rise, 14°S. *Earth and Planetary Science Letters*, *258*, 61–72.
- Goldstein, S. J. (1995). Uranium-series chronology of subsurface basalts, 9°31'N East Pacific Rise. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 142, pp. 37–39). College Station, Texas: Ocean Drilling Program.
- Goldstein, S. J., Murrell, M. T., & Janecky, D. R. (1989). Th and U isotopic systematics of basalts from the Juan de Fuca and Gorda rRidges by mass spectrometry. *Earth and Planetary Science Letters*, *96*, 134–146.
- Goldstein, S. J., Murrell, M. T., Janecky, D. R., Delaney, J. R., & Clague, D. A. (1991). Geochronology and petrogenesis of MORB from the Juan de Fuca and Gorda Ridges by 238U-230Th disequilibrium. *Earth and Planetary Science Letters*, *107*, 25–41.
- Goldstein, S. J., Murrell, M. T., & Williams, R. W. (1993). 231Pa and 23Th chronology of Mid-Ocean Ridge basalts. *Earth and Planetary Science Letters*, *115*, 151–159.
- Goldstein, S. L., Soffer, G., Langmuir, C. H., Lehnert, K. A., Graham, D. W., & Michael, P. J. (2008). Origin of a 'Southern Hemisphere' geochemical signature in the Arctic upper mantle. *Nature*, *453*, 89–93. <https://doi.org/10.1038/nature06919>
- Goss, A. R., Perfit, M. R., Ridley, W. I., Rubin, K. H., Kamenov, G. D., Soule, S. A., et al. (2010). Geochemistry of lavas from the 2005-2006 eruption at the East Pacific Rise, 9°46'N - 9°56'N: Implications for ridge crest plumbing and decadal changes in magma chamber compositions. *Geochemistry, Geophysics, Geosystems*, *11*, Q05T09. <https://doi.org/10.1029/2009GC002977>
- Graham, D. W. (2007). Submitted data set: Galapagos Spreading Center He data, Earthchem Library.
- Graham, D. W., Blichert-Toft, J., Russo, C. J., Rubin, K. H., & Albarède, F. (2006). Cryptic striations in the upper mantle revealed by hafnium isotopes in southeast Indian ridge basalts. *Nature*, *440*(7081), 199–202. <https://doi.org/10.1038/nature04582s>
- Graham, D. W., Castillo, P. R., Lupton, J. E., & Batiza, R. (1996). Correlated He and Sr isotope ratios in South Atlantic near-ridge seamounts and implications for mantle dynamics. *Earth and Planetary Science Letters*, *144*, 491–503.
- Graham, D. W., Christie, D. M., Harpp, K. S., & Lupton, J. E. (1993). Mantle plume helium in submarine basalts from the Galápagos platform. *Science*, *262*(5142), 2023–2026.
- Graham, D. W., Hanan, B. B., Lupton, J. E., Hoernle, K., Werner, R., Christie, D. M., & Sinton, J. M. (2014). Helium isotope variations and mantle plume-spreading ridge interactions along the Galápagos spreading center. A Natural Laboratory for the Earth Sciences. 393-414.
- Graham, D. W., Jenkins, W. J., Kurz, M. D., & Batiza, R. (1987). Helium isotope disequilibrium and geochronology of glassy submarine basalts. *Nature*, *326*, 384–386.
- Graham, D. W., Jenkins, W. J., Schilling, J.-G., Thompson, G., Kurz, M. D., & Humphris, S. E. (1992). Helium isotope geochemistry of Mid-Ocean Ridge basalts from the South Atlantic. *Earth and Planetary Science Letters*, *110*, 133–147.
- Graham, D. W., Johnson, K. T. M., Douglas-Priebe, L. M., & Lupton, J. E. (1999). Hotspot-ridge interaction along the Southeast Indian Ridge near Amsterdam and St. Paul Islands: Helium isotope evidence. *Earth and Planetary Science Letters*, *167*, 297–310.
- Graham, D. W., Lupton, J. E., Spera, F. J., & Christie, D. M. (2001). Upper-mantle dynamics revealed by helium isotope variations along the Southeast Indian Ridge. *Nature*, *409*, 701–703.
- Graham, D. W., Michael, P. J., & Shea, T. (2016). Extreme incompatibility of helium during mantle melting: Evidence from undegassed Mid-Ocean Ridge basalts. *Earth and Planetary Science Letters*, *454*, 192–202. <https://doi.org/10.1016/j.epsl.2016.09.016>
- Graham, D. W., Zindler, A., Kurz, M. D., Jenkins, W. J., Batiza, R., & Staudigel, H. (1988). He, Pb, Sr and Nd isotope constraints on magma genesis and mantle heterogeneity beneath young Pacific seamounts. *Contributions to Mineralogy and Petrology*, *99*, 446–463.
- Gregg, T. K. P., Fornari, D. J., Perfit, M. R., Ridley, W. I., & Kurz, M. D. (2000). Using submarine lava pillars to record Mid-Ocean Ridge eruption dynamics. *Earth and Planetary Science Letters*, *178*, 195–214.
- Gribble, R. F., Stern, R. J., Bloomer, S. H., Stüben, D., O'Hearn, T., & Newman, S. (1996). MORB mantle and subduction components interact to generate basalts in the southern Mariana Trough back-arc basin. *Geochimica et Cosmochimica Acta*, *60*, 2153–2166.
- Gribble, R. F., Stern, R. J., Newman, S., Bloomer, S. H., & O'Hearn, T. (1998). Chemical and isotopic composition of lavas from the northern Mariana Trough: Implications for magmagenesis in back-arc basins. *Journal of Petroleum*, *39*, 125–134.
- Haase, K., Brandl, P., Devey, C., Hauff, F., Melcher, B., Garbe-Schönberg, D., et al. (2016). Compositional variation and 226Ra- 230Th model ages of Axial lavas from the southern Mid-Atlantic Ridge, 8°48's. *Geochemistry, Geophysics, Geosystems*, *1*, 199–218. <https://doi.org/10.1002/2015GC006016>
- Haase, K. M. (2002). Geochemical constraints on magma sources and mixing processes in easter microplate MORB (se pacific); a case study of plume-ridge interaction. *Chemical Geology*, *182*, 335–355.



- Haase, K. M., Devey, C. W., Mertz, D. F., Stoffers, P., & Garbe-Schönberg, D. (1996). Geochemistry of lavas from Mohns Ridge, Norwegian-Greenland Sea: Implications for melting conditions and magma sources near Jan Mayen. *Contributions to Mineralogy and Petrology*, *123*, 223–237.
- Haase, K. M., Regelous, M., Duncan, R. A., Brandl, P. A., Stronck, N., & Grevenmeyer, I. (2011). Insights into mantle composition and mantle melting beneath Mid-Ocean Ridges from postspreading volcanism on the Fossil Galapagos Rise. *Geochemistry, Geophysics, Geosystems*, *12*, Q0AC11. <https://doi.org/10.1029/2010GC003482>
- Hahn, D., Castillo, P. R., & Hilton, D. R. (2009). A deep mantle source for high  $^3\text{He}/^4\text{He}$  ocean island basalts (OIB) inferred from Pacific near-ridge seamount lavas. *Geophysical Research Letters*, *36*, L20316. <https://doi.org/10.1029/2009GL040560>
- Hahn, D., Hilton, D. R., Castillo, P. R., Hawkins, J. W., Hanan, B. B., & Hauri, E. H. (2012). An overview of the volatile systematics of the Lau Basin - resolving the effects of source variation, magmatic degassing and crustal contamination. *Geochimica et Cosmochimica Acta*, *85*, 88–113.
- Hamelin, B., & Allègre, C. J. (1985). Large-scale regional units in the depleted upper mantle revealed by an isotope study of the Southwest Indian Ridge. *Nature*, *315*, 196–199. <https://doi.org/10.1038/315196a0>
- Hamelin, B., Dupre, B., & Allègre, C. J. (1985). Pb-Sr-Nd isotope data of Indian Ocean Ridges: New evidence of large-scale mapping of mantle heterogeneities. *Earth and Planetary Science Letters*, *76*, 288–298.
- Hamelin, B., Dupre, B., & Allègre, C. J. (1984). Lead-strontium isotopic variations along the East Pacific Rise and the Mid-Atlantic Ridge: A comparative study. *Earth and Planetary Science Letters*, *67*, 340–350.
- Hamelin, C., Bezos, A., Dosso, L., Escartin, J., Cannat, M., & Mevel, C. (2013). Atypically depleted upper mantle component revealed by Hf isotopes at Lucky Strike segment. *Chemical Geology*, *341*, 128–139.
- Hamelin, C., Dosso, L., Hanan, B., Barrat, J.-A., & Ondréas, H. (2010). Sr-Nd-Hf isotopes along the Pacific Antarctic Ridge from 41 to 53°S. *Geophysical Research Letters*, *37*, L10303. <https://doi.org/10.1029/2010GL042979>
- Hamelin, C., Dosso, L., Hanan, B. B., Moreira, M., Kositsky, A. P., & Thomas, M. Y. (2011). Geochemical portray of the Pacific Ridge: New isotopic data and statistical techniques. *Earth and Planetary Science Letters*, *302*, 154–162.
- Hanan, B. B., Blichert-Toft, J., Hemond, C., Sayit, K., Agranier, A., Graham, D. W., & Albarede, F. (2013). Pb and Hf isotope variations along the southeast Indian Ridge and the dynamic distribution of MORB source domains in the upper mantle. *Earth and Planetary Science Letters*, *375*, 196–208.
- Hanan, B. B., Blichert-Toft, J., Kingsley, R. H., & Schilling, J.-G. (2000). Depleted Iceland mantle plume geochemical signature: Artifact of multicomponent mixing? *Geochemistry, Geophysics, Geosystems*, *1*(4), 1003. <https://doi.org/10.1029/1999GC000009>
- Hanan, B. B., Blichert-Toft, J., Pyle, D. G., & Christie, D. M. (2004). Contrasting origins of the upper mantle revealed by hafnium and lead isotopes from the southeast Indian Ridge. *Nature*, *432*, 91–94.
- Hanan, B. B., Kingsley, R. H., & Schilling, J.-G. (1986). Pb isotope evidence in the South Atlantic for migrating ridge-hot spot interactions. *Nature*, *322*, 137–144.
- Hanan, B. B., & Schilling, J.-G. (1989). Easter microplate evolution: Pb isotope evidence. *Journal of Geophysical Research*, *94*, 7432–7448. <https://doi.org/10.1029/JB094iB06p07432>
- Hannigan, R. E., Asish, R. B., & Friedrich, T. (2001). Mantle reservoir geochemistry from statistical analysis of ICP-MS trace element data of equatorial mid-Atlantic MORB glasses. *Chemical Geology*, *175*(3–4), 397–428. [https://doi.org/10.1016/S0009-2541\(00\)00335-1](https://doi.org/10.1016/S0009-2541(00)00335-1)
- Harpp, K. S. (1995). Geochemical variations in a single basaltic flow at 9°30'N on the East Pacific Rise. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 142, pp. 9–22). College Station, Texas: Ocean Drilling Program.
- Harpp, K. S., Fornari, D. J., Geist, D. J., & Kurz, M. D. (2003). Genovesa Submarine Ridge: A manifestation of plume-ridge interaction in the northern Galapagos Islands. *Geochemistry, Geophysics, Geosystems*, *4*(9), 8511. <https://doi.org/10.1029/2003GC000531>
- Harpp, K. S., Wanless, V. D., Otto, R. H., Hoernle, K., & Werner, R. (2005). The Cocos and Carnegie aseismic Ridges: A trace element record of long-term plume-spreading center interaction. *Journal of Petroleum*, *46*, 109–133.
- Harpp, K. S., & White, W. M. (2001). Tracing a mantle plume: Isotopic and trace element variations of Galapagos seamounts. *Geochemistry, Geophysics, Geosystems*, *2*(6), 1042–1046. <https://doi.org/10.1029/2000GC000137>
- Hart, R., Dymond, J., Hogan, L. G., & Schilling, J.-G. (1983). Mantle plume noble gas component in glassy basalts from Reykjanes Ridge. *Nature*, *305*, 403–407.
- Hart, S. R. (1971). K, Rb, Cs, Sr and Ba contents and Sr isotope ratios of ocean floor basalts. *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, *268*, 573–587.
- Hart, S. R., Glassley, W. E., & Karig, D. E. (1972). Basalts and seafloor spreading behind the Mariana Island Arc. *Earth and Planetary Science Letters*, *15*, 12–18.
- Hart, S. R., Schilling, J.-G., & Powell, J. L. (1973). Basalts from Iceland and along the Reykjanes Ridge: Sr isotope geochemistry. *Nature*, *246*, 104–107.
- Hawkins, J. W., Lonsdale, P. F., Macdougall, J. D., & Volpe, A. M. (1990). Petrology of the axial ridge of the Mariana Trough backarc spreading center. *Earth and Planetary Science Letters*, *100*, 226–250.
- Hawkins, J. W., & Melchior, J. T. (1985). Petrology of Mariana Trough and Lau Basin basalts. *Journal of Geophysical Research*, *90*, 11,431–11,468. <https://doi.org/10.1029/JB090iB13p11431>
- Hays, M. R. (2004). Intra-transform volcanism along the Siqueiros fracture zone 8°20'N to 8°30'N, East Pacific Rise, Thesis Or Dissertation, Masters, University Of Florida. carbon cycle and associated redox processes through time. *Philosophical Transactions of the Royal Society, B: Biological Sciences*, *361*, 931–950. [https://doi.org/10.1016/0016-7037\(95\)00038-2](https://doi.org/10.1016/0016-7037(95)00038-2)
- Hedge, C. E., Futa, K., Engel, C. G., & Fisher, R. L. (1979). The rare earth abundances and Rb-Sr systematics of basalts, gabbro, anorthosite and minor granite rocks from the Indian Ocean rRdge system, western Indian Ocean. *Contributions to Mineralogy and Petrology*, *68*, 373–376.
- Hedge, C. E., & Peterman, Z. E. (1970). The strontium isotopic composition of basalts from the Gorda and Juan de Fuca Rises, northeastern Pacific Ocean. *Contributions to Mineralogy and Petrology*, *27*, 114–120.
- Heft, K., Gillis, K. M., Pollock, M. A., Karson, J. A., & Klein, E. M. (2008). Role of upwelling hydrothermal fluids in the development of alteration patterns at fast spreading ridges: Evidence from the sheeted dike complex at Pito Deep. *Geochemistry, Geophysics, Geosystems*, *9*, Q05O07. <https://doi.org/10.1029/2007GC001926>
- Hegner, E., & Tatsumoto, M. (1987). Pb, Sr, and Nd isotopes in basalts and sulfides from the Juan de Fuca Ridge. *Journal of Geophysical Research*, *92*, 11,380–11,386. <https://doi.org/10.1029/JB092iB11p11380>
- Hegner, E., & Tatsumoto, M. (1989). Pb, Sr, and Nd isotopes in seamount basalts from the Juan de Fuca Ridge and Kodiak-Bowie Seamount Chain, northeast Pacific. *Journal of Geophysical Research*, *94*, 17,839–17,846. <https://doi.org/10.1029/JB094iB12p17839>

- Hekinian, R. (1971). Chemical and mineralogical differences between abyssal hill basalts and ridge tholeiites in the eastern Pacific Ocean. *Marine Geology*, *11*, 77.
- Hekinian, R., Bideau, R. D., Hebert, R., & Niu, Y. (1995). Magmatism in the Garrett Transform Fault (East Pacific Rise near 13°27'S). *Journal of Geophysical Research*, *100*(B6), 10,163–10,185. <https://doi.org/10.1029/94JB02125>
- Hekinian, R., & Fouquet, Y. (1985). Volcanism and metallogenesis of axial and off-axial structures on the East Pacific Rise near 13°N. *Economic Geology*, *80*, 221–249.
- Hekinian, R., Francheteau, J., Armijo, R., Cogne, J. P., Constantin, M., Girardeau, J., et al. (1996). Petrology of the Easter microplate region in the South Pacific. *Journal of Volcanology and Geothermal Research*, *72*, 259–289.
- Hekinian, R., & Hoffert, M. (1975). Rate of palagonitization and manganese coating on basaltic rocks from the Rift Valley in the Atlantic Ocean near 36deg50'N. *Marine Geology*, *19*, 91–109.
- Hekinian, R., Moore, J. G., & Bryan, W. B. (1976). Volcanic rocks and processes of the Mid-Atlantic Ridge Rift Valley near 36°49'N. *Contributions to Mineralogy and Petrology*, *58*, 83–110.
- Hekinian, R., Stoffers, P., Ackerman, D., Revillon, S., Maia, M., & Bohn, M. (1999). Ridge-hotspot interaction: The Pacific-Antarctic Ridge and the Foundation Seamounts. *Marine Geology*, *160*, 199–223.
- Hekinian, R., Stoffers, P., Devey, C. W., Ackerman, D., Hemond, C., O'Connor, J., et al. (1997). Intraplate versus ridge volcanism on the Pacific-Antarctic Ridge near 37°S–111degW. *Journal of Geophysical Research*, *102*(B6), 12,265–12,286. <https://doi.org/10.1029/96JB03856>
- Hekinian, R., Thompson, G., & Bideau, R. D. (1989). Axial and off-axial heterogeneity of basaltic rocks from the East Pacific Rise at 12°35'N–12°51'N. *Journal of Geophysical Research*, *94*, 17,437–17,463. <https://doi.org/10.1029/1998JB900048>
- Hekinian, R., & Walker, D. (1987). Diversity and spatial zonation of volcanic rocks from the East Pacific Rise near 21°N. *Contributions to Mineralogy and Petrology*, *96*, 265–280.
- Hemond, C., Hofmann, A. W., Vlastelic, I., & Nauret, F. (2006). Origin of MORB enrichment and relative trace element compatibilities along the Mid-Atlantic Ridge between 10 and 24°N. *Geochemistry, Geophysics, Geosystems*, *7*, Q12010. <https://doi.org/10.2029/2006GC1317>
- Herbrich, A., Hauff, F., Hoernle, K., Werner, R., Garbe-Schönberg, D., & White, S. M. (2016). A 1.5 ma record of plume-ridge interaction at the western Galápagos Spreading Center (91°40'–92°00'W). *Geochimica et Cosmochimica Acta*, *185*, 141–159. <https://doi.org/10.1016/j.gca.2016.04.036>
- Hermes, O. D., & Schilling, J.-G. (1976). Olivine from Reykjanes Ridge and Iceland tholeiites, and its significance to the two-mantle source model. *Earth and Planetary Science Letters*, *28*, 345–355.
- Hertogen, J., Janssens, M. J., & Palme, H. (1980). Trace elements in ocean ridge basalt glasses: Implications for fractionations during mantle evolution and petrogenesis. *Geochimica et Cosmochimica Acta*, *44*, 2125–2143.
- Hildenbrand, A., Weis, D., Madureira, P., & Marques, A. F. A. (2014). Recent plate re-organization at the Azores Triple Junction: Evidence from combined geochemical and geochronological data on Faial, S. Jorge and Terceira volcanic islands. *Lithos*, *210–211*, 27–39. <https://doi.org/10.1016/j.lithos.2014.09.009>
- Hilton, D. R., Thirlwall, M. F., Taylor, R. N., Murton, B. J., & Nichols, A. (2000). Controls on magmatic degassing along the Reykjanes Ridge with implications for the helium paradox. *Earth and Planetary Science Letters*, *183*, 43–50.
- Hinds, J. A. (2005). Construction of the oceanic crustal layer 2A: A detailed petrographic and geochemical study of lavas from 9deg30 'N and 9°50 'N East Pacific Rise. Thesis Or Dissertation, Master'S, University Of Florida; Monographi.
- Hiyagon, H., Ozima, M., Marty, B., Zashu, S., & Sakai, H. (1992). Noble gases in submarine glasses from mid-ocean ridges and Loihi seamount: Constraints on the early history of the Earth. *Geochimica et Cosmochimica Acta*, *56*, 1301–1316.
- Hoernle, K., Hauff, F., Kokfelt, T. F., Haase, K. M., Garbe-Schönberg, D., & Werner, R. (2011). On- and off-axis chemical heterogeneities along the South Atlantic Mid-Ocean-Ridge (5–11°S): Shallow or deep recycling of ocean crust and/or intraplate volcanism? *Earth and Planetary Science Letters*, *306*, 86–97.
- Hofmann, A. W., Jochum, K. P., Seufert, H. M., & White, W. M. (1986). Nb and Pb in oceanic basalts: New constraints on mantle evolution. *Earth and Planetary Science Letters*, *79*, 33–45.
- Hofmann, A. W., & White, W. M. (1983). Ba, Rb and Cs in the Earth's mantle. *Zeitschrift für Naturforschung*, *38A*, 256–266.
- Holm, P. M. (2002a). Sr, Nd and Pb isotopic composition of in situ lower crust at the Southwest Indian Ridge: Results from ODP Leg 176. *Chemical Geology*, *184*, 195–216.
- Holm, P. M. (2002b). Data report: On the composition of the lower ocean crust-major and trace element analyses of gabbroic rocks from Hole 735B, 500–1500 mbsf. In J. H. Natland, H. J. B. Dick, D. J. Miller, & R. P. von Herzen (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 176, pp. 1–13). College Station, Texas: Ocean Drilling Program.
- Humler, E., & Whitechurch, H. (1988). Petrology of basalts from the Central Indian Ridge (lat. 25°23'S, long. 70°04'E): Estimates of frequencies and fractional volumes of magma injections in a two-layered reservoir. *Earth and Planetary Science Letters*, *88*, 169–181.
- Humphris, S. E., Thompson, G., Schilling, J.-G., & Kingsley, R. H. (1985). Petrological and geochemical variations along the Mid-Atlantic Ridge between 46degS and 32degS: Influence of the Tristan de Cunha mantle plume. *Geochimica et Cosmochimica Acta*, *49*, 1445–1464.
- Ikeda, Y., Nagao, K., Stern, R. J., Yuasa, M., & Newman, S. (1998). Noble gas in pillow basalt glasses from the northern Mariana Trough back-arc basin. *The Island Arc*, *7*, 471–478.
- Ingle, S., Ito, G., Mahoney, J. J., Chazey Iii, W., Sinton, J. M., Rotella, M., & Christie, D. M. (2010). Mechanisms of geochemical and geophysical variations along the western Galapagos Spreading Center. *Geochemistry, Geophysics, Geosystems*, *11*, Q04003. <https://doi.org/10.1029/2009GC002694>
- Ito, E., White, W. M., & Goepel, C. (1987). The O, Sr, Nd, and Pb isotope geochemistry of Mid-Ocean Ridge basalts. *Chemical Geology*, *62*, 157–176.
- Ito, E., White, W. M., von Drach, V., Hofmann, A. W., & James, D. E. (1980). Isotopic studies of ocean ridge basalts. *Carn Inst, Dtm, Annual Rep Director*, 465–471.
- Jambon, A., Deruelle, B., Dreibus, G., & Pineau, F. (1995). Chlorine and bromine abundance in MORB: The contrasting behaviour of the Mid-Atlantic Ridge and East Pacific Rise and implications for chlorine geodynamic cycle. *Chemical Geology*, *126*, 101–117.
- Jambon, A., & Zimmermann, J. L. (1990). Water in oceanic basalts: Evidence for dehydration of recycled crust. *Earth and Planetary Science Letters*, *101*, 323–331.
- Jambon, A., & Zimmermann, L. (1987). Major volatiles from a North Atlantic MORB glass and calibration to He: A size fraction analysis. *Chemical Geology*, *62*, 177–189.
- Janney, P. E., Le Roex, A. P., & Carlson, R. W. (2005). Hafnium isotope and trace element constraints on the nature of mantle heterogeneity beneath the central Southwest Indian Ridge (13°E to 47°E). *Journal of Petrology*, *46*(12), 2427–2464. <https://doi.org/10.1093/petrology/egi060>

- Jenner, F. E., Arculus, R. J., Mavrogenes, J. A., Dyriw, N., Nebel, O., & Hauri, E. H. (2012). Chalcophile element systematics in volcanic glasses from the northwestern Lau Basin. *Geochemistry, Geophysics, Geosystems*, 13, Q06014. <https://doi.org/10.1029/2012GC004088>
- Jenner, F. E., Oneill, H. S. C., Arculus, R. J., & Mavrogenes, J. A. (2010). The magnetite crisis in the evolution of arc-related magmas and the initial concentration of Au, Ag and Cu. *Journal of Petroleum*, 51, 2445–2464.
- Jenner, G. A., Cawood, P. A., Rautenschlein, M., & White, W. M. (1987). Composition of back-arc basin volcanics, Valu Fa s, Lau Basin: Evidence for a slab-derived component in their mantle source. *Journal of Volcanology and Geothermal Research*, 32, 209–222.
- Jochum, K. P., & Hofmann, A. W. (1997). Constraints on Earth evolution from antimony in mantle-derived rocks. *Chemical Geology*, 139, 39–49.
- Jochum, K. P., Hofmann, A. W., Ito, E., Seufert, H. M., & White, W. M. (1983). K, U and Th in Mid-Ocean Ridge basalt glasses and heat production, K/U and K/Rb in the mantle. *Nature*, 306, 431–436.
- Jochum, K. P., Hofmann, A. W., & Seufert, H. M. (1993). Tin in mantle-derived rocks: Constraints on Earth evolution. *Geochimica et Cosmochimica Acta*, 57, 3585–3595.
- Jochum, K. P., & Verma, S. P. (1996). Extreme enrichment of Sb, Tl and other trace elements in altered MORB. *Chemical Geology*, 130, 289–299.
- Johnson, J. R. (1979). Transitional basalts and tholeiites from the East Pacific Rise, 9°N. *Journal of Geophysical Research*, 84(B4), 1635–1651. <https://doi.org/10.1029/JB084iB04p01635>
- Johnson, K. T. M., & Sinton, J. M. (1990). Petrology, tectonic setting, and the formation of back-arc basin basalts in the north Fiji Basin. *Geologisches Jahrbuch*, D92, 517–545.
- Johnson, R. W., Jaques, A. L., Langmuir, C. H., Perfit, M. R., Staudigel, H., Dunkley, P. N., et al. (1987). Ridge subduction and forearc volcanism: Petrology and geochemistry of rocks dredged from the western Solomon Arc and Woodlark Basin. In B. Taylor & N. F. Exon (Eds.), *Marine Geology, Geophysics, and Geochemistry of the Woodlark Basin - Solomon Islands*, Earth Science Series (Vol. 7, pp. 155–226). Houston, TX: Circum-Pacific Council for Energy and Mineral Resources.
- Juteau, T., Eissen, J. P., Francheteau, J., Needham, D., Choukroune, P., Rangin, C., et al. (1980). Homogeneous basalts from the East Pacific Rise at 21°N: Steady state magma reservoirs at moderately fast spreading centers. *Oceanologica Acta*, 3.
- Kadko, D., Koski, R. A., Tatsumoto, M., & Bouse, R. (1985). An estimate of hydrothermal fluid residence times and vent chimney growth rates based on 210 Pb/Pb ratios and mineralogic studies of sulfides dredged from the Juan de Fuca Ridge. *Earth and Planetary Science Letters*, 76, 35–44.
- Kamenetsky, V. S., Binns, R. A., Gemmill, J. B., Crawford, A. J., Mernagh, T. P., Maas, R., & Steele, D. (2001). Parental basaltic melts and fluids in eastern Manus backarc Basin: Implications for hydrothermal mineralisation. *Earth and Planetary Science Letters*, 184, 685–702.
- Kamenetsky, V. S., Maas, R., Sushchevskaya, N. M., Norman, M. D., Cartwright, I., & Peyve, A. (2001). Remnants of Gondwanan continental lithosphere in oceanic upper mantle: Evidence from the South Atlantic Ridge. *Geology*, 29, 243–246.
- Kamenetsky, V. S., Crawford, A. J., Eggins, S. M., & Mühe, R. K. (1997). Phenocryst and melt inclusion chemistry of near-axis seamounts, Valu Fa Ridge, Lau Basin: Insight into mantle wedge melting and the addition of subduction components. *Earth and Planetary Science Letters*, 151, 205–223.
- Kamenetsky, V. S., Eggins, S. E., Crawford, A. J., Green, D. H., Gasparon, M., & Falloon, T. J. (1998). Calcic melt inclusions in primitive olivine at 43°N MAR: Evidence for melt-rock reaction/melting involving clinopyroxene-rich lithologies during MORB generation. *Earth and Planetary Science Letters*, 160, 115–132.
- Kamenetsky, V. S., Everard, J. L., Crawford, A. J., Varne, R., Eggins, S. M., & Lanyon, R. (2000). Enriched end-member of primitive MORB melts: Petrology and geochemistry of glasses from Macquarie Island (SW Pacific). *Journal of Petroleum*, 41, 411–430.
- Karsten, J. L. (1988). Spatial and temporal variations in the petrology, morphology and tectonics of a migrating spreading center: The endeavor segment, Juan de Fuca Ridge, (Thesis Or Dissertation). Doctoral, University Wash; Monographic.
- Karsten, J. L., Delaney, J. R., Rhodes, J. M., & Lias, R. A. (1990). Spatial and temporal evolution of magmatic systems beneath the endeavor segment, Juan de Fuca Ridge: Tectonic and petrologic constraints. *Journal of Geophysical Research*, 95, 19,235–19,256. <https://doi.org/10.1029/JB095iB12p19235>
- Karsten, J. L., Klein, E. M., & Sherman, C. (1996). Subduction zone geochemical characteristics in ocean ridge basalts from the southern Chile Ridge: Implications of modern ridge subduction systems for the Archaen. *Lithos*, 37, 143–161.
- Kay, R., Hubbard, N. J., & Gast, P. W. (1970). Chemical characteristics and origin of oceanic ridge volcanic rocks. *Journal of Geophysical Research*, 75, 1585–1613. <https://doi.org/10.1029/JB075i008p01585>
- Kay, R., Sun, S. S., & Lee-Hu, C. N. (1978). Pb and Sr isotopes in volcanic rocks from the Aleutian Islands and Pribilof Islands, Alaska. *Geochimica et Cosmochimica Acta*, 42(3), 263–273.
- Keller, N. S., Arculus, R. J., Hermann, J., & Richards, S. (2008). Submarine back-arc lava with arc signature: Fonualei Spreading Center, northeast Lau Basin, Tonga. *Journal of Geophysical Research*, 113, B08S07. <https://doi.org/10.1029/2007JB005451>
- Kelley, K. A., & Cottrell, E. (2009). Water and the oxidation state of subduction zone magmas. *Science*, 325, 605–607.
- Kempton, P. D., Hawkesworth, C. J., & Fowler, M. (1991). Geochemistry and isotopic composition of Gabbros from Layer 3 of the Indian Ocean crust, Hole 735B. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 118, pp. 127–143). College Station, Texas: Ocean Drilling Program.
- Kempton, P. D., & Hunter, A. G. (1997). A Sr-, Nd-, Pb-, O-isotope study of plutonic rocks from MARK, Leg 153: Implications for mantle heterogeneity and magma chamber processes. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 153, pp. 305–319). College Station, Texas: Ocean Drilling Program.
- Kempton, P. D., & Casey, J. F. (1997). Petrology and geochemistry of crosscutting diabase dikes, Sites 920 and 921. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 153, pp. 363–377). College Station, Texas: Ocean Drilling Program.
- Kempton, P. D., Fitton, J. G., Saunders, A. D., Nowell, G. M., Taylor, R. N., Hardarson, B. S., & Pearson, G. (2000). The Iceland plume in space and time: A Sr-Nd-Pb-Hf study of the North Atlantic rifted margin. *Earth and Planetary Science Letters*, 177, 255–271.
- Kempton, P. D., Pearce, J. A., Barry, T. L., Fitton, J. G., Langmuir, C. H., & Christie, D. M. (2002). Sr-Nd-Pb-Hf isotope results from ODP Leg 187: Evidence for mantle dynamics of the Australian-Antarctic discordance and origin of the Indian MORB source. *Geochemistry, Geophysics, Geosystems*, 3(12), 1074. <https://doi.org/10.1029/2002GC000320>
- Kent, A. J. R., Peate, D. W., Newman, S., Stolper, E., & Pearce, J. A. (2002). Chlorine in submarine glasses from the Lau Basin: Seawater contamination and constraints on the composition of slab-derived fluids. *Earth and Planetary Science Letters*, 202, 361–377.
- Kingsley, R. H., Blichert-Toft, J., Fontignie, D., & Schilling, J.-G. (2007). Hafnium, neodymium, and strontium isotope and parent-daughter element systematics in basalts from the plume-ridge interaction system of the Salas y Gomez Seamount Chain and Easter Microplate. *Geochemistry, Geophysics, Geosystems*, 8, Q04005. <https://doi.org/10.1029/2006GC001401>
- Kingsley, R. H., & Schilling, J.-G. (1995). Carbon in Mid-Atlantic Ridge basalt glasses from 28°N to 63°N: Evidence for a carbon-enriched Azores mantle plume. *Earth and Planetary Science Letters*, 129, 31–53.

- Kingsley, R. H., & Schilling, J.-G. (1998). Plume-ridge interaction in the Easter-Salas y Gomez Seamount Chain--Easter Microplate system: Pb isotope evidence. *Journal of Geophysical Research*, *103*, 24,159–24,177. <https://doi.org/10.1029/98JB01496>
- Kingsley, R. H., Schilling, J.-G., Dixon, J. E., Swart, P., Poreda, R., & Simons, K. (2002). D/H ratios in basalt glasses from the Salas y Gomez mantle plume interacting with the East Pacific Rise: Water from old D-rich recycled crust or primordial water from the lower mantle? *Geochemistry, Geophysics, Geosystems*, *3*(4), 1025. <https://doi.org/10.1029/2001GC000199>
- Klein, E. M., Langmuir, C. H., & Staudigel, H. (1991). Geochemistry of basalts from the Southeast Indian Ridge, 115°E–138°E. *Journal of Geophysical Research*, *96*(B2), 2089–2107. <https://doi.org/10.1029/90JB01384>
- Klein, E. M., Langmuir, C. H., Zindler, A., Staudigel, H., & Hamelin, B. (1988). Isotope evidence of a mantle convection boundary at the Australian Antarctic Discordance. *Nature*, *333*, 623–629.
- Kokfelt, T. F., Lundstrom, C. C., Hoernle, K., Hauff, F., & Werner, R. (2005). Plume-ridge interaction studied at the Galápagos Spreading Center: Evidence from 226Ra–230Th–238U and 231Pa–235U isotopic disequilibria. *Earth and Planetary Science Letters*, *234*(1–2), 165–187. <https://doi.org/10.1016/j.epsl.2005.02.031>
- Krolikowska-Ciaglo, S., Deyhle, A. T., Hauff, F., & Hoernle, K. (2007). Boron isotope geochemistry and U-Pb systematics of altered MORB from the Australian Antarctic Discordance (ODP Leg 187). *Chemical Geology*, *242*, 455–469.
- Krolikowska-Ciaglo, S., Hauff, F., & Hoernle, K. (2005). Sr-Nd isotope systematics in 14–28 MA low-temperature altered Mid-Ocean Ridge basalt from the Australian Antarctic Discordance, Ocean Drilling Program Leg 187. *Geochemistry, Geophysics, Geosystems*, *6*, Q01001. <https://doi.org/10.1029/2004GC000802>
- Kruber, C., Thorseth, I., & Pedersen, R. B. (2008). Seafloor alteration of basaltic glass: Textures, geochemistry, and endolithic microorganisms. *Geochemistry, Geophysics, Geosystems*, *9*, Q12002. <https://doi.org/10.1029/2008GC002119>
- Kumagai, H., Dick, H. J. B., & Kaneoka, I. (2003). Noble gas signatures of Abyssal gabbros and peridotites at an Indian Ocean core complex. *Geochemistry, Geophysics, Geosystems*, *4*(12), 9107. <https://doi.org/10.1029/2003GC000540>
- Kumagai, H., & Kaneoka, I. (1998). Variations of noble gas abundances and isotope ratios in a single MORB pillow. *Geophysical Research Letters*, *25*, 3891–3894. <https://doi.org/10.1029/1998GL900047>
- Kumagai, H., & Kaneoka, I. (2003). Relationship between submarine MORB glass textures and atmospheric component of MORBs. *Chemical Geology*, *200*, 1–24.
- Kurz, M. D. (1982). Helium isotope geochemistry of oceanic volcanic rocks: Implications for mantle heterogeneity and degassing. *Whoi Techn Reports*. <https://doi.org/10.1575/1912/2211>
- Kurz, M. D., Curtice, J., Fornari, D. J., Geist, D. J., & Moreira, M. (2009). Primitive neon from the center of the Galápagos hotspot. *Earth and Planetary Science Letters*, *286*, 23–34. <https://doi.org/10.1016/j.epsl.2009.06.008>
- Kurz, M. D., Warren, J. E., & Curtice, J. (2009). Mantle deformation and noble gases: Helium and neon in oceanic mylonites. *Chemical Geology*, *266*, 10–18.
- Kurz, M. D., Jenkins, W. J., Schilling, J.-G., & Hart, S. R. (1982). Helium isotopic variations in the mantle beneath the central North Atlantic Ocean. *Earth and Planetary Science Letters*, *58*, 1–14.
- Kurz, M. D., Le Roex, A. P., & Dick, H. J. B. (1998). Isotope geochemistry of the oceanic mantle near the Bouvet Triple Junction. *Geochimica et Cosmochimica Acta*, *62*, 841–852.
- Kurz, M. D., Moreira, M., Curtice, J., Lott, I., Mahoney, J. J., & Sinton, J. M. (2005). Correlated helium, neon, and melt production on the super-fast spreading East Pacific Rise near 17°S. *Earth and Planetary Science Letters*, *232*, 125–142.
- Kutza, S. (2002). Petrology and geochemistry of Mid-Ocean Ridge basalts from the East Pacific Rise (9°24' and 10°06' N): Implications for off-axis magmatic processes. Thesis Or Dissertation, Master'S, University Of Florida.
- Kyser, T. K., & O'Neil, J. R. (1984). Hydrogen isotope systematics of submarine basalts. *Geochimica et Cosmochimica Acta*, *48*, 2123–2133.
- Labidi, J., Cartigny, P., Bircik, J. L., Assayag, N., & Bourrand, J. (2012). Determination of multiple sulfur isotopes in glasses: A reappraisal of the MORB δ34S. *Chemical Geology*, *334*, 189–198.
- Labidi, J., Cartigny, P., Hamelin, C., Moreira, M., & Dosso, L. (2014). Sulfur isotope budget (<sup>32</sup>S, <sup>33</sup>S, <sup>34</sup>S and <sup>36</sup>S) in Pacific-Antarctic ridge basalts: A record of mantle source heterogeneity and hydrothermal sulfide assimilation. *Geochimica et Cosmochimica Acta*, *133*, 47–67. <https://doi.org/10.1016/j.gca.2014.02.023>
- Lagabriele, Y., Auzende, J. M., Eissen, J. P., Janin, M. C., & Cottin, J. (1994). Geology and geochemistry of a 800 m section through young upper oceanic crust in the North Fiji Basin (Southwest Pacific). *Marine Geology*, *116*, 113–132.
- Langmuir, C. H. (1999). Submitted data set: Major and trace element composition of basalt glasses from the CHEPR Cruise, East Pacific Rise. Earthchem Library.
- Langmuir, C. H., Bender, J. F., Bence, A. E., Hanson, G. N., & Taylor, S. R. (1977). Petrogenesis of basalts from the FAMOUS area: Mid-Atlantic Ridge. *Earth and Planetary Science Letters*, *36*, 133–156.
- Laschek, D. (1985). Geochemische untersuchungen an basalten vom Galapagos Spreading Center und vom East Pacific Rise, (Thesis Or Dissertation). Doctoral, University Karlsruhe; Monographic.
- Lawson, K., Searle, R. C., Pearce, J. A., Browning, P., & Kempton, P. D. (1996). Detailed volcanic geology and geochemistry of the MARNOCK area (Mid-Atlantic Ridge north of Kane Transform). In C. J. Macleod, P. A. Taylor, & C. L. Walker (Eds.), *Tectonic, Magmatic, Hydrothermal and Biological Segmentation of Mid-Ocean Ridges, Geological Society of America Special Publications* (pp. 118, 61–102).
- Layne, G., Kent, A. J. R., & Bach, W. (2009). 37Cl systematics of a backarc spreading system: The Lau Basin. *Geology*, *37*, 427–430.
- Le Roex, A. P. (1998). Submitted data set: Major and trace element and mineral compositions of basalts from the Agulhas 32 Cruise, southern Mid-Atlantic Ridge.
- Le Roex, A. P., Dick, H. J. B., & Fisher, R. L. (1989). Petrology and geochemistry of MORB from 25°E to 46°E along the Southwest Indian Ridge - Evidence for contrasting styles of mantle enrichment. *Journal of Petrology*, *30*, 947–986. <https://doi.org/10.1093/petrology/30.4.947>
- Le Roex, A. P., Dick, H. J. B., Erlank, A. J., Reid, A. M., Frey, F. A., & Hart, S. R. (1983). Geochemistry, mineralogy and petrogenesis of lavas erupted along the Southwest Indian Ridge between the Bouvet Triple Junction and 11°E. *Journal of Petroleum*, *24*, 267–318.
- Le Roex, A. P., Dick, H. J. B., Levent, G., Reid, A. M., & Erlank, A. J. (1987). Local and regional heterogeneity in MORB from the Mid-Atlantic Ridge between 54.5°S and 51°S: Evidence for geochemical enrichment. *Geochimica et Cosmochimica Acta*, *51*, 541–555.
- Le Roex, A. P., Dick, H. J. B., Reid, A. M., & Erlank, A. J. (1982). Ferrobasalt from the Spiess Ridge Segment of the Southwest Indian Ridge. *Earth and Planetary Science Letters*, *60*, 437–451.
- Le Roex, A. P., Dick, H. J. B., Reid, A. M., Frey, F. A., Erlank, A. J., & Hart, S. R. (1985). Petrology and geochemistry of basalts from the American-Antarctic Ridge, Southern Ocean: Implications for the westward influence of the Bouvet mantle plume. *Contributions to Mineralogy and Petrology*, *90*, 367–380.

- Le Roex, A. P., Dick, H. J. B., & Watkins, R. T. (1992). Petrogenesis of anomalous K-enriched MORB from the Southwest Indian Ridge: 11°53'E to 14°38'E. *Contributions to Mineralogy and Petrology*, *110*, 253–268.
- Le Roex, A. P., Erlank, A. J., & Needham, H. D. (1981). Geochemical and mineralogical evidence for the occurrence of at least three distinct magma types in the FAMOUS region. *Contributions to Mineralogy and Petrology*, *77*, 24–37.
- Le Roex, P. J., Le Roex, A. P., & Schilling, J.-G. (2002a). Crystallization processes beneath the southern Mid-Atlantic Ridge (40–55°S): Evidence for high-pressure initiation of crystallization. *Contributions to Mineralogy and Petrology*, *142*, 582–602.
- Le Roex, P. J., Le Roex, A. P., & Schilling, J.-G. (2002b). MORB melting processes beneath the southern Mid-Atlantic Ridge (40–55°S): A role for pyroxenite and residual garnet. *Contributions to Mineralogy and Petrology*, *144*, 206–229.
- Le Roex, P. J., Le Roex, A. P., Schilling, J.-G., Shimizu, N., Perkins, W. W., & Pearce, N. J. G. (2001). Mantle heterogeneity beneath the southern Mid-Atlantic Ridge: Trace element evidence for contamination of ambient asthenospheric mantle. *Earth and Planetary Science Letters*, *203*, 479–498.
- Le Roex, P. J., Shirey, S. B., Hauri, E. H., & Perfit, M. R. (2006). The effects of variable sources, processes and contaminants on the composition of northern EPR MORB (8–10°N and 12–14°N): Evidence from volatiles (H<sub>2</sub>O, CO<sub>2</sub>, S) and halogens (F, Cl). *Earth and Planetary Science Letters*, *251*, 209–231. <https://doi.org/10.1016/j.epsl.2006.09.012>
- Le Roux, P. J. (2000). The geochemistry of selected Mid-Ocean Ridge basalts from the southern Mid-Atlantic Ridge (40–55°S), (Thesis or Dissertation, Doctoral). University Cape Town; Monographic.
- Le Voyer, M., Cottrell, E., Kelley, K. A., Brounce, M., & Hauri, E. H. (2014). The effect of primary versus secondary processes on the volatile content of MORB glasses: An example from the equatorial Mid-Atlantic Ridge (5°N–3°S). *Journal of Geophysical Research: Solid Earth*, *120*, 125–144. <https://doi.org/10.1002/2014JB011160>
- Lecuyer, C., & Gruau, G. (1996). Oxygen and strontium isotope compositions of Hess Deep Gabbros (Holes 894F and 894G): High-temperature interaction of seawater with the oceanic crust layer 3. In C. Mevel, K. M. Gillis, J. F. Allan, & P. S. Meyer (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 147, pp. 227–234). College Station, Texas: Ocean Drilling Program.
- Leybourne, M. I., & van Wagoner, N. A. (1991). Heck and Heckle seamounts, northeast Pacific Ocean: High extrusion rates of primitive and highly depleted Mid-Ocean Ridge basalt on off-ridge seamounts. *Journal of Geophysical Research*, *96*, 16,275–16,293. <https://doi.org/10.1029/91JB00595>
- Lias, R. A. (1986). Geochemistry and petrogenesis of basalts erupted along the Juan de Fuca Ridge, Thesis or Dissertation, Doctoral, University Mass, Amherst; Monographic.
- Lock, G. (1992). Character and distribution of the Indian Ocean domain: A study of the mantle source compositions of Lau Basin volcanics (SW Pacific) and the Indian Ocean Mid-Ocean Ridge basalts, (Thesis or Dissertation, Doctoral). University Koeln; Monographic.
- Lopez, M. M., Perez, J. R., Urrutia, J. F., Verma, S. P., & Terrell, D. J. (1978). Geochemistry and petrology of some volcanic rocks dredged from the Gulf of California. *Geochemical Journal*, *12*, 127–132.
- Lundstrom, C. C., Sampson, D. E., Perfit, M. R., Gill, J. B., & Williams, Q. (1999). Insights into Mid-Ocean Ridge basalt petrogenesis: U-series disequilibria from the Siqueiros Transform, Lamont Seamounts, and the East Pacific Rise. *Journal of Geophysical Research*, *104*, 13,035–13,048. <https://doi.org/10.1029/1999JB900081>
- Lupton, J. E., Arculus, R. J., Evans, L. J., & Graham, D. W. (2012). Mantle hotspot neon in basalts from the Northwest Lau Back-arc Basin. *Geophysical Research Letters*, *39*, L08308. <https://doi.org/10.1029/2012GL051201>
- Lupton, J. E., Arculus, R. J., Greene, R. R., Evans, L. J., & Goddard, C. I. (2009). Helium isotope variations in seafloor basalts from the Northwest Lau Back-arc Basin: Mapping the influence of the Samoan hotspot. *Journal of Geophysical Research*, *36*, L17313. <https://doi.org/10.1029/2009GL039468>
- Lupton, J. E., & Craig, H. (1975). Excess <sup>3</sup>He in oceanic basalts: Evidence for terrestrial primordial helium. *Earth and Planetary Science Letters*, *26*, 133–139.
- Lupton, J. E., Graham, D. W., Delaney, J. R., & Johnson, H. P. (1993). Helium isotope variations in Juan de Fuca Ridge basalts. *Geophysical Research Letters*, *20*, 1851–1854. <https://doi.org/10.1029/93GL01271>
- Lustrino, M. (2006). Composition, petrography, and mineral chemistry of ODP Site 1224 Eocene ferrobasalts (Leg 200; North Pacific Ocean). In J. Kasahara, R. A. Stephen, G. D. Acton, & F. A. Frey (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 200, pp. 1–36). College Station, Texas: Ocean Drilling Program.
- Lytle, M., Kelley, K. A., Hauri, E. H., Gill, J. B., Papia, D., & Arculus, R. J. (2012). Tracing mantle sources and Samoan Arc influence in the northwestern Lau Back-arc Basin. *Geochemistry, Geophysics, Geosystems*, *13*, Q10019. <https://doi.org/10.1029/2012GC004233>
- Macdougall, J. D. (1976). Fission track studies of basalt from Leg 34. *Initial Reports of the Deep Sea Drilling Project*, *34*, 455.
- Macdougall, J. D., & Lugmair, G. W. (1985). Extreme isotopic homogeneity among basalts from the southern East Pacific Rise: Mantle or mixing effect. *Nature*, *313*, 209–211.
- Macdougall, J. D., & Lugmair, G. W. (1986). Sr and Nd isotopes in basalts from the East Pacific Rise: Significance for mantle heterogeneity. *Earth and Planetary Science Letters*, *77*, 273–284.
- Machado, N., Ludden, J. N., Brooks, J. N., & Thompson, G. (1982). Fine-scale isotopic heterogeneity in the sub-Atlantic mantle. *Nature*, *295*, 226–228.
- Machida, S., Orihashi, Y., Magnani, M., Neo, N., Wilson, S., Tanimizu, M., et al. (2014). Regional mantle heterogeneity regulates melt production along the Reunion hotspot-influenced Central Indian Ridge. *Geochemical Journal*, *48*, 433–449. <https://doi.org/10.2343/geochemj.2.0320>
- Macpherson, C. G., Hilton, D. R., & Hammerschmidt, K. (2010). No slab-derived CO<sub>2</sub> in Mariana Trough back-arc basalts: Implications for carbon subduction and for temporary storage of CO<sub>2</sub> beneath slow spreading ridges. *Geochemistry, Geophysics, Geosystems*, *11*, Q11007. <https://doi.org/10.1029/2010GC003293>
- Macpherson, C. G., Hilton, D. R., Matthey, D. P., & Sinton, C. W. (2000). Evidence for an 18O-depleted mantle plume from contrasting 18O/16O ratios of back-arc lavas from the Manus Basin and Mariana Trough. *Earth and Planetary Science Letters*, *176*, 171–183.
- Mahoney, J. J., Graham, D. W., Christie, D. M., Johnson, K. T. M., Hall, L. S., & Vonderhaar, D. L. (2002). Between a hotspot and a cold spot: Isotopic variation in the Southeast Indian Ridge asthenosphere, 86°E–118°E. *Journal of Petrology*, *43*, 1155–1176.
- Mahoney, J. J., Le Roex, A. P., Peng, Z., Fisher, R. L., & Natland, J. H. (1992). Southwestern limits of Indian Ocean ridge mantle and the origin of low <sup>206</sup>Pb/<sup>204</sup>Pb mid-ocean ridge basalt: Isotope systematics of the central southwest Indian Ridge (17°–50°E). *Journal of Geophysical Research*, *97*, 19,771–19,790. <https://doi.org/10.1029/92JB01424>
- Mahoney, J. J., Natland, J. H., White, W. M., Poreda, R. J., Bloomer, S. H., Fisher, R. L., & Baxter, A. N. (1989). Isotopic and geochemical provinces of the western Indian Ocean Spreading Centers. *Journal of Geophysical Research*, *94*, 4033–4052. <https://doi.org/10.1029/JB094iB04p04033>
- Mahoney, J. J., Sinton, J. M., Kurz, M. D., Macdougall, J. D., Spencer, K., & Lugmair, G. W. (1993). Isotope and trace element characteristics of a super-fast spreading ridge: East Pacific Rise 13–23°S. *Earth and Planetary Science Letters*, *121*, 171–191.

- Maia, M., Hemond, C., & Gente, P. (2001). Contrasted interactions between plume, upper mantle, and lithosphere: Foundation chain case. *Geochemistry, Geophysics, Geosystems*, 2, 1028–1029. <https://doi.org/10.1029/2000GC000117>
- Makishima, A., & Masuda, A. (1994). Ce isotope ratios of N-type MORB. *Chemical Geology*, 118, 1–8.
- Mallick, S., Dick, H. J. B., Sachi-Kocher, A., & Salters, V. J. M. (2014). Isotope and trace element insights into heterogeneity of Subridge mantle. *Geochemistry, Geophysics, Geosystems*, 15, 1–16. <https://doi.org/10.1002/2014GC005314>
- Mallick, S., Standish, J.-J., & Bizimis, M. (2015). Constraints on the mantle mineralogy of an ultra-slow ridge: Hafnium isotopes in Abyssal peridotites and basalts from the 9–25°E Southwest Indian Ridge. *Earth and Planetary Science Letters*, 410, 42–53.
- Martin, C. E. (1991). Os isotopic characteristics of mantle derived rocks. *Geochimica et Cosmochimica Acta*, 55, 1421–1434.
- Marty, B., Zashu, S., & Ozima, M. (1983). Two noble gas components in a Mid-Atlantic Ridge basalt. *Nature*, 302(5905), 238.
- Marty, B., & Jambon, A. (1987). C3He in volatile fluxes from the solid Earth: Implications for carbon geodynamics. *Earth and Planetary Science Letters*, 83, 16–26.
- Marty, B., & Ozima, M. (1986). Noble gas distribution in oceanic basalt glasses. *Geochimica et Cosmochimica Acta*, 50, 1093–1097.
- Marty, B., & Tolstikhin, I. N. (1998). CO<sub>2</sub> fluxes from mid-ocean ridges, arcs and plumes. *Chemical Geology*, 145, 233–248.
- Marty, B., & Zimmermann, L. (1999). Volatiles (He, C, N, Ar) in Mid-Ocean Ridge basalts: Assessment of shallow-level fractionation and characterization of source composition. *Geochimica et Cosmochimica Acta*, 63, 3619–3634.
- Mathez, E. A. (1976). Sulfur solubility and magmatic sulfides in submarine basalt glass. *Journal of Geophysical Research*, 81, 4269–4276. <https://doi.org/10.1029/JB081i023p04269>
- Mattey, D. P., Carr, R. H., Wright, I. P., & Pillinger, C. T. (1984). Carbon isotopes in submarine basalts. *Earth and Planetary Science Letters*, 70, 196–206.
- Melson, W. G., & O'Hearn, T. (2003). Smithsonian volcanic glass file.
- Mertz, D. F., Devey, C. W., Todt, W., Stoffers, P., & Hofmann, A. W. (1991). Sr-Nd-Pb isotope evidence against plume-asthenosphere mixing north of Iceland. *Earth and Planetary Science Letters*, 107, 243–255.
- Mertz, D. F., & Haase, K. M. (1997). The radiogenic isotope composition of the high-latitude North Atlantic mantle. *Geology*, 25, 411–414.
- Meyer, P. S., & Bryan, W. B. (1996). Petrology of basaltic glasses from the Tag Segment: Implications for a deep hydrothermal heat source. *Geophysical Research Letters*, 23, 3435–3438. <https://doi.org/10.1029/96GL01033>
- Meyzen, C. M., Blichert-Toft, J., Ludden, J. N., Humler, E., Mevel, C., & Albarede, F. (2007). Isotopic portrayal of the Earth's upper mantle flow field. *Nature*, 447, 1069–1074.
- Meyzen, C. M., Ludden, J. N., Humler, E., Luais, B., Toplis, M. J., Mevel, C., & Storey, M. (2005). New insights into the origin and distribution of the Dupal isotope anomaly in the Indian Ocean mantle from MORB of the Southwest Indian Ridge. *Geochemistry, Geophysics, Geosystems*, 6, Q11K11. <https://doi.org/10.1029/2005GC000979>
- Michael, P. J. (1988). The concentration, behaviour and storage of H<sub>2</sub>O in the suboceanic upper mantle: Implications for mantle metasomatism. *Geochimica et Cosmochimica Acta*, 52, 555–566.
- Michael, P. J. (1995). Regionally distinctive sources of depleted MORB: Evidence from trace elements and H<sub>2</sub>O. *Earth and Planetary Science Letters*, 131, 301–320.
- Michael, P. J., & Chase, R. L. (1987). The influence of primary magma composition, H<sub>2</sub>O and pressure on Mid-Ocean Ridge basalt differentiation. *Contributions to Mineralogy and Petrology*, 96, 245–263.
- Michael, P. J., Chase, R. L., & Allan, J. F. (1989). Petrology and geologic variations along the Southern Explorer Ridge, northeast Pacific Ocean. *Journal of Geophysical Research*, 94, 13,895–13,918. <https://doi.org/10.1029/JB094iB10p13895>
- Michael, P. J., & Cornell, W. C. (1998). Influence of spreading rate and magma supply on crystallization and assimilation beneath Mid-Ocean Ridges: Evidence from chlorine and major element chemistry of Mid-Ocean Ridge basalts. *Journal of Geophysical Research*, 103, 18,325–18,356. <https://doi.org/10.1029/98JB00791>
- Michael, P. J., Forsyth, D. W., Blackman, D. K., Fox, P. J., Hanan, B. B., Harding, A. J., et al. (1994). Mantle control of a dynamically evolving spreading center: Mid-Atlantic Ridge 31–34°S. *Earth and Planetary Science Letters*, 121, 451–468.
- Michael, P. J., & Graham, D. W. (2015). The behavior and concentration of CO<sub>2</sub> in the suboceanic mantle: Inferences from undegassed ocean ridge and ocean island basalts. *Lithos*, 236–237, 338–351. <https://doi.org/10.1016/j.lithos.2015.08.020>
- Michael, P. J., & Graham, D. W. (2015). The behavior and concentration of CO<sub>2</sub> in the suboceanic mantle: Inferences from ungasped ocean ridge and ocean island basalts. *Lithos*, 236–237, 351. <https://doi.org/10.1016/j.lithos.2015.08.020>
- Michael, P. J., Langmuir, C. H., Dick, H. J. B., Snow, J. E., Goldstein, S. L., Graham, D. W., et al. (2003). Magmatic and amagmatic seafloor generation at the ultraslow-spreading Gakkel Ridge, Arctic Ocean. *Nature*, 423, 956–961.
- Michael, P. J., & Schilling, J.-G. (1989). Chlorine in mid-ocean ridge magmas: Evidence for assimilation of seawater-influenced components. *Geochimica et Cosmochimica Acta*, 53, 3131–3143. [https://doi.org/10.1016/0016-7037\(89\)90094-X](https://doi.org/10.1016/0016-7037(89)90094-X)
- Michael, P. J., & Schilling, J.-G. (1989). chlorine in Mid-Ocean Ridge magmas: evidence for assimilation of seawater-influenced components. *Geochimica et Cosmochimica Acta*, 53, 3131–3143.
- Michard, A., Montigny, R., & Schlich, R. (1986). Geochemistry of the mantle beneath the Rodriguez Triple Junction and the South-East Indian Ridge. *Earth and Planetary Science Letters*, 78, 104–111.
- Miller, D. J., & Kelley, J. (2004). Low-temperature alteration of basalt over time: A synthesis of results from Ocean Drilling Program Leg 187. In R. B. Pedersen, D. M. Christie, & D. J. Miller (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 187, pp. 1–29). College Station, Texas: Ocean Drilling Program. Retrieved from [https://www-odp.tamu.edu/publications/187\\_SR/206/206.htm](https://www-odp.tamu.edu/publications/187_SR/206/206.htm)
- Miyashiro, A., Shido, F., & Ewing, M. (1969). Diversity and origin of Abyssal tholeiite from the Mid-Atlantic Ridge near 24° and 30°N latitude. *Contributions to Mineralogy and Petrology*, 23, 38–52.
- Moeller, H. (2002). Magma genesis and mantle sources at the Mid-Atlantic Ridge east of Ascension Island, (Thesis Or Dissertation, Doctoral). Christian-Albrechts Universitaet; Monographic.
- Moore, A., Coogan, L. A., Costa, F., & Perfit, M. R. (2014). Primitive melt replenishment and crystal-mush disaggregation in the weeks preceding the 2005–2006 eruption at 9°50'N, EPR. *Earth and Planetary Science Letters*, 403, 15–26.
- Moore, J. G., Normark, W. R., Hess, G. R., & Meyer, C. E. (1977). Petrology of basalt from the East Pacific Rise near 21°N latitude. *Journal of Research of the U. S. Geological Survey*, 5, 753–759.
- Moore, J. G., & Schilling, J.-G. (1973). Vesicles, water and sulfur in Reykjanes Ridge basalts. *Contributions to Mineralogy and Petrology*, 41, 105–118.
- Moreira, M. (1995). Thomas Staudacher, Philippe Sarda, Jean-Guy Schilling, and Claude Jean Allègre. "A primitive plume neon component in MORB: The Shona ridge-anomaly, South Atlantic (51–52°S)". *Earth and Planetary Science Letters*, 133(3–4), 367–377.
- Moreira, M., Dosso, L., & Ondreas, H. (2008). Helium isotopes on the Pacific-Antarctic ridge (52.5°–41.5°S). *Geophysical Research Letters*, 35, L10306. <https://doi.org/10.1029/2008GL033286>

- Moreira, M., Escartin, J., Gayer, E., Hamelin, C., Bezos, A., Guillon, F., & Cannat, M. (2011). Rare gas systematics on Lucky Strike basalts (37°N, North Atlantic): Evidence for efficient homogenization in a long-lived magma chamber system? *Geophysical Research Letters*, *38*, L08304. <https://doi.org/10.1029/2011GL046794>
- Moreira, M., Kunz, J., & Allegre, C. J. (1998). Rare gas systematics in popping rock: Isotopic and elemental compositions in the upper mantle. *Science*, *279*, 1178–1181.
- Moreira, M., Valbracht, P. J., Staudacher, T., & Allegre, C. J. (1996). Rare gas systematics in Red Sea ridge basalts. *Geophysical Research Letters*, *23*, 2453–2456. <https://doi.org/10.1029/96GL02301>
- Mougel, B., Agraniér, A., Hemond, C., & Gente, P. (2014). A highly unradiogenic lead isotopic signature revealed by volcanic rocks from the East Pacific Rise. *Nature Communications*, *5*, 1–7.
- Mougel, B., Moreira, M., & Agraniér, A. (2015). A "high 4He/3He" mantle material detected under the East Pacific Rise (15°4'N). *Geophysical Research Letters*, *45*, 1375–1383. <https://doi.org/10.1002/2014GL062921>
- Muenow, D. W., Liu, N. W. K., Garcia, M. O., & Saunders, A. D. (1980). Volatiles in submarine volcanic rocks from the spreading axis of the East Scotia Sea back-arc basin. *Earth and Planetary Science Letters*, *47*, 149–153.
- Mühe, R. K., Bohrmann, H., Garbe-Schönberg, D., & Kassens, H. (1997). E-MORB glasses from the Gakkel Ridge (Arctic Ocean) at 87°N: Evidence for the Earth's most northerly volcanic activity. *Earth and Planetary Science Letters*, *152*, 1–9.
- Mühe, R. K., Bohrmann, H., Hörmann, P. K., Thiede, J., & Stoffers, P. (1991). Spinifex basalts with komatiite-tholeiite trend from the Nansen-Gakkel-Ridge (Arctic Ocean). *Tectonophysics*, *190*, 95–108.
- Mühe, R. K., Devey, C. W., & Bohrmann, H. (1993). Isotope and trace element geochemistry of MORB from the Gakkel Ridge at 86°N. *Earth and Planetary Science Letters*, *120*, 103–109.
- Murton, B. J., Taylor, R. N., & Thirlwall, M. F. (2002). Plume-ridge interaction: A geochemical perspective from the Reykjanes Ridge. *Journal of Petroleum*, *43*, 1987–2012.
- Murton, B. J., Tindle, A. G., Milton, J. A., & Sauter, D. (2005). Heterogeneity in southern Central Indian Ridge MORB: Implications for ridge-hot spot interaction. *Geochemistry, Geophysics, Geosystems*, *6*, Q03E20. <https://doi.org/10.1029/2004GC000798>
- Natland, J. H. (1989). Partial melting of a lithologically heterogeneous mantle: Inferences from crystallization histories of magnesian abyssal tholeiites from the Siqueiros Fracture Zone. Magmatism In The Ocean Basins, Saunders-A-D, Norry-M-J, (Eds.). *Geological Society, London, Special Publications*, *42*(1), 41–70. <https://doi.org/10.1144/GSLSP.1989.042.01.05>
- Natland, J. H. (1991). Indian Ocean crust. In P. A. Floyd (Ed.), *Oceanic Basalts* (pp. 289–310). Glasgow, UK: Blackie & Sons.
- Natland, J. H., & Melson, W. G. (1980). Comparison of basalt glasses from the East Pacific Rise and Siqueiros Fracture Zone near 9°N. *Initial Reports of the Deep Sea Drilling Project*, *54*, 705–723.
- Nauret, F., Abouchami, W., Galer, S. J. G., Hofmann, A. W., Hémond, C., Chauvel, C., & Dymet, J. (2006). Correlated trace element-Pb isotope enrichments in Indian MORB along 18–20°S, Central Indian Ridge. *Earth and Planetary Science Letters*, *245*(1-2), 137–152. <https://doi.org/10.1016/j.epsl.2006.03.015>
- Nauret, F., Snow, J. E., Hellebrand, E., & Weis, D. (2011). Geochemical composition of K-rich lavas from the Lena Trough (Arctic Ocean). *Journal of Petroleum*, *52*, 1185–1206.
- Neumann, E. R., & Schilling, J.-G. (1984). Petrology of basalts from the Mohs-Knipovich Ridge: The Norwegian-Greenland Sea. *Contributions to Mineralogy and Petrology*, *85*, 209–223.
- Newman, S., Finkel, R. C., & Macdougall, J. D. (1983). 230Th-238U disequilibrium systematics in oceanic tholeiites from 21°N on the East Pacific Rise. *Earth and Planetary Science Letters*, *65*, 17–33.
- Newman, S., Stolper, E., & Stern, R. J. (2000). H<sub>2</sub>O and CO<sub>2</sub> in magmas from the Mariana arc and back arc system. *Geochemistry, Geophysics, Geosystems*, *1*(5), 1013. <https://doi.org/10.1029/1999GC000027>
- Newsom, H. E., White, W. M., Jochum, K. P., & Hofmann, A. W. (1986). Siderophile and chalcophile element abundances in oceanic basalts, Pb isotope evolution and growth of the Earth's core. *Earth and Planetary Science Letters*, *80*, 299–313.
- Niedermann, S., & Bach, W. (1998). Anomalously nucleogenic neon in North Chile Ridge basalt glasses suggesting a previously degassed mantle source. *Earth and Planetary Science Letters*, *160*, 447–462.
- Niedermann, S., Erzinger, J., & Bach, W. (1997). Noble gas evidence for a lower mantle component in MORBs from the southern East Pacific Rise: Decoupling of helium and neon isotope systematics. *Geochimica et Cosmochimica Acta*, *61*, 2697–2715.
- Nielsen, R. L., Crum, J., Bourgeois, R., Hascall, K., Forsythe, L., Fisk, M. R., & Christie, D. M. (1995). Melt inclusions in high-An plagioclase from the Gorda Ridge: An example of the local diversity of MORB parent magmas. *Contributions to Mineralogy and Petrology*, *122*, 34–50.
- Nielsen, R. L., Sours-Page, R. E., & Harpp, K. S. (2000). Role of a Cl-bearing flux in the origin of depleted ocean floor magmas. *Geochemistry, Geophysics, Geosystems*, *1*(5), 1007. <https://doi.org/10.1029/1999GC000017>
- Nielsen, S., Shimizu, N., Lee, C.-T., & Behn, M. (2014). Chalcophile behavior of thallium during MORB melting and implications for the sulfur content of the mantle. *Geochemistry, Geophysics, Geosystems*, *15*, 4905–4919. <https://doi.org/10.1002/2014GC005536>
- Nishio, Y., Ishii, T., Gamo, T., & Sano, Y. (1999). Volatile element isotopic systematics of the Rodriguez Triple Junction Indian Ocean MORB: Implications for mantle heterogeneity. *Earth and Planetary Science Letters*, *170*, 241–253.
- Nishio, Y., Nakai, S. I., Ishii, T., & Sano, Y. (2007). Isotope systematics of Li, Sr, Nd, and volatiles in Indian Ocean MORBs of the Rodriguez Triple Junction: Constraints on the origin of the Dupal anomaly. *Geochimica et Cosmochimica Acta*, *71*, 745–759.
- Nishio, Y., Sasaki, S., Gamo, T., Hiyagon, H., & Sano, Y. (1998). Carbon and helium isotope systematics of north Fiji basin glasses: Carbon geochemical cycle in the subduction zone. *Earth and Planetary Science Letters*, *154*, 127–138.
- Niu, Y., & Batiza, R. (1994). Magmatic processes at a slow spreading ridge segment: 26°S Mid-Atlantic Ridge. *Journal of Geophysical Research*, *99*, 19,719–19,740. <https://doi.org/10.1029/94JB01663>
- Niu, Y., & Batiza, R. (1997). Trace element evidence from seamounts for recycled oceanic crust in the Eastern Pacific mantle. *Earth and Planetary Science Letters*, *148*, 471–483.
- Niu, Y., Bideau, R. D., Hekinian, R., & Batiza, R. (2001). Mantle compositional control on the extent of mantle melting, crust production, gravity anomaly, ridge morphology, and ridge segmentation: A case study at the Mid-Atlantic Ridge 3–35°N. *Earth and Planetary Science Letters*, *186*, 383–399.
- Niu, Y., Collerson, K. D., Batiza, R., Wendt, J. I., & Regelous, M. (1999). Origin of enriched-type Mid-Ocean Ridge basalt at ridges far from mantle plumes: The East Pacific Rise at 11°20'N. *Journal of Geophysical Research*, *104*, 7067–7087. <https://doi.org/10.1029/1998JB900037>
- Niu, Y., Gilmore, T., Mackie, S., Greig, A., & Bach, W. (2002). Mineral chemistry, whole-rock compositions, and petrogenesis of Leg 176 gabbros: Data and discussion. In J. H. Natland, H. J. B. Dick, D. J. Miller, & R. P. von Herzen (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 176, pp. 1–60). College Station, Texas: Ocean Drilling Program. Retrieved from [http://www-odp.tamu.edu/publications/176\\_SR/chap\\_08/chap\\_08.htm](http://www-odp.tamu.edu/publications/176_SR/chap_08/chap_08.htm)

- Niu, Y., Regelous, M., Wendt, J. I., Batiza, R., & O'Hara, M. J. (2002). Geochemistry of near-EPR seamounts: Importance of source vs. process and the origin of enriched mantle component. *Earth and Planetary Science Letters*, *199*, 327–345.
- Niu, Y., Waggoner, D. G., Sinton, J. M., & Mahoney, J. J. (1996). Mantle source heterogeneity and melting processes beneath seafloor spreading centers: The East Pacific Rise, 18°–19°S. *Journal of Geophysical Research*, *101*, 27,711–27,733. <https://doi.org/10.1029/96JB01923>
- Nowell, G. M., Kempton, P. D., Noble, S. R., Fitton, J. G., Saunders, A. D., Mahoney, J. J., & Taylor, R. N. (1998). High precision Hf isotope measurements of MORB and OIB by thermal ionisation mass spectrometry: insights into the depleted mantle. *Chemical Geology*, *149*(3–4), 211–233.
- O'Nions, R. K., Hamilton, P. J., & Evensen, N. M. (1977). Variations in  $^{143}\text{Nd}/^{144}\text{Nd}$  and  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios in oceanic basalts. *Earth and Planetary Science Letters*, *34*, 13–22.
- O'Nions, R. K., & Pankhurst, R. J. (1974). Petrogenetic significance of isotope and trace element variations in volcanic rocks from the mid-Atlantic. *Journal of Petrology*, *15*, 603–634.
- Ozima, M., & Zashu, S. (1983). Noble gases in submarine pillow volcanic glasses. *Earth and Planetary Science Letters*, *62*, 24–40.
- Pan, Y., & Batiza, R. (2003). Magmatic processes under Mid-Ocean Ridges: A detailed, mineralogic study of lavas from East Pacific Rise 9°30'N, 10°30'N, and 11°20'N. *Geochemistry, Geophysics, Geosystems*, *4*(11), 8623. <https://doi.org/10.1029/2002GC000309>
- Pan, Y., Batiza, R., & Dixon, J. E. (1998). Major element chemistry of volcanic glasses from the Easter Seamount Chain: Constraints on melting conditions in the plume channel. *Journal of Geophysical Research*, *103*, 5287–5304. <https://doi.org/10.1029/97JB03617>
- Pandey, S. K., Shrivastava, J. P., & Roonwal, G. S. (2009). Petrography and mineral chemistry of neovolcanics occurring between Pacific and Nazca Plate boundaries. *Journal of the Geological Society of India*, *74*, 559–572.
- Paquet, M., Cannat, M., Brunelli, D., Hamelin, C., & Humler, E. (2016). Effect of melt/lithosphere interactions on MORB chemistry at the easternmost Southwest Indian Ridge (61° to 67°E). *Geochemistry, Geophysics, Geosystems*, *17*, 4605–4640. <https://doi.org/10.1002/2016GC006385>
- Park, K. H., & Staudigel, H. (1990). Radiogenic isotope ratios and initial seafloor alteration in submarine Serocki Volcano basalts. In R. Detrick, J. Honnorez, W. B. Bryan, & T. Juteau (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 106–109, pp. 117–121). College Station, Texas: Ocean Drilling Program.
- Patchett, P. J. (1983). Hafnium isotope results from Mid-Ocean Ridges and Kerguelen. *Lithos*, *16*, 47–51.
- Patchett, P. J., & Tatsumoto, M. (1980). Hafnium isotope variations in oceanic basalts. *Geophysical Research Letters*, *7*, 1077–1080. <https://doi.org/10.1029/GL0071012p01077>
- Paulick, H., Muenker, C., & Schuth, S. (2010). The influence of small-scale mantle heterogeneities on Mid-Ocean Ridge volcanism: Evidence from the southern Mid-Atlantic Ridge (7°30'S to 11°30'S) and Ascension Island. *Earth and Planetary Science Letters*, *296*, 299–310.
- Pearce, J. A., Stern, R. J., Bloomer, S. H., & Fryer, P. (2005). Geochemical mapping of the Mariana arc-basin system: Implications for the nature and distribution of subduction components. *Geochemistry, Geophysics, Geosystems*, *6*, Q07006. <https://doi.org/10.1029/2004GC000895>
- Peate, D. W., Kokfelt, T. F., Hawkesworth, C. J., van Calsteren, P. W., Hergt, J. M., & Pearce, J. A. (2001). U-series isotope data on Lau Basin glasses: The role of subduction-related fluids during melt generation in back-arc basins. *Journal of Petrology*, *42*, 1449–1470.
- Pedersen, R. B., Christie, D. M., & Pyle, D. G. (2004). Regional and local mantle heterogeneities associated with the Australian Antarctic Discordance: Sr and Nd isotopic results from ODP Leg 187. In R. B. Pedersen, D. M. Christie, & D. J. Miller (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 187, pp. 1–23). College Station, Texas: Ocean Drilling Program.
- Pedersen, R. B., Malpas, J., & Falloon, T. J. (1996). Petrology and geochemistry of gabbroic and related rocks from Site 894, Hess Deep. In C. Mével, K. M. Gillis, & J. F. Allen (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 147, pp. 3–19). College Station, Texas: Ocean Drilling Program.
- Perfit, M. R. (2000). Submitted data set: Major element compositions of basalt glasses from the East Pacific Rise, 9°N. Earth Chem Library.
- Perfit, M. R., & Fornari, D. J. (1983). Geochemical studies of abyssal lavas recovered by DSRV Alvin from eastern Galapagos rift, Inca transform, and Ecuador rift. 2. Phase chemistry and crystallization history. *Journal of Geophysical Research*, *88*, 10,530–10,550. <https://doi.org/10.1029/JB088iB12p10530>
- Perfit, M. R., Fornari, D. J., Malahoff, A., & Embley, R. W. (1983). Geochemical studies of abyssal lavas recovered by dsrv alvin from eastern Galapagos rift, inca transform, and ecuador rift 3. trace element abundances and petrogenesis. *Journal of Geophysical Research*, *88*, 10,551–10,572. <https://doi.org/10.1029/JB088iB12p10551>
- Perfit, M. R., Fornari, D. J., Ridley, W. I., Kirk, P. D., Casey, J. F., Kastens, K. A., et al. (1996). Recent volcanism in the Siqueiros transform fault: Picritic basalts and implications for MORB magma genesis. *Earth and Planetary Science Letters*, *141*, 91–108.
- Perfit, M. R., Fornari, D. J., Smith, M. C., Bender, J. F., Langmuir, C. H., & Haymon, R. (1994). Small-scale spatial and temporal variations in Mid-Ocean Ridge crest magmatic processes. *Geology*, *22*, 375–379.
- Perfit, M. R., Langmuir, C. H., Baekisapa, M., Chappell, B. W., Johnson, R. W., Staudigel, H., & Taylor, S. R. (1987). Geochemistry and petrology of volcanic rocks from the Woodlark basin: Addressing questions of ridge subduction. In B. Taylor & N. F. Exon (Eds.), *Marine Geology, Geophysics, and Geochemistry of the Woodlark Basin - Solomon Islands, Earth Science Series* (Vol. 7, pp. 113–154). Houston, TX: Circum-Pacific Council for Energy and Mineral Resources.
- Perfit, M. R., Ridley, W. I., & Jonasson, I. R. (1998). Geologic, petrologic and geochemical relationships between magmatism and massive sulfide mineralization along the eastern Galapagos Spreading Center. In T. Barry & M. Hannington (Eds.), *Volcanic-Associated Massive Sulfide Deposits: Processes and Examples in Modern and Ancient Settings, Reviews in Economic Geology* (Vol. 8, pp. 75–100).
- Perfit, M. R., Smith, M. C., Walters, R., Ridley, W. I., & Fornari, D. J. (2013). Compilation of East Pacific Rise 9°–10°N basalt geochemistry, Submitted Data Set.
- Petrini, R., Joron, J.-L., Ottonello, G., Bonatti, E., & Seyler, M. (1988). Basaltic dykes from Zabargad Island, RSea: Petrology and geochemistry. *Tectonophysics*, *150*, 229–248.
- Pineau, F., & Javoy, M. (1983). Carbon isotopes and concentrations in Mid-Ocean Ridge basalts. *Earth and Planetary Science Letters*, *62*, 239–257.
- Pineau, F., & Javoy, M. (1994). Strong degassing at ridge crests: The behaviour of dissolved carbon and water in basalt glasses at 14°N, Mid-Atlantic Ridge. *Earth and Planetary Science Letters*, *123*, 179–198.
- Poldervaart, A., & Green, J. (1965). Chemical analyses of submarine basalts. *American Mineralogist*, *50*, 1723–1728.
- Pollock, M. A., Klein, E. M., Karson, J. A., & Coleman, D. S. (2009). Compositions of dikes and lavas from the Pito Deep Rift: Implications for crustal accretion at superfast spreading centers. *Journal of Geophysical Research*, *114*, B03207. <https://doi.org/10.1029/2007JB005436>
- Poreda, R., & Radicati di Brozolo, F. (1984). Neon isotope variations in Mid-Atlantic Ridge basalts. *Earth and Planetary Science Letters*, *69*(2), 277–289.
- Poreda, R. J. (1985). Helium-3 and deuterium in back-arc basalts: Lau Basin and the Mariana Trough. *Earth and Planetary Science Letters*, *73*, 244–254.



- Poreda, R. J., Schilling, J.-G., & Craig, H. (1986). Helium and hydrogen isotopes in ocean ridge basalts north and south of Iceland. *Earth and Planetary Science Letters*, *78*, 1–17.
- Price, R. C., Johnson, L. E., & Crawford, A. J. (1990). Basalts from the north Fiji basin: The generation of back arc basins magmas by mixing of depleted and enriched mantle sources. *Contributions to Mineralogy and Petrology*, *105*, 106–121.
- Price, R. C., Kennedy, A. K., Riggs-Sneeringer, M., & Frey, F. A. (1986). Geochemistry of basalts from the Indian Ocean triple junction: Implications for the generation and evolution of Indian Ocean ridge basalts. *Earth and Planetary Science Letters*, *78*, 379–396.
- Prinzhofer, A., Lewin, E., & Allègre, C. J. (1989). Stochastic melting of the marble cake mantle: evidence from local study of the East Pacific Rise at 12°50'N. *Earth and Planetary Science Letters*, *92*, 189–206.
- Prytulak, J., Avanzinelli, R., Koetsier, G., Kreissig, K., Beier, C., & Elliott, T. (2014). Melting versus contamination effects on 238U-230Th-226Ra and 235U-233Pa disequilibria in lavas from São Miguel, Azores. *Chemical Geology*, *381*, 94–109.
- Puchelt, H., & Emmermann, R. (1983). Petrogenetic implications of tholeiitic basalt glasses from the East Pacific Rise and the Galapagos Spreading Center. *Chemical Geology*, *38*, 39–56.
- Puchelt, H., Prichard, H., Berner, Z., & Maynard, J. (1996). Sulfide mineralogy, sulfur content, and sulfur isotope composition of mafic and ultramafic rocks from Leg 147. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 147, pp. 91–101). College Station, Texas: Ocean Drilling Program.
- Pyle, D. G. (1994). Geochemistry of Mid-Ocean Ridge basalt within and surrounding the Australian-Antarctic discordance. Thesis Or Dissertation, Doctoral, Osu; Monographic. *Geochemistry Geology*, *188*, 55–08.
- Pyle, D. G., Christie, D. M., & Mahoney, J. J. (1992). Resolving an isotopic boundary within the Australian-Antarctic discordance. *Earth and Planetary Science Letters*, *112*, 161–178.
- Pyle, D. G., Christie, D. M., Mahoney, J. J., & Duncan, R. A. (1995). Geochemistry and geochronology of ancient southeast Pacific seafloor. *Journal of Geophysical Research*, *100*, 22,261–22,282. <https://doi.org/10.1029/95JB01424>
- Raquin, A., & Moreira, M. (2009). Atmospheric 38Ar/36Ar in the mantle: Implications for the nature of the terrestrial parent bodies. *Earth and Planetary Science Letters*, *287*, 551–558. <https://doi.org/10.1016/j.epsl.2009.09.003>
- Ray, D., Misra, S., Widdowson, M., & Langmuir, C. H. (2014). A common parentage for Deccan Continental Flood Basalt and Central Indian Ocean Ridge Basalt? A geochemical and isotopic approach. *Journal of Asian Earth Sciences*, *84*, 188–200.
- Regelous, M., Niu, Y., Wendt, J. I., Batiza, R., Greig, A., & Collerson, K. D. (1999). Variations in the geochemistry of magmatism on the East Pacific Rise at 10°30'N since 800 ka. *Earth and Planetary Science Letters*, *168*, 45–63.
- Rehkaemper, M., Halliday, A. N., Fitton, J. G., Lee, D. C., Wieneke, M., & Arndt, N. (1999). Ir, Ru, Pt, and Pd in basalts and komatiites: New constraints for the geochemical behavior of the platinum-group elements in the mantle. *Geochimica et Cosmochimica Acta*, *63*, 3915–3934.
- Rehkaemper, M., & Hofmann, A. W. (1997). Recycled ocean crust and sediment in Indian Ocean MORB. *Earth and Planetary Science Letters*, *147*, 93–106.
- Reynolds, J. R. (1995). Segment-scale systematics of Mid-Ocean Ridge magmatism and geochemistry, (Thesis Or Dissertation, Doctoral). Columbia University; Monographic.
- Reynolds, J. R., & Langmuir, C. H. (1997). Petrological systematics of the Mid-Atlantic Ridge south of Kane, implications for ocean crust formations. *Journal of Geophysical Research*, *102*, 14,915–14,946. <https://doi.org/10.1029/97JB00391>
- Rhodes, J. M., Morgan, C., & Lias, R. A. (1990). Geochemistry of axial seamount lavas: Magmatic relationship between the Cobb Hotspot and the Juan de Fuca Ridge. *Journal of Geophysical Research*, *95*, 12,713–12,733. <https://doi.org/10.1029/JB095iB08p12713>
- Ridley, W. I., Perfit, M. R., & Adams, M. L. (1978). Petrology of basalts from Deep Sea Drilling Project, Leg 38. *Initial Reports of the Deep Sea Drilling Project*, *38*, 731–739.
- Ridley, W. I., Perfit, M. R., Jonasson, I. R., & Smith, M. (1994). Hydrothermal alteration in oceanic ridge volcanoes: A detailed study at the Galapagos Fossil Hydrothermal Field. *Geochimica et Cosmochimica Acta*, *58*, 2477–2494.
- Robinson, C.-J., Bickle, M.-J., Minshull, T. A., White, R. S., & Nichols, A. (2001). Low degree melting under the Southwest Indian Ridge: The roles of mantle temperature, conductive cooling and wet melting. *Earth and Planetary Science Letters*, *188*, 383–398.
- Robinson, C.-J., White, R. S., Bickle, M.-J., & Minshull, T. A. (1996). Restricted melting under the very slow-spreading Southwest Indian Ridge. *Geological Society, London, Special Publications*, *118*, 131–141.
- Rodehorst, U., Schmincke, H. U., & Sumita, M. (1998). Geochemistry and petrology of Pleistocene ash layers erupted at Las Canadas edifice (Tenerife). In *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 157, pp. 315–328). College Station, Texas: Ocean Drilling Program. <https://doi.org/10.1029/1998RG000054>
- Rogers, N. W. (1993). The isotope and trace element geochemistry of basalts from volcanic islands of the southern Red Sea. In H. M. Pichard, T. Alabaster, N. B. W. Harris, & C. R. Neary (Eds.), *Magmatic Processes and Plate Tectonics, Special Publications* (pp. 76, 455–467). London: Geological Society.
- Rosendahl, B. R., Furbish, W. J., & Meadows, G. (1980). Picritic basalts from the Siqueiros Transform Fault. *Initial Reports of the Deep Sea Drilling Project*, *54*, 71–78.
- Rotella, M., Sinton, J. M., Mahoney, J. J., & Chazey Iii, W. (2009). Geochemical evidence for low magma supply and inactive propagation at the Galapagos 93.25°W overlapping spreading center. *Geochemistry, Geophysics, Geosystems*, *10*, Q09005. <https://doi.org/10.1029/2009GC002445>
- Rowe, E. C., & Schilling, J. G. (1979a). Fluorine in Iceland and Reykjanes Ridge basalts. *Nature*, *279*, 33–37. <https://doi.org/10.1038/279033a0>
- Roy-Barman, M., & Allègre, C. J. (1994). 187Os/186Os ratios of mid-ocean ridge basalts and abyssal peridotites. *Geochimica et Cosmochimica Acta*, *58*, 5043–5054.
- Roy-Barman, M., Wasserburg, G. J., Papanastassiou, D. A., & Chaussidon, M. (1998). Osmium isotopic compositions and re-os concentrations in sulfide globules from basaltic glasses. *Earth and Planetary Science Letters*, *154*, 331–347.
- Rubin, K., & H., & Macdougall, J. D. (1990). Dating of neovolcanic MORB using (226Ra/230Th) disequilibrium. *Earth and Planetary Science Letters*, *101*, 313–322.
- Rubin, K. H., & Macdougall, J. D. (1988). 226Ra excesses in Mid-Ocean Ridge basalts and mantle melting. *Nature*, *355*, 158–161.
- Ryan, J. G., & Langmuir, C. H. (1987). The systematics of lithium abundances in young volcanic rocks. *Geochimica et Cosmochimica Acta*, *51*, 1727–1741.
- Ryan, J. G., & Langmuir, C. H. (1993). The systematics of boron abundances in young volcanic rocks. *Geochimica et Cosmochimica Acta*, *57*, 1489–1498.
- Saal, A. E., Hauri, E. H., Langmuir, C. H., & Perfit, M. R. (2002). Vapour undersaturation in primitive mid-ocean-ridge basalt and the volatile content of Earth's upper mantle. *Nature*, *419*, 451–455.

- Sakai, H., Des Marais, D. J., Ueda, A., & Moore, J. G. (1984). Concentrations and isotope ratios of carbon, nitrogen and sulfur in ocean-floor basalts. *Geochimica et Cosmochimica Acta*, *48*, 2433–2441.
- Salters, V. J. M. (1996). The generation of mid-ocean ridge basalts from the Hf and Nd isotope perspective. *Earth and Planetary Science Letters*, *141*, 109–123.
- Salters, V. J. M., & Hart, S. R. (1991). The mantle sources of ocean ridges, islands and arcs: The Hf-isotope connection. *Earth and Planetary Science Letters*, *104*, 364–380.
- Salters, V. J. M., Mallick, S., Hart, S. R., Langmuir, C. E., & Stracke, A. (2011). Domains of depleted mantle: New evidence from hafnium and neodymium isotopes. *Geochemistry, Geophysics, Geosystems*, *12*, Q08001. <https://doi.org/10.1029/2011GC003617>
- Salters, V. J. M., & White, W. M. (1998). Hf isotope constraints on mantle evolution. *Chemical Geology*, *145*, 447–460.
- Sano, Y., Nishio, Y., Gamo, T., Jambon, A., & Marty, B. (1998). Noble gas and carbon isotopes in Mariana Trough basalt glasses. *Applied Geochemistry*, *13*, 441–449.
- Sarda, P., Moreira, M., Staudacher, T., Schilling, J.-G., & AlLegré, C. J. (2000). Rare gas systematics on the southernmost Mid-Atlantic Ridge: Constraints on the lower mantle and the Dupal Source. *Journal of Geophysical Research*, *105*, 5973–5996. <https://doi.org/10.1029/1999JB900282>
- Sarda, P., Staudacher, T., & AlLegré, C. J. (1985). 40Ar/36Ar in MORB glasses: Constraints on atmosphere and mantle evolution. *Earth and Planetary Science Letters*, *72*, 357–375.
- Sarda, P., Staudacher, T., & AlLegré, C. J. (1988). Neon isotopes in submarine basalts. *Earth and Planetary Science Letters*, *91*, 73–88.
- Saunders, A. D. (1983). Geochemistry of basalts recovered from the Gulf of California during Leg 65 of the Deep Sea Drilling Project. *Initial Reports of the Deep Sea Drilling Project*, *65*, 591–621.
- Saunders, A. D., Fornari, D. J., Joron, J.-L., Tarney, J., & Treuil, M. (1982). Geochemistry of basic igneous rocks, Gulf of California, Deep Sea Drilling Project Leg 64. *Initial Reports of the Deep Sea Drilling Project*, *64*, 595–642.
- Scheidegger, K. F., & Corliss, J. B. (1981). Petrogenesis and secondary alteration of upper layer 2 basalts of the Nazca Plate. *Geological Society of America Memoirs*, *154*, 77–107.
- Schiano, P., Birck, J. L., & AlLegré, C. J. (1997). Osmium-strontium-neodymium-lead isotopic covariations in Mid-Ocean Ridge basalt glasses and the heterogeneity of the upper mantle. *Earth and Planetary Science Letters*, *150*, 363–379.
- Schiffman, P., Zierenberg, R. A., Chadwick, W., Clague, D. A., & Lowenstern, J. (2010). Contamination of basaltic lava by seawater: Evidence found in a lava pillar from Axial Seamount, Juan de Fuca Ridge. *Geochemistry, Geophysics, Geosystems*, *11*, Q04004. <https://doi.org/10.1029/2009GC003009>
- Schilling, J.-G. (1971). Sea-floor evolution: Rare-earth evidence. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, *A268*, 663–706.
- Schilling, J.-G. (1973). Iceland mantle plume: Geochemical study of Reykjanes Ridge. *Nature*, *242*, 565–571.
- Schilling, J.-G. (1975). Rare-Earth variations across normal segments of the Reykjanes Ridge, 60°–53°N, Mid-Atlantic Ridge, 29°S, and the East Pacific Rise, 2°–19°S, and evidence on the composition of the underlying low-velocity layer. *Journal of Geophysical Research*, *80*(11), 1459–1473. <https://doi.org/10.1029/JB080i011p01459>
- Schilling, J.-G., Anderson, R. N., & Vogt, P. R. (1976). Rare Earth, Fe and Ti variations along the Galapagos spreading center and their relationship to the Galapagos mantle plume. *Nature*, *261*, 108–113.
- Schilling, J.-G., Bergeron, M. B., & Evans, R. (1980). Halogens in the mantle beneath the North Atlantic. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, *297*, 147–178.
- Schilling, J.-G., Fontignie, D., Blichert-Toft, J., Kingsley, R. H., & Tomza, U. (2003). Pb-Hf-Nd-Sr isotope variations along the Galapagos spreading center (101°–83°W): Constraints on the dispersal of the Galapagos mantle plume. *Geochemistry, Geophysics, Geosystems*, *4*(10), 8512. <https://doi.org/10.1029/2002GC000495>
- Schilling, J.-G., Hanan, B. B., Mccully, B. L., Kingsley, R. H., & Fontignie, D. (1994). Influence of the Sierra Leone mantle plume on the equatorial Mid-Atlantic Ridge. *Journal of Geophysical Research*, *99*, 12,005–12,028. <https://doi.org/10.1029/94JB00337>
- Schilling, J.-G., Kingsley, R. H., & Devine, J. D. (1982). Galapagos hot spot spreading center system 1. Spatial petrological and geochemical variations (83°W–101°W). *Journal of Geophysical Research*, *87*, 5593–5610. <https://doi.org/10.1029/JB087iB07p05593>
- Schilling, J.-G., Kingsley, R. H., Fontignie, D., Poreda, R. J., & Xue, S. (1999). Dispersion of the Jan Mayen and Iceland mantle plumes in the Arctic: A He-Pb-Nd-Sr isotope tracer study of basalts from the Kolbeinsey, Mohns, and Knipovich Ridges. *Journal of Geophysical Research*, *104*, 10,543–10,569. <https://doi.org/10.1029/1999JB900057>
- Schilling, J.-G., Kingsley, R. H., Hanan, B. B., & Mccully, B. L. (1992). Nd-Sr-Pb isotopic variations along the Gulf of Aden: Evidence for Afar mantle plume-continental lithosphere interaction. *Journal of Geophysical Research*, *97*, 10,927–10,966. <https://doi.org/10.1029/92JB00415>
- Schilling, J.-G., Ruppel, C., Davis, A. N., Mccully, B. L., Tighe, S. A., Kingsley, R. H., & Lin, J. (1995). Thermal structure of the mantle beneath the equatorial Mid-Atlantic Ridge: Inferences from the spatial variation of dredged basalt glass compositions. *Journal of Geophysical Research*, *100*, 10,057–10,076. <https://doi.org/10.1029/95JB00668>
- Schilling, J.-G., Sigurdsson, H., Davis, A. N., & Hey, R. N. (1985). Easter microplate evolution. *Nature*, *317*, 325–331.
- Schilling, J. G., Zajac, M., Evans, R., Johnston, T., White, W., Devine, J. D., & Kingsley, R. (1983). Petrologic and geochemical variations along the Mid-Atlantic Ridge from 29°N to 73°N. *American Journal of Science*, *283*, 510–586. <https://doi.org/10.2475/ajs.283.6.510>
- Schrader, E. L., Rosendahl, B. R., Furbish, W. J., & Meadows, G. (1980). Picritic basalts from the Siqueiros Fracture Zone. *Initial Reports of the Deep Sea Drilling Project*, *54*, 71–78.
- Scott, S. D., Chase, R. L., Hannington, M. D., Michael, P. J., Mcconachy, T. F., & Shea, G. T. F. (1990). Sulphide deposits, tectonics and petrogenesis of Southern Explorer Ridge, Northeast Pacific Ocean. In J. Malpas, E. M. Moores, A. Panayiotou, & C. Xenophontos (Eds.), *Ophiolites; Oceanic Crustal Analogues: Proceedings of The Symposium, Troodos 1987* (pp. 719–733). Nicosia, Cyprus: Geol. Surv. Dept. Min. Agric. and Nat. Res.
- Selo, M., & Storzer, D. (1979). Chronologie des evenements volcaniques de la zone FAMOUS. *Comptes Rendus de Academie des Sciences, Paris, Serie II*, *289*, 1125–1128.
- Shaw, A. M., Behn, M. D., Humphris, S. E., Sohn, R. A., & Gregg, P. M. (2010). Deep pooling of low degree melts and volatile fluxes at the 85°E segment of the Gakkel Ridge: Evidence from olivine-hosted melt inclusions and glasses. *Earth and Planetary Science Letters*, *289*, 311–322. <https://doi.org/10.1016/j.epsl.2009.11.018>
- Shaw, A. M., Hauri, E. H., Behn, M., Hilton, D. R., Macpherson, C. G., & Sinton, J. M. (2012). Long-term preservation of slab signatures in the mantle inferred from hydrogen isotopes. *Nature Geoscience*, *5*, 224–228.
- Shaw, A. M., Hilton, D. R., Macpherson, C. G., & Sinton, J. M. (2004). The CO<sub>2</sub>-He-Ar-H<sub>2</sub>O systematics of the Manus back-arc basin: Resolving source composition from degassing and contamination effects. *Geochimica et Cosmochimica Acta*, *68*, 1837–1856.

- Shibata, T., & Fox, P. J. (1975). Fractionation of abyssal tholeiites: Samples from the Oceanographer Fracture Zone (35°N, 35°W). *Earth and Planetary Science Letters*, 27, 62–72.
- Shibata, T., Thompson, G., & Frey, F. A. (1979). Tholeiitic and alkali basalts from the Mid-Atlantic Ridge at 43°N. *Contributions to Mineralogy and Petrology*, 70, 127–141.
- Shirey, S. B., Bender, J. F., & Langmuir, C. H. (1987). Three-component isotopic heterogeneity near the Oceanographer Transform, Mid-Atlantic Ridge: Evidence for recycling of continental lithosphere. *Nature*, 325, 217–223.
- Shorttle, O., Moussallam, Y., Hartley, M., MacLennan, J., Ivanov, A., Murton, B. J., & Pedersen, R. B. (2015). Fe-XANES analyses of Reykjanes Ridge basalts: Implications for oceanic crust's role in the solid Earth oxygen cycle. *Earth and Planetary Science Letters*, 427, 272–285.
- Sigurdsson, H. (1977). Spinels in Leg 37 basalts and peridotites: Phase chemistry and zoning. *Initial Reports of the Deep Sea Drilling Project*, 37, 883–891.
- Sigurdsson, H. (1981). First-order major element variation in basalt glasses from the Mid-Atlantic Ridge: 29°N to 73°N. *Journal of Geophysical Research*, 86, 9483–9502. <https://doi.org/10.1029/JB086iB10p09483>
- Silant'ev, S. A., Belyatsky, B., Beltenev, V. E., & Vikentyev, I. V. (2001). The distribution of isotope signatures in MAR peridotites between 12 and 36°N and two main kinds of mantle substratum below ridge axis. *InterRidge News*, 10, 27–33.
- Simons, K., Dixon, J. E., Schilling, J.-G., Kingsley, R. H., & Poreda, R. J. (2002). Volatiles in basaltic glasses from the Easter-Salas y Gomez Seamount Chain and Easter Microplate: Implications for geochemical cycling of volatile elements. *Geochemistry, Geophysics, Geosystems*, 3(7), 1039. <https://doi.org/10.1029/2001GC000173>
- Sims, K. W. W., Blichert-Toft, J., Fornari, D. J., Perfit, M. R., Goldstein, S. J., Johnson, P., et al. (2003). Aberrant youth: Chemical and isotopic constraints on the origin of off-axis lavas from the East Pacific Rise, 9°–10°N. *Geochemistry, Geophysics, Geosystems*, 4(10), 8621. <https://doi.org/10.1029/2002GC000443>
- Sims, K. W. W., De Paolo, D. J., Murrell, M. T., Baldrige, W. S., Goldstein, S. J., & Clague, D. A. (1995). Mechanisms of magma generation beneath Hawaii and Mid-Ocean Ridges: Uranium/thorium and samarium/neodymium isotopic evidence. *Science*, 267, 508–512.
- Sims, K. W. W., Goldstein, S. J., Blichert-Toft, J., Donnellan, N. C. B., Perfit, M. R., Kelemen, P., et al. (2002). Chemical and isotopic constraints on the generation and transport of magma beneath the East Pacific Rise. *Geochimica et Cosmochimica Acta*, 66, 3481–3504.
- Sinton, J. M. (1999). Submitted data set: Major and trace element compositions of basalts from the southern East Pacific Rise, Earthchem Library.
- Sinton, J. M., Ford, L. L., Chappell, B. W., & McCulloch, M. T. (2003). Magma genesis and mantle heterogeneity in the Manus back-arc basin, Papua New Guinea. *Journal of Petrology*, 44, 159–195.
- Sinton, J. M., Price, R. C., Johnson, K. T. M., Staudigel, H., & Zindler, A. (1993). Petrology and geochemistry of submarine lavas from the Lau and North Fiji back-arc basins. In L. W. Kroenke & J. V. Eads (Eds.), *Basin Formation, Ridge Crest Processes and Metallogenesis in the North Fiji Basin*, *Earth Science Series* (Vol. 15, pp. 119–135). Houston, TX: Circum-Pacific Council for Energy and Mineral Resources.
- Sinton, J. M., Smaglik, S. M., Mahoney, J. J., & Macdonald, K. C. (1991). Magmatic processes at superfast spreading Mid-Ocean Ridges: Glass compositional variations along the East Pacific Rise 13°–23°S. *Journal of Geophysical Research*, 96(B4), 6133–6155. <https://doi.org/10.1029/90JB02454>
- Smith, A. D. (1999). The Nd-Sr-Pb isotopic record in abyssal tholeiites from the Gulf of California region, western Mexico: No evidence for a gulf mouth plume. *International Geology Review*, 41, 921–931.
- Smith, M. C., Perfit, M. R., Fornari, D. J., Ridley, W. I., Edwards, M., Kurras, G. J., & von Damm, K. L. (2001). Magmatic processes and segmentation at a fast spreading Mid-Ocean Ridge: Detailed investigation of an axial discontinuity on the East Pacific Rise crest at 9°37'. *Geochemistry, Geophysics, Geosystems*, 2(10), 1040. <https://doi.org/10.1029/2000GC000134>
- Smith, M. C., Perfit, M. R., & Jonasson, I. R. (1994). Petrology and geochemistry of basalts from the southern Juan de Fuca Ridge: Controls on the spatial and temporal evolution of Mid-Ocean Ridge basalts. *Journal of Geophysical Research*, 99(B3), 4787–4812. <https://doi.org/10.1029/93JB02158>
- Smith, S. E., Casey, J. F., Bryan, W. B., Dmitriev, L. V., Silant'ev, S. A., & Magakyan, R. (1998a). Geochemistry of basalts from the Hayes Transform region of the Mid-Atlantic Ridge. *Journal of Geophysical Research*, 103(B3), 5305. <https://doi.org/10.1029/97JB03208>
- Snow, J. E., & Reisberg, L. R. (1995). Os isotope systematics of the MORB mantle: Results from altered abyssal peridotites. *Earth and Planetary Science Letters*, 133, 411–421.
- Soule, S. A., Nakata, D. S., Fornari, D. J., Fundis, A., Perfit, M. R., & Kurz, M. D. (2012). CO<sub>2</sub> variability in Mid-Ocean Ridge basalts from syn-emplacement degassing: Constraints on eruption dynamics. *Earth and Planetary Science Letters*, 327–328, 39–49.
- Stakes, D., Shervais, J. W., & Hopson, C. A. (1984). The volcanic-tectonic cycle of the FAMOUS and AMAR valleys, Mid-Atlantic Ridge (36°47'N): Evidence from basalt glass and phenocryst compositional variations for a steady-state magma chamber beneath the valley midsections. *Journal of Geophysical Research*, 89, 6995–7028. <https://doi.org/10.1029/JB089iB08p06995>
- Standish, J.-J., Dick, H. J. B., Michael, P. J., Melson, W. G., & O'Hearn, T. (2008). MORB generation beneath the ultraslow spreading Southwest Indian Ridge (9–25°E): major element chemistry and the importance of process versus source. *Geochemistry, Geophysics, Geosystems*, 9, Q05004. <https://doi.org/10.1029/2008GC001959>
- Standish, J.-J., & Sims, K. W. W. (2010). Young off-axis volcanism along the ultraslow-spreading Southwest Indian Ridge. *Nature Geoscience*, 3, 286–292.
- Staudacher, T., Sarda, P., Richardson, S. H., AlLègre, C. J., Sagna, I., & Dmitriev, L. V. (1989). Noble gases in basalt glasses from a Mid-Atlantic Ridge topographic high at 14°N: Geodynamic consequences. *Earth and Planetary Science Letters*, 96, 119–133.
- Stolper, E., & Newman, S. (1994). The role of water in the petrogenesis of Mariana Trough magmas. *Earth and Planetary Science Letters*, 121, 293–325.
- Straub, S. (2003). The evolution of the Izu Bonin-Mariana volcanic arcs (NW Pacific) in terms of major element chemistry. *Geochemistry, Geophysics, Geosystems*, 4(2), 1018. <https://doi.org/10.1029/2002GC000357>
- Stroncić, N., & Devey, C. W. (2011). Recycled gabbro signature in hotspot magmas unveiled by plume-ridge interactions. *Nature Geoscience*, 4, 393–397.
- Stroncić, N., & Niedermann, S. (2015). Atmospheric contamination of the primary Ne and Ar signal in Mid-Ocean Ridge basalts and its implications for ocean crust formation. *Geochimica et Cosmochimica Acta*, 172, 306–321.
- Sturm, M. E., Klein, E. M., Graham, D. W., & Karsten, J. L. (1999). Age constraints on crustal recycling to the mantle beneath the Southern Chile Ridge: He-Pb-Sr-Nd isotope systematics. *Journal of Geophysical Research*, 104, 5097–5114. <https://doi.org/10.1029/1998JB900107>
- Subbarao, K. V. (1972). The strontium isotopic composition of basalts from the East Pacific and Chile Rises and abyssal hills in the Eastern Pacific Ocean. *Contributions to Mineralogy and Petrology*, 37, 111.

- Subbarao, K. V., & Hedge, C. E. (1973). K, Rb, Sr and  $87\text{Sr}/86\text{Sr}$  in rocks from mid-Indian Ocean Ridge. *Earth and Planetary Science Letters*, *18*, 223.
- Subbarao, K. V., & Hekinian, R. (1978). Alkali-enriched rocks from the central Eastern Pacific Ocean. *Marine Geology*, *26*, 249–268.
- Subbarao, K. V., Hekinian, R., & Chandrasekharam, D. (1977). Large ion lithophile elements and Sr and Pb isotopic variations in volcanic rocks from the Indian Ocean. In J. R. Hertzler, H. M. Bolli, T. A. Davies, J. B. Saunders, & J. G. Sclater (Eds.), *Indian Ocean Geology and Biostratigraphy: Studies Following Deep-Sea Drilling Legs* (pp. 259–278). Washington, DC: American Geophysical Union.
- Subbarao, K. V., & Reddy, V. V. (1981). Geochemical studies on oceanic basalts from the Indian Ocean. *Tectonophysics*, *75*(1-2), 69–89. [https://doi.org/10.1016/0040-1951\(81\)90210-9](https://doi.org/10.1016/0040-1951(81)90210-9)
- Sun, S. S., & Crawford, A. J. (1980). Lead isotope studies of young volcanic rocks from mid-ocean ridges, ocean islands and island arcs. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, *297*, 409–445.
- Sun, S. S., Nesbitt, R. W., & Sharaskin, A. Y. (1979). Geochemical characteristics of mid-ocean ridge basalts. *Earth and Planetary Science Letters*, *44*, 119–138.
- Sun, S. S., Tatsumoto, M., & Schilling, J.-G. (1975). Mantle plume mixing along the Reykjanes Ridge axis: Lead isotopic evidence. *Science*, *190*, 143–147.
- Sun, W., Arculus, R. J., Kamenetsky, V. S., & Hirose, K. (2004). Release of gold-bearing fluids in convergent margin magmas prompted by magnetite crystallization. *Nature*, *431*, 975–978.
- Sun, W., Bennett, V. C., & Kamenetsky, V. S. (2004). The mechanism of re enrichment in arc magmas: Evidence from Lau Basin basaltic glasses and primitive melt inclusions. *Earth and Planetary Science Letters*, *222*, 101–114.
- Sun, W., Binns, R. A., Fan, A. C., Kamenetsky, V. S., Wysoczanski, R., Wei, G. J., et al. (2007). Chlorine in submarine volcanic glasses from the eastern Manus Basin. *Geochimica et Cosmochimica Acta*, *71*, 1542–1552.
- Sunkel, G. (1990). Origin of petrological and geochemical variations of Lau Basin lavas (SW Pacific). *Marine Mining*, *9*, 205–234.
- Tatsumoto, M. (1966). Genetic relations of oceanic basalts as indicated by lead isotopes. *Science*, *153*, 1094–1101.
- Tatsumoto, M. (1978). Isotopic composition of lead in oceanic basalt and its implication to mantle evolution. *Earth and Planetary Science Letters*, *38*, 63–87.
- Tatsumoto, M., Hedge, C. E., & Engel, A. E. J. (1965). Potassium, rubidium, strontium, thorium, uranium and the ratio strontium-87 to strontium-86 in oceanic tholeiite basalts. *Science*, *150*, 886–888.
- Taylor, R. N., Thirlwall, M. F., Murton, B. J., Hilton, D. R., & Gee, M. A. M. (1997). Isotopic constraints on the influence of the Icelandic plume. *Earth and Planetary Science Letters*, *148*, E1–E8. [https://doi.org/10.1016/S0012-821X\(97\)00038-1](https://doi.org/10.1016/S0012-821X(97)00038-1)
- Thompson, G., Bryan, W. B., Frey, F. A., Dickey, J. S. Jr., & Suen, C. J. (1976). Petrology and geochemistry of basalts from DSDP Leg 34, Nazca Plate. *Initial Reports of the Deep Sea Drilling Project*, *34*, 215–226.
- Thompson, G., Bryan, W. B., & Humphris, S. E. (1989). Axial volcanism on the East Pacific Rise, 10–12°N. In A. D. Saunders & M. J. Norry (Eds.), *Magmatism in the Ocean Basins, Geological Society, London, Special Publications* (Vol. 42).
- Thompson, R. N. (1980). Major-element chemistry of basaltic glasses in Hole 418A lavas and a dyke: Deep Sea Drilling Project Legs 52 and 53. *Initial Reports of the Deep Sea Drilling Project*, *51-53*, 973–976.
- Tian, L., Castillo, P. R., Hilton, D. R., Hawkins, J. W., Hanan, B. B., & Pietruszka, A. J. (2011). Major and trace element and Sr-Nd isotope signatures of the northern Lau Basin lavas: Implications for the composition and dynamics of the back-arc basin mantle. *Journal of Geophysical Research*, *116*, B11201. <https://doi.org/10.1029/2011JB008791>
- Timm, C., Hoernle, K., Werner, R., Hauff, F., van den Bogaard, P., Michael, P. J., et al. (2011). Age and geochemistry of the oceanic Manihiki Plateau, SW Pacific: New evidence for a plume origin. *Earth and Planetary Science Letters*, *304*, 135–146.
- Todd, E., Gill, J. B., Wysoczanski, R., Hergt, J. M., Wright, I. C., Leybourne, M. I., & Mortimer, N. (2011). Hf isotopic evidence for small-scale heterogeneity in the mode of mantle wedge enrichment: Southern Havre Trough and South Fiji Basin back-arc. *Geochemistry, Geophysics, Geosystems*, *12*, Q09011. <https://doi.org/10.1029/2011GC003683>
- Turner, S., Beier, C., Niu, Y., & Cook, C. (2011). U-Th-Ra disequilibria and the extent of off-axis volcanism across the East Pacific Rise at 90°30'N, 10°30'N, and 11°20'N. *Geochemistry, Geophysics, Geosystems*, *12*, Q0AC12. <https://doi.org/10.1029/2010GC003403>
- Ulrich, M., Hemond, C., Nonnotte, P., & Jochum, K. P. (2012). OIB/seamount recycling as a possible process for E-MORB genesis. *Geochemistry, Geophysics, Geosystems*, *13*, Q0AC19. <https://doi.org/10.1029/2012GC004078>
- Unni, C. K. (1976). Chlorine and bromine degassing during submarine and subaerial volcanism, (Thesis Or Dissertation, Doctoral). Uri; Monographic, 544.
- Vallier, T. L., Jenner, G. A., Frey, F. A., Gill, J. B., Davis, A. S., Volpe, A. M., et al. (1991). Subalkaline andesite from Valu Fa Ridge, a back-arc spreading center in southern Lau Basin: Petrogenesis, comparative chemistry, and tectonic implications. *Chemical Geology*, *91*, 227–256.
- van der Zwan, F. M., Devey, C. W., Augustin, N., Almeev, R., Bantan, R., & Basaham, A. (2015). Hydrothermal activity at the ultraslow- to slow-spreading Red Sea Rift traced by chlorine in basalt. *Chemical Geology*, *405*, 63–81. <https://doi.org/10.1016/j.chemgeo.2015.04.001>
- Verma, S. P. (1981). Seawater alteration effects on  $87\text{Sr}/86\text{Sr}$ , K, Rb, Cs, Ba and Sr in oceanic igneous rocks. *Chemical Geology*, *34*, 81–89.
- Verma, S. P. (1982). Magnetic properties and incompatible element geochemistry of some igneous rocks from Deep Sea Drilling Project Leg 64. *Initial Reports of the Deep Sea Drilling Project*, *64*, 667–671.
- Verma, S. P. (1983). Strontium and neodymium isotope geochemistry of igneous rocks from the North East Pacific and Gulf of California. *Chemical Geology*, *1*, 339–356.
- Verma, S. P., & Schilling, J.-G. (1982). Galapagos hot spot spreading center system 2.  $87\text{Sr}/86\text{Sr}$  and large ion lithophile element variations (85°W–101°W). *Journal of Geophysical Research*, *87*, 10,838–10,856. <https://doi.org/10.1029/JB087iB13p10838>
- Verma, S. P., Schilling, J.-G., & Waggoner, D. G. (1983). Neodymium isotopic evidence for Galapagos hotspot - spreading center system evolution. *Nature*, *306*, 654–657.
- Vidal, P., & Clauer, N. (1981). Pb and Sr isotopic systematics of some basalts and sulfides from the East Pacific Rise at 21°N (Project Rita). *Earth and Planetary Science Letters*, *55*, 237–246.
- Vils, F., Pelletier, L., Kalt, A., Muentener, O., & Ludwig, T. (2008). The lithium, boron and beryllium content of serpentinized peridotites from ODP Leg 209 (Sites 1272A and 1274A): Implications for lithium and boron budgets of oceanic lithosphere. *Geochimica et Cosmochimica Acta*, *72*, 5475–5504.
- Vils, F., Tonarini, S., Kalt, A., & Seitz, H.-M. (2009). Boron, lithium and strontium isotopes as tracers of seawater-serpentinite interaction at Mid-Atlantic Ridge, ODP Leg 209. *Earth and Planetary Science Letters*, *286*, 414–425.
- Vlastelic, I., Aslanian, D., Dosso, L., Bougault, H., Olivet, J. L., & Geli, L. (1999). Large-scale chemical and thermal division of the Pacific mantle. *Nature*, *399*, 345–350. <https://doi.org/10.1038/20664>
- Vlastelic, I., & Dosso, L. (2005). Initiation of a plume-ridge interaction in the South Pacific recorded by high-precision Pb isotopes along Hollister Ridge. *Geochemistry, Geophysics, Geosystems*, *6*, Q05011. <https://doi.org/10.1029/2004GC000902>

- Vlastelic, I., Dosso, L., Bougault, H., Aslanian, D., Geli, L., Etoubleau, J., et al. (2000a). Chemical systematics of an intermediate spreading ridge: The Pacific-Antarctic Ridge between 56°S and 66°S. *Journal of Geophysical Research*, *105*, 2915–2936. <https://doi.org/10.1029/1999JB900234>
- Vlastelic, I., Dosso, L., Bougault, H., Aslanian, D., Geli, L., Etoubleau, J., et al. (2000). Chemical systematics of an intermediate spreading ridge: The Pacific-Antarctic Ridge between 56°S and 66°S. *Journal of Geophysical Research*, *105*, 2915–2936. <https://doi.org/10.1029/1999JB900234>
- Vlastelic, I., Dosso, L., Guillou, H., Bougault, H., Geli, L., Etoubleau, J., & Joron, J.-L. (1998). Geochemistry of the Hollister Ridge: Relation with the Louisville hotspot and the Pacific-Antarctic Ridge. *Earth and Planetary Science Letters*, *160*, 777–793.
- Volker, F., Altherr, R., Jochum, K. P., & Mcculloch, M. T. (1997). Quaternary volcanic activity of the Southern Red Sea: New data and assessment of models on magma sources and afar plume-lithosphere interaction. *Tectonophysics*, *278*, 15–29.
- Volker, F., Mcculloch, M. T., & Altherr, R. (1993). Submarine basalts from the Red Sea: new Pb, Sr, and Nd isotopic data. *Geophysical Research Letters*, *20*, 927–930. <https://doi.org/10.1029/93GL00050>
- Volpe, A. M., & Goldstein, S. J. (1993). 226Ra-230Th disequilibrium in axial and off-axis Mid-Ocean Ridge basalts. *Geochimica et Cosmochimica Acta*, *57*, 1233–1241.
- Volpe, A. M., Macdougall, J. D., & Hawkins, J. W. (1987). Mariana Trough basalts (MTB): Trace element and Sr-Nd isotopic evidence for mixing between MORB-like and arc-like melts. *Earth and Planetary Science Letters*, *82*, 241–254.
- Volpe, A. M., Macdougall, J. D., & Hawkins, J. W. (1988). Lau Basin basalts (LBB): Trace element and Sr-Nd isotopic evidence for heterogeneity in backarc basin mantle. *Earth and Planetary Science Letters*, *90*, 174–186.
- Volpe, A. M., Macdougall, J. D., Lugmair, G. W., Hawkins, J. W., & Lonsdale, P. F. (1990). Fine-scale isotopic variation in Mariana Trough basalts: Evidence for heterogeneity and a recycled component in backarc basin mantle. *Earth and Planetary Science Letters*, *100*, 251–264.
- Waggoner, D. G. (1989). An isotopic and trace element study of mantle heterogeneity beneath the Norwegian-Greenland Sea, (Thesis Or Dissertation, Doctoral). Uri; Monographic.
- Wallace, P. (2002). Volatiles in submarine basaltic glasses from the northern Kerguelen Plateau (ODP Site 1140): Implications for source region compositions, magmatic processes, and plateau subsidence. *Journal of Petroleum*, *43*, 1311–1326.
- Wallace, P., & Carmichael, I. (1992). Sulfur in basaltic magmas. *Geochimica et Cosmochimica Acta*, *56*, 1863–1874.
- Wallace, P., & Roberge, J. (2004). Data report: Volatiles in submarine basaltic glasses from the Australian Antarctic Discordance (ODP Leg 187). In R. B. Pedersen, D. M. Christie, & D. J. Miller (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* (Vol. 187, pp. 1–8). College Station, Texas: Ocean Drilling Program.
- Wanless, V. D., Behn, M. D., Shaw, A. M., & Plank, T. (2014). Variations in melting dynamics and mantle compositions along the Eastern Volcanic Zone of the Gakkel Ridge: Insights from olivine-hosted melt inclusions. *Contributions to Mineralogy and Petrology*, *167*, 1005–1027. <https://doi.org/10.1007/s00410-014-1005-7>
- Wanless, V. D., Perfit, M. R., Klein, E. M., White, S. M., & Ridley, W. I. (2012). Reconciling geochemical and geophysical observations of magma supply and melt distribution at the 9°N overlapping spreading center, East Pacific Rise. *Geochemistry, Geophysics, Geosystems*, *13*, Q11005. <https://doi.org/10.1029/2012GC004168>
- Wanless, V. D., Perfit, M. R., Ridley, W. I., & Klein, E. M. (2010). Dacite petrogenesis on Mid-Ocean Ridges: Evidence for oceanic crustal melting and assimilation. *Journal of Petroleum*, *51*, 2377–2410.
- Wanless, V. D., Perfit, M. R., Ridley, W. I., Wallace, P., Grimes, C. B., & Klein, E. M. (2011). Volatile abundances and oxygen isotopes in basaltic to dacitic lavas on Mid-Ocean Ridges: The role of assimilation at spreading centers. *Chemical Geology*, *287*, 54–65.
- Wanless, V. D., & Shaw, A. M. (2012). Lower crustal crystallization and melt evolution at mid-ocean ridges. *Nature Geoscience*, *5*, 651–655. <https://doi.org/10.1038/ngeo1552>
- Wanless, V. D., Shaw, A. M., Behn, M. D., Soule, S. A., Escartin, J., & Hamelin, C. (2015). Magmatic plumbing at Lucky Strike volcano based on olivine-hosted melt inclusion compositions. *Geochemistry, Geophysics, Geosystems*, *16*, 126–147. <https://doi.org/10.1002/2014GC005517>
- Warren, J. M., Shimizu, N., Sakaguchi, C., Dick, H. J. B., & Nakamura, E. (2009). An assessment of upper mantle heterogeneity based on abyssal peridotite isotopic compositions. *Journal of Geophysical Research*, *114*, B12203. <https://doi.org/10.1029/2008JB006186>
- Waters, C., Sims, K. W. W., Perfit, M. R., Blichert-Toft, J., & Blusztajn, J. (2011). Perspective on the genesis of E-MORB from chemical and isotopic heterogeneity at 9°N East Pacific Rise. *Journal of Petroleum*, *52*, 565–602.
- Waters, C., Sims, K. W. W., Soule, S. A., Blichert-Toft, J., Dunbar, N., Plank, T., et al. (2013). Recent volcanic accretion at 9°-10°N East Pacific Rise as resolved by combined geochemical and geological observations. *Geochemistry, Geophysics, Geosystems*, *14*, 2547–2574. <https://doi.org/10.1002/ggge.20134>
- White, W. M. (1977). Geochemistry of igneous rocks from the central North Atlantic: The Azores and the Mid-Atlantic Ridge, (Thesis Or Dissertation, Doctoral). Uri; Monographic.
- White, W. M. (1979). Geochemistry of basalts from the FAMOUS area: A re-examination, *Carn Inst, Dtm, Yearbook*, *0*, 325–331.
- White, W. M. (1993). 238U/204Pb in MORB and open system evolution of the depleted mantle. *Earth and Planetary Science Letters*, *115*, 211–226.
- White, W. M., & Hofmann, A. W. (1982). Sr and Nd isotope geochemistry of oceanic basalts and mantle evolution. *Nature*, *296*, 821–825.
- White, W. M., Hofmann, A. W., & Puchelt, H. (1987). Isotope geochemistry of pacific Mid-Ocean Ridge basalts. *Journal of Geophysical Research*, *92*, 4881–4893. <https://doi.org/10.1029/JB092iB06p04881>
- White, W. M., & Schilling, J.-G. (1978). The nature and origin of geochemical variation in Mid-Atlantic Ridge basalts from the Central North Atlantic. *Geochimica et Cosmochimica Acta*, *42*, 1501–1516.
- Whitmarsh, R. B., Ross, D. A., Ali, S., Boudreaux, J. E., Coleman, R. G., Fleisher, R. L., et al. (1974). Site 226. *Initial Reports of the Deep Sea Drilling Project*, *23*, 595–600.
- Wilson, S., Murton, B. J., & Taylor, R. N. (2013). Mantle composition controls the development of an oceanic core complex. *Geochemistry, Geophysics, Geosystems*, *40*, 979–995. <https://doi.org/10.1002/ggge.20046>
- Yi, W., Halliday, A. N., Alt, J. C., Lee, D. C., Rehkaemper, M., Garcia, M. O., & Su, Y. (2000). Cadmium, indium, tin, tellurium, and sulfur in oceanic basalts: Implications for chalcophile element fractionation in the Earth. *Journal of Geophysical Research*, *105*, 18,927–18,948. <https://doi.org/10.1029/2000JB900152>
- Yonover, R. N. (1989). Petrological effects of rift failure at the Galapagos Spreading Center near 95.5°W including analyses of glass inclusions by laser mass spectrometry and ion microprobe, (Thesis or Dissertation, Doctoral). Univ Hawaii; Monographic.
- Yu, D., Fontignie, D., & Schilling, J.-G. (1997). Mantle plume - ridge interactions in the Central North Atlantic: A Nd isotope study of Mid-Atlantic Ridge basalts from 30°N to 50°N. *Earth and Planetary Science Letters*, *146*, 259–272.

- Zellmer, G. F., Rubin, K. H., Groenvold, K., & Jurado-Chichay, Z. (2008). On the recent bimodal magmatic processes and their rates in the Torfajökull–Veidivötn area, Iceland. *Earth and Planetary Science Letters*, *269*, 388–398.
- Zhang, G.-L., Zong, C.-L., Yin, X.-B., & Li, H. (2012). Geochemical constraints on a mixed pyroxenite-peridotite source for East Pacific Rise basalts. *Chemical Geology*, *330-331*, 176–187.
- Zhang, Y., Meng, F. X., & Niu, Y. L. (2016). Hf isotope systematics of seamounts near the East Pacific Rise (EPR) and geodynamic implications. *Lithos*, *262*, 107–119.
- Zindler, A., Staudigel, H., & Batiza, R. (1984). Isotope and trace element geochemistry of young Pacific seamounts: Implications for the scale of upper mantle heterogeneity. *Earth and Planetary Science Letters*, *70*, 175–195.